



LIBRARY
OF THE
UNIVERSITY
OF ILLINOIS

570.5
ILL
v. 5
COP. 52

~~NATURAL~~
HISTORY

JAN 6 - 1950



Digitized by the Internet Archive
in 2011 with funding from
University of Illinois Urbana-Champaign

ILLINOIS BIOLOGICAL MONOGRAPHS

PUBLISHED QUARTERLY
UNDER THE AUSPICES OF THE GRADUATE SCHOOL
BY THE UNIVERSITY OF ILLINOIS

VOLUME V

Urbana, Illinois
1919

EDITORIAL COMMITTEE

STEPHEN ALFRED FORBES

WILLIAM TRELEASE

HENRY BALDWIN WARD

570.5
ILL
v. 5
cop. 2

~~NATIONAL~~
~~HISTORICAL~~

TABLE OF CONTENTS

VOLUME V

NUMBERS	PAGES
1. The Skull of <i>Amiurus</i> . By James E. Kindred. With 8 plates.....	1-120
(Distributed July 7, 1919)	
2. Contributions to the Life Histories of <i>Gordius Robustus</i> Leidy and <i>Paragordius Varius</i> (Leidy). By Henry Gustav May. With 21 plates.....	121-238
3-4. Studies on Myxosporidia: a Synopsis of Genera and Species. By Rokusaburo Kudo. With 25 plates and 2 text figures.....	239-506
(Distributed December 31, 1920)	

ILLINOIS BIOLOGICAL MONOGRAPHS

Vol. V

July-October, 1919

Nos. 3 and 4

EDITORIAL COMMITTEE

STEPHEN ALFRED FORBES

WILLIAM TRELEASE

HENRY BALDWIN WARD

PUBLISHED UNDER THE
AUSPICES OF THE GRADUATE SCHOOL BY
THE UNIVERSITY OF ILLINOIS

COPYRIGHT, 1920 BY THE UNIVERSITY OF ILLINOIS
DISTRIBUTED DECEMBER 31, 1920.

STUDIES ON MYXOSPORIDIA

A SYNOPSIS OF GENERA AND SPECIES
OF MYXOSPORIDIA

WITH 25 PLATES AND 2 TEXTFIGURES

BY
ROKUSABURO KUDO

Contributions from the
Zoological Laboratory of the University of Illinois
No. 158

TABLE OF CONTENTS

Introduction.....	7
General Remarks on Recent Observations.....	8
Myxosporidia Recorded in the Present Paper (List I).....	9
Distribution of Myxosporidia.....	13
A. Geographical Distribution.....	13
B. Distribution of Myxosporidia in Animals.....	25
C. Distribution of Myxosporidia in the Organs of the Host.....	37
D. The Effect of Environment on the Organel Distribution of Myxosporidia in Hosts.....	44
The Spore.....	47
Definition of Terms Used for Descriptions.....	50
Classification of Myxosporidia.....	52
Descriptions of Genera and Species.....	60
Suborder Eurysporea.....	60
Family Ceratomyxidae.....	60
Genus Leptotheca.....	60
Genus Ceratomyxa.....	65
Genus Myxoproteus.....	81
Genus Wardia.....	82
Genus Mitraspora.....	84
Suborder Sphaerosporea.....	86
Family Chloromyxidae.....	87
Genus Chloromyxum.....	87
Family Sphaerosporidae.....	99
Genus Sphaerospora.....	100
Genus Sinuolinea.....	104
Suborder Platysporea.....	106
Family Myxidiidae.....	106
Genus Myxidium.....	107
Genus Sphaeromyxa.....	118
Genus Zschokkella.....	122
Family Myxosomatidae.....	123
Genus Myxosoma.....	123
Genus Lentospora.....	125
Family Myxobolidae.....	128
Genus Myxobolus.....	128
Genus Henneguya.....	158
Genus Hoferellus.....	173
Myxosporidia Genera et Species Incertae.....	174

Keys to the Genera and Species of Myxosporidia	178
Key to the Genera of Myxosporidia	178
Key to the Species	179
Summary	196
Appendix: New Myxosporidia from Australia	197
Bibliography	200
General Explanation of Figures	210
Explanation of Plates	212
Index	261

INTRODUCTION

Ten years have elapsed since Auerbach (1910) published *Die Cnidosporidien* in which he gave a synopsis of the genera and species of Myxosporidia known up to that time. During this period new genera and a number of new species have been added to the list of this particular group of parasitic protozoa from the various parts of the world. It is, therefore, desirable to have a complete monographic work including all the forms reported up to the present time.

The main objects of the present paper are: 1) to describe a new genus and a number of new species which have come under the observation of the writer; 2) to collect all the genera and species recorded by various authors; 3) to propose a new classification by which some of the confusion now existing may, probably, be avoided; 4) to show the geographical, zoological and organal distribution in the light of more recent observations; and 5) to present a complete list of the names of the hosts in which Myxosporidia occur.

The writer believes himself to be in possession of as complete references as possible under present conditions. However, he may be unaware of some works which have not reached him owing to the war.

The Myxosporidia recorded by Labbé (1899) are arranged in almost the same order as that author listed them, with some slight change such as placing the type species at the front of each genus or removing a few species to other genera, while those species which have been described since 1898 are arranged chronologically, no matter whether names are given the species or not.

Some of the references are omitted, especially when they can be found in Gurley (1894), Thélohan (1895), Labbé (1899), or Auerbach (1910). The description of each species is given according to the first observer. The observations of subsequent investigators are then mentioned in the second place.

Each species is described according to the following scheme:

- 1) Specific name
- 2) Synonyms and literature
- 3) Habitat, including the locality and the date of observation
- 4) Vegetative form
- 5) Spore
- 6) Remarks

I wish to express my appreciation to Professor Henry B. Ward whose kindness has made the completion of this paper possible.

GENERAL REMARKS ON RECENT OBSERVATIONS

The total number of species of Myxosporidia reported up to date and described in the following pages, excluding 12 ambiguous forms, reaches 237 of which 125 are species which have been observed since 1910.*

The distribution of these new forms is as follows:

Africa.....	6 species
Asia	23 species
Australia.....	1 species
Europe.....	31 species
North America.....	63 species
South America.....	1 species

Thus, the majority of the species were observed in other lands than Europe, nearly half being recorded from North American waters. It is not hard to anticipate from the observations made by Awerinzew, Davis, Kudo, Mavor, Johnston and Bancroft, and others, that further investigations on the parasites in the localities where the study of the protozoa under consideration was neglected, will bring out not only new and interesting forms which will be quite different from the comparatively well studied European species, but also many important facts that will clear unknown or doubtful phases concerning the life history and structure of Myxosporidia.

* Three species are included here which have been described (in Nipponese) by Miyairi in 1909.

MYXOSPORIDIA RECORDED IN THE PRESENT PAPER

LIST I

Order MYXOSPORIDIA Bütschli

I Suborder EURYSPOREA nom. nov. (see page 56)

I Family CERATOMYXIDAE Doflein

Genus 1 LEPTOTHECA Thélohan
[15 species]

- 1) *L. agilis* Thélohan (type species)
- 2) *L. elongata* Thélohan
- 3) *L. polymorpha* (Thél.) Labbé
- 4) *L. parva* Thélohan
- 5) *L. renicola* Thélohan
- 6) *L. hepseti* Thélohan
- 7) *L. perlata* (Gurley) Labbé
- 8) *L. sp.* Awerinzew
- 9) *L. macrospora* Auerbach
- 10) *L. informis* Auerbach
- 11) *L. longipes* Auerbach
- 12) *L. fusiformis* Davis
- 13) *L. scissura* Davis
- 14) *L. lobosa* Davis
- 15) *L. glomerosa* Davis

- 17) *C. sp.* Georgévitch
- 18) *C. coris* Georgévitch
- 19) *C. herouardi* Georgévitch
- 20) *C. mesospora* Davis
- 21) *C. sphairophora* Davis
- 22) *C. taenia* Davis
- 23) *C. attenuata* Davis
- 24) *C. recurvata* Davis
- 25) *C. lunata* Davis
- 26) *C. abbreviata* Davis
- 27) *C. flagellifera* Davis
- 28) *C. agglomerata* Davis
- 29) *C. amorphia* Davis
- 30) *C. monospora* Davis
- 31) *C. streptospora* Davis
- 32) *C. aggregata* Davis
- 33) *C. undulata* Davis
- 34) *C. navicularia* Davis
- 35) *C. spinosa* Davis

Genus 2 CERATOMYXA Thélohan
[35 species]

- 1) *C. arcuata* Thélohan (type species)
- 2) *C. sphaerulosa* Thélohan
- 3) *C. pallida* Thélohan
- 4) *C. globurifera* Thélohan
- 5) *C. appendiculata* Thélohan
- 6) *C. truncata* Thélohan
- 7) *C. reticularis* Thélohan
- 8) *C. inaequalis* Doflein
- 9) *C. linozpora* Doflein
- 10) *C. ramosa* Awerinzew
- 11) *C. drepanopsellae* Awerinzew
- 12) *C. tylosuri* Awerinzew
- 13) *C. (?) spari* Awerinzew
- 14) *C. sp. (?)* Awerinzew
- 15) *C. sp. (?)* Awerinzew
- 16) *C. acadensis* Mavor

Genus 3 MYXOPROTEUS Doflein
[3 species]

- 1) *M. ambiguus* (Thélohan) Doflein (type species)
- 2) *M. cordiformis* Davis
- 3) *M. cornutus* Davis

Genus 4 WARDIA nov. gen.
[2 species]

- 1) *W. ovinocua* nov. spec. (type species)
- 2) *W. ohlmacheri* (Gurley) Kudo

Genus 5 MITRASPORA Fujita emend.
Kudo [3 species]

- 1) *M. cyprini* Fujita (type species)
- 2) *M. caudata* (Parisi) Kudo
- 3) *M. elongata* nov. spec.

II Suborder SPHAEROSPOREA nom. nov. (see page 57)

I Family CHLOROMYXIDAE Thélohan

- Genus 1 CHLOROMYXUM Mingazzini [22 species]
- | | |
|------------------------------------------------|------------------------------------|
| 1) <i>C. leydigi</i> Mingazzini (type species) | 11) <i>C. sp.</i> Awerinzew |
| 2) <i>C. caudatum</i> Thélohan | 12) <i>C. thymalli</i> Lebzelter |
| 3) <i>C. quadratum</i> Thélohan | 13) <i>C. koi</i> Fujita |
| 4) <i>C. fluviatile</i> Thélohan | 14) <i>C. magnum</i> Awerinzew |
| 5) <i>C. mucronatum</i> Gurley | 15) <i>C. funduli</i> Hahn |
| 6) <i>C. diploxys</i> (Gurley) Thélohan | 16) <i>C. misgurni</i> Kudo |
| 7) <i>C. protei</i> Joseph | 17) <i>C. fujitai</i> Kudo |
| 8) <i>C. truttiae</i> Léger | 18) <i>C. clupeiidae</i> Hahn |
| 9) <i>C. cristatum</i> Léger | 19) <i>C. granulorum</i> Davis |
| 10) <i>C. dubium</i> Auerbach | 20) <i>C. trijugum</i> nov. spec. |
| | 21) <i>C. catostomi</i> nov. spec. |
| | 22) <i>C. wardi</i> nov. spec. |

II Family SPHAEROSPORIDAE Davis

- Genus 1 SPHAEROSPORA Thélohan [10 species]
- | | |
|------------------------------------------------|-------------------------------------------|
| 1) <i>S. divergens</i> Thélohan (type species) | 9) <i>S. (?) sp.</i> Southwell et Prashad |
| 2) <i>S. elegans</i> Thélohan | 10) <i>S. carassii</i> nov. spec. |

Genus 2 SINUOLINEA Davis [5 species]

- | | |
|---------------------------------|--------------------------------------------|
| 3) <i>S. rostrata</i> Thélohan | 1) <i>S. dimorpha</i> Davis (type species) |
| 4) <i>S. masovica</i> Cohn | 2) <i>S. capsularis</i> Davis |
| 5) <i>S. platessae</i> Woodcock | 3) <i>S. arborescens</i> Davis |
| 6) <i>S. angulata</i> Fujita | 4) <i>S. opacita</i> Davis |
| 7) <i>S. sp.</i> Davis | 5) <i>S. brachiophora</i> Davis |
| 8) <i>S. polymorpha</i> Davis | |

III Suborder PLATYSPOREA nom. nov. (see page 57)

I Family MYXIDIIDAE Thélohan

- Genus 1 MYXIDIUM Bütschli [26 species]
- | | |
|--------------------------------------------------|-----------------------------------------|
| 1) <i>M. lieberkühni</i> Bütschli (type species) | 21) <i>M. gadi</i> Georgévitch |
| 2) <i>M. incurvatum</i> Thélohan | 22) <i>M. glutinosum</i> Davis |
| 3) <i>M. sphaericum</i> Thélohan | 23) <i>M. phyllium</i> Davis |
| 4) <i>M. histophilum</i> Thélohan | 24) <i>M. striatum</i> Cunha et Fonseca |
| 5) <i>M. sp.</i> Gurley | 25) <i>M. kagayamai</i> nov. spec. |
| 6) <i>M. danilewskyi</i> Laveran | 26) <i>M. americanum</i> nov. spec. |
| 7) <i>M. giganteum</i> Doflein | |
| 8) <i>M. barbatulae</i> Cépède | |
| 9) <i>M. giardi</i> Cépède | |
| 10) <i>M. pfeifferi</i> Auerbach | |
| 11) <i>M. inflatum</i> Auerbach | |
| 12) <i>M. bergense</i> Auerbach | |
| 13) <i>M. procerum</i> Auerbach | |
| 14) <i>M. mackiei</i> Bosanquet | |
| 15) <i>M. macrocapsulare</i> Auerbach | |
| 16) <i>M. sp.</i> Awerinzew | |
| 17) <i>M. depressum</i> Parisi | |
| 18) <i>M. oviforme</i> Parisi | |
| 19) <i>M. anguillae</i> Ishii | |
| 20) <i>M. sp.</i> Mavor | |
- Genus 2 SPHAEROMYXA Thélohan [7 species]
- | |
|------------------------------------------------|
| 1) <i>S. balbianii</i> Thélohan (type species) |
| 2) <i>S. immersa</i> (Lutz) Thélohan |
| 3) <i>S. incurvata</i> Doflein |
| 4) <i>S. sabrazezi</i> Laveran et Mesnil |
| 5) <i>S. hellandi</i> Auerbach |
| 6) <i>S. exneri</i> Awerinzew |
| 7) <i>S. gasterostei</i> Georgévitch |
- Genus 3 ZSCHOKKELLA Auerbach [4 species]
- | |
|---------------------------------------------|
| 1) <i>Z. hildae</i> Auerbach (type species) |
| 2) <i>Z. nova</i> Klokacewa |
| 3) <i>Z. acheilognathi</i> Kudo |
| 4) <i>Z. globulosa</i> Davis |

II Family MYXOSOMATIDAE Poche

Genus 1 MYXOSOMA Thélohan
[3 species]

- 1) *M. dujardini* Thélohan (type species)
- 2) *M. (?) lobatum* Nemeček
- 3) *M. funduli* Kudo

Genus 2 LENTOSPORA Plehn
[6 species]

- 1) *L. cerebralis* (Hofer) Plehn (type species)
- 2) *L. multiplicata* Reuss
- 3) *L. encephalina* Mulsov
- 4) *L. asymmetrica* Parisi
- 5) *L. acuta* (Fujita) Kudo
- 6) *L. dermatobia* Ishii

III Family MYXOBOLIDAE Thélohan

Genus 1 MYXOBOLUS Bütschli
[63 species]

- 1) *M. mülleri* Bütschli (type species)
- 2) *M. piriformis* Thélohan
- 3) *M. unicusulatus* Gurley
- 4) *M. fuhrmanni* Auerbach
- 5) *M. oculi-leucisci* Trojan
- 6) *M. toyamai* Kudo
- 7) *M. notatus* Mavor
- 8) *M. sp.* Kudo
- 9) *M. rohita* Southwell et Prasad
- 10) *M. seni* Southwell et Prasad
- 11) *M. misgurni* nov. spec.
- 12) *M. pfeifferi* Thélohan
- 13) *M. inaequalis* Gurley
- 14) *M. dispar* Thélohan
- 15) *M. ellipsoides* Thélohan
- 16) *M. exiguus* Thélohan
- 17) *M. oviformis* Thélohan
- 18) *M. lintoni* Gurley
- 19) *M. globosus* Gurley
- 20) *M. oblongus* Gurley
- 21) *M. transovalis* Gurley
- 22) *M. obesus* Gurley
- 23) *M. cycloides* Gurley
- 24) *M. sphaeralis* Gurley
- 25) *M. anurus* Cohn
- 26) *M. sp.* Gurley
- 27) *M. sp.* Gurley
- 28) *M. sp.* Gurley
- 29) *M. cyprini* Doflein
- 30) *M. neurobius* Schuberg et Schröder
- 31) *M. aeglefini* Auerbach
- 32) *M. gigas* Auerbach
- 33) *M. volgensis* Reuss
- 34) *M. scardinii* Reuss
- 35) *M. physophilus* Reuss
- 36) *M. macrocapsularis* Reuss
- 37) *M. sandrae* Reuss
- 38) *M. bramae* Reuss

39) *M. cyprinicola* Reuss40) *M. balleri* Reuss41) *M. squamae* Keysselitz42) *M. cordis* Keysselitz43) *M. musculi* Keysselitz44) *M. sp.* Miyairi45) *M. sp.* Wegener46) *M. permagnus* Wegener47) *M. rotundus* Nemeček48) *M. minutus* Nemeček49) *M. sp.* Lebzelter50) *M. magnus* Awerinzew51) *M. carassii* Klokacewa52) *M. sp.* Southwell53) *M. funduli* Kudo54) *M. pleuronectidae* (Hahn)55) *M. capsulatus* Davis56) *M. nodularis* Southwell et Prasad57) *M. hylae* Johnston et Bancroft58) *M. aureatus* Ward59) *M. miyairii* nov. spec.60) *M. koi* nov. spec.61) *M. orbiculatus* nov. spec.62) *M. discrepans* nov. spec.63) *M. mesentericus* nov. spec.Genus 2 HENNEGUYA Thélohan
[32 species]

- 1) *H. psorospermica* Thélohan (type species)
- 2) *H. texta* (Cohn) Labbé
- 3) *H. minuta* (Cohn) Labbé
- 4) *H. oviperda* (Cohn) Labbé
- 5) *H. lobosa* (Cohn) Labbé
- 6) *H. peri-intestinalis* Cépède
- 7) *H. media* Thélohan
- 8) *H. brevis* Thélohan
- 9) *H. schizura* (Gurley) Labbé
- 10) *H. creplini* (Gurley) Labbé

- 11) *H. linearis* (Gurley) Labbé
- 12) *H. gurleyi* Kudo
- 13) *H. strongylura* (Gurley) Labbé
- 14) *H. monura* (Gurley) Labbé
- 15) *H. kolesnikovii* (Gurley) Labbé
- 16) *H. macrura* (Gurley) Thélohan
- 17) *H. zschokkei* (Gurley) Doflein
- 18) *H. sp.* (Gurley) Labbé
- 19) *H. sp.* (Gurley) Labbé
- 20) *H. tenuis* Vaney et Conte
- 21) *H. nüsslini* Schuberg et Schröder
- 22) *H. legeri* Cépède
- 23) *H. acerinae* Schröder
- 24) *H. gigantea* Nemeček
- 25) *H. (?) sp.* Nemeček
- 26) *H. gasterostei* Parisi
- 27) *H. neapolitana* Parisi
- 28) *H. wisconsinensis* Mavor
- 29) *H. brachyura* Ward
- 30) *H. salminicola* Ward

- 31) *H. miyairii* nov. spec.
- 32) *H. mictospora* nov. spec.

Genus 3 HOFERELLUS Berg
[1 species]

- 1) *H. cyprini* Doflein
- Appendix: Myxosporidia of unknown genera
and species [11 forms]
- 1) Gen. et spec. incert. Leydig
 - 2) Gen. et spec. incert. Leydig
 - 3) Gen. et spec. incert. Leydig
 - 4) Gen. et spec. incert. Heckel et Kner
 - 5) Gen. et spec. incert. Borne
 - 6) Gen. incert. *merlucii* Perugia
 - 7) Gen. incert. *congri* Perugia
 - 8) Gen. et spec. incert. Linton
 - 9) Gen. et spec. incert. Mingazzini
 - 10) Gen. et spec. incert. Nufer
 - 11) Gen. et spec. incert. Mavor
 - 12) Gen. et spec. incert. Mavor

DISTRIBUTION OF MYXOSPORIDIA

A. GEOGRAPHICAL DISTRIBUTION

As will be seen from List III, Myxosporidia are common parasites of fish in various parts of the world.

It is interesting to notice that the same species are found among fresh-water or marine fish from waters in widely separated countries. It is possible to think that Myxosporidia in marine fish may be carried into remote waters by the migration of their hosts, while those infecting fresh-water fish may be brought from one place to another by the transportation of infected fish for breeding purpose, etc. It should be noted in this connection that no intermediate host has yet been found in relation to myxosporidiosis.

The followings are the common species found in different localities:

<i>Leptotheca parva</i> Thél.	Marseille, Bergen
<i>Ceratomyxa sphaerulosa</i> Thél.	Monaco, Roscoff, Bergen
<i>C. appendiculata</i> Thél.	Roscoff, Marseille, Rovigno
<i>C. drepanopsettae</i> Awerinzew	Murman coast, Bergen, Woods Hole
<i>Chloromyxum leydigi</i> Ming.	Roscoff, Monaco, Napoli, Rovigno, Beaufort
<i>C. quadratum</i> Thél.	Roscoff, Marseille, Napoli, Beira
<i>Sphaerospora elegans</i> Thél.	Bretagne, Karlsruhe, Lago di Garda
<i>S. divergens</i> Thél.	Napoli, Roscoff, Smalfjorden
<i>Myxidium lieberkühni</i> Bütsch.	Lago Maggiore, France, Germany, Lake Mendota, Georgian Bay
<i>M. incurvatum</i> Thél.	Napoli, Monaco, Roscoff, Bergen, Beaufort
<i>M. bergense</i> Auerbach.	Bergen, St. Andrews
<i>M. oviforme</i> Parisi	Napoli, Norwegian coast
<i>Sphaeromyxa balbianii</i> Thél.	Roscoff, Napoli, Beaufort
<i>Myxosoma dujardini</i> Thél.	France, Germany, Tokio(?)

On the other hand, some species are limited to certain localities. Five species classified in the genus *Sinuolinea* by Davis are reported only from Beaufort, N. C., U. S. A. The two species of the genus *Wardia* have been found solely in the state of Illinois, U. S. A.

More detailed data are shown in the following list.

LIST II

ASIA

I NIPPON

Myxosporidia of fresh water fish

1) Northern Part (Hokkaido)

Sapporo: *Mitraspora cyprini* Fujita

Chloromyxum koi Fujita

Sphaerospora angulata Fujita

Lentospora acuta (Fujita) Kudo

2) Central part (Hondo)

Tokio: *Mitraspora cyprini* Fujita
Chloromyxum misgurni Kudo
Chloromyxum fujitai Kudo
Sphaerospora carassii nov. spec.
Myxidium kagayamai nov. spec.
Zschokkella acheilognathi Kudo
Myxosoma dujardini (?) Thélohan
Myxobolus toyamai Kudo
Myxobolus misgurni, nov. spec.
Myxobolus koi nov. spec.

Numazu: *Myxidium anguillae* Ishii
Lentospora dermatobia Ishii

3) Southern part (Kiushiu)

Fukuoka: *Myxobolus* sp. Miyairi
Myxobolus miyairii nov. spec.
Henneguya miyairii nov. spec.

II INDIA

A. Myxosporidia of fresh-water fish

Katwan, Mirzapore (U.P.): *Myxobolus* sp. Southwell
 Mirpur, Decca district: *Myxobolus rohita* Southwell et Prashad
Myxobolus seni Southwell et Prashad
Myxobolus nodularis Southwell et Prashad

B. Myxosporidian of reptiles

Bombay: *Myxidium mackiei* Bosaquet

III BURMA

In the vicinity of Ruby Mines: *Sphaerospora* sp. Southwell et Prashad

IV KAMTSCHATKA

?*Henneguya salminicola* Ward

AUSTRALIA

Myxosporidian of amphibia

In the vicinity of Sidney: *Myxobolus hylae* Johnston et Bancroft

AFRICA

A. Myxosporidia of fresh-water fish

Nile: *Myxobolus unicusulatus* Gurley
Henneguya strongylura Gurley

B. Myxosporidia of marine fish

1) Indian Ocean

Algoa Bay: *Chloromyxum magnum* Awerinzew
 Beira: *Chloromyxum quadratum* Thélohan
 East London: *Chloromyxum magnum* Awerinzew
 Lorenço Marques: *Ceratomyxa tylosuri* Awerinzew
Ceratomyxa spari Awerinzew
Ceratomyxa sp(?). Awerinzew
Ceratomyxa sp (?). Awerinzew
Sphaeromyxa exneri Awerinzew

2) South Atlantic Ocean

Lüderitz Bay: *Chloromyxum magnum* Awerinzew

NORTH AMERICA

I UNITED STATES

A. Myxosporidia of fresh-water fish

- 1) From Rivers emptying into Atlantic Ocean
 Carlus, Va. (tribt. of Potomac River): *Myxobolus transovalis* Gurley
 Columbia, S. C. (Santee River): *Myxobolus globosus* Gurley
 Kinston, N. C. (Neuse River): *Myxobolus globosus* Gurley
 West Falmouth, Mass.: *Myxobolus* sp. Kudo
 Woodbury, N. J. (Delaware River): *Henneguya monura* Gurley
- 2) From Lakes and Rivers opening into the Gulf of Mexico
 Fox River, trib. Mississippi: *Myxobolus globosus* Gurley
 Lake Mendota, Wis.: *Myxidium lieberkühni* Bütschli
Henneguya wisconsinensis Mavor et Strasser
 Neches River, Palestin, Tex.: *Henneguya macrura* (Gurley) Thélohan
 Storm Lake, Ia.: *Henneguya gurleyi* Kudo
 Stony Creek, Ill.: *Chloromyxum trijugum* nov. spec.
Myxobolus orbiculatus nov. spec.
Henneguya miclospora nov. spec.
 Homer Park, Ill.: *Chloromyxum trijugum* nov. spec.
Myxobolus orbiculatus nov. spec.
 Salt Fork, Urbana, Ill.: *Wardia ovinocua* nov. gen. nov. spec.
Chloromyxum catostomi nov. spec.
Myxobolus discrepans, nov. spec.
 Crystal Lake, Urbana, Ill.: *Mitraspora elongata* nov. spec.
Myxidium americanum nov. spec.
Myxobolus mesentericus nov. spec.
- 3) From the rivers opening into the Great Lakes
 Black River, Ohio: Gen. et spec. incert. Linton
 Put-In-Bay, Ohio: *Myxobolus aureatus* Ward
Henneguya brachyura Ward

B. Myxosporidia of marine fish (Atlantic Ocean)

- Beaufort, N. C.: *Leptotheca fusiformis* Davis
Leptotheca scissura Davis
Leptotheca lobosa Davis
Leptotheca glomerata Davis
Ceratomyxa mesospora Davis
Ceratomyxa sphairophora Davis
Ceratomyxa taenia Davis
Ceratomyxa attenuata Davis
Ceratomyxa recurvata Davis
Ceratomyxa lunata Davis
Ceratomyxa abbreviata Davis
Ceratomyxa flagellifera Davis
Ceratomyxa agglomerata Davis
Ceratomyxa amorpha Davis
Ceratomyxa monospora Davis
Ceratomyxa streptospora Davis
Ceratomyxa aggregata Davis
Ceratomyxa undulata Davis

- Ceratomyxa navicularia* Davis
Ceratomyxa spinosa Davis
Myxoproteus cordiformis Davis
Myxoproteus cornutus Davis
Chloromyxum leydigi Mingazzini
Chloromyxum granulosum Davis
Sphaerospora polymorpha Davis
Sinuolinea dimorpha Davis
Sinuolinea capsularis Davis
Sinuolinea arborescens Davis
Sinuolinea opacita Davis
Sinuolinea brachiophora Davis
Myxidium incurvatum Thélohan
Myxidium glutinosum Davis
Myxidium phyllium Davis
Sphaeromyxa balbianii Thélohan
Zschokkella globulosa Davis
Myxobolus capsulatus Davis
 Woods Hole, Mass: *Ceratomyxa drepanopectae* Awerinzew
Chloromyxum funduli Hahn
Chloromyxum clupeiidae Hahn
Myxosoma funduli Kudo
Myxobolus lintoni Gurley
Myxobolus funduli Kudo
Myxobolus pleuronectidae Hahn
 Locality unrecorded: *Henneguya schizura* (Gurley) Labbé
 C. Myxosporidian of Amphibia
 Sycamore, Ill.: *Wardia ohlmacheri* (Gurley) Kudo

II CANADA

- A. Myxosporidia of fresh-water fish
 Georgian Bay (south. part): *Myxidium lieberkühni* Bütschli
Myxobolus notatus Mavor
 Gen. et spec. incert. Mavor
 B. Myxosporidia of marine fish (Atlantic Ocean)
 Passamaquoddy Bay (at or
 near the mouth of St. Croix
 River), New Brunswick: *Ceratomyxa acadiensis* Mavor
Myxidium bergense Auerbach
M. sp. Mavor
 Gen. et spec. incert. Mavor

III ALASKA

- Klutina Lake: *Chloromyxum wardi* nov. spec.
 Stickeen River: *Henneguya salminicola* Ward

SOUTH AMERICA

- A. Myxosporidia of fresh-water fish from the waters connected with Atlantic Ocean
 Guiana: *Myxobolus inaequalis* Gurley
 Surinam: *Myxobolus inaequalis* Gurley
 Locality?: *Henneguya linearis* (Gurley) Labbé

B. Myxosporidian of marine fish (Atlantic Ocean)

Rio de Janeiro: *Myxidium striatum* Cunha et Fonseca

C. Myxosporidian of Amphibia

Brazil: *Sphaeromyxa immersa* (Lutz) Thélohan

EUROPE

I ITALY

A. Myxosporidia of fresh-water fish from lakes and rivers opening into Adriatic Sea

Lago di Como: *Mitraspora caudata* (Parisi) Kudo*Myxidium lieberkühni* BütschliLago di Garda: *Sphaerospora elegans* Thélohan*Henneguya gasterostei* ParisiLago di Varamo: *Henneguya minuta* (Cohn)Lago Maggiore: *Myxidium lieberkühni* BütschliMilano: *Myxidium lieberkühni* Bütschli*Myxobolus pfeifferi* ThélohanPavia: *Myxobolus gigas* Auerbach*Myxobolus ellipsoides* Thélohan*Henneguya peri-intestinalis* CépèdeTicino River: *Henneguya minuta* (Cohn)

B. Myxosporidia of marine fish

1) Ligurian Sea

Genova: *Chloromyxum leydigi* MingazziniGen. incert. *merluccii* PerugiaGen. incert. *congru* Perugia

2) Tyrrhenian Sea

Napoli: *Leptotheca agilis* Thélohan*Leptotheca elongata* Thélohan*Ceratomyxa arcuata* Thélohan*Ceratomyxa appendiculata* Thélohan*Ceratomyxa truncata* Thélohan*Ceratomyxa inaequalis* Doflein*Ceratomyxa linospora* Doflein*Myxoproteus ambiguus* (Thél.) Doflein*Chloromyxum leydigi* Mingazzini*Chloromyxum quadratum* Thélohan*Sphaerospora divergens* Thélohan*Myxidium incurvatum* Thélohan*Myxidium giganteum* Doflein*Myxidium depressum* Parisi*Myxidium oviforme* Parisi*Sphaeromyxa balbianii* Thélohan*Sphaeromyxa incurvata* Doflein*Sphaeromyxa sabrazesi* Laveran et Mesnil*Lentospora asymmetrica* Parisi*Myxobolus exiguus* Thélohan*Myxobolus mülleri* Bütschli*Henneguya neapolitana* Parisi

II MONACO

Myxosporidia of fish from Ligurian Sea

Leptotheca elongata Thélohan*Ceratomyxa sphaerulosa* Thélohan

Ceratomyxa arcuata Thélohan
Ceratomyxa pallida Thélohan
Ceratomyxa herouardi Georgévitch
Ceratomyxa sp. Georgévitch
Chloromyxum leydigi Mingazzini
Myxidium incurvatum Thélohan
Sphaeromyxa sabrazezi Laveran et Mesnil

III FRANCE

A. Myxosporidia of fresh-water fish

- 1) From Rivers opening into Atlantic Ocean
 - Aigne: *Myxobolus pfeifferi* Thélohan
 - Bretagne: *Sphaerospora elegans* Thélohan
 - Lorraine: *Myxobolus oviformis* Thélohan
 - Nancy: *Myxobolus pfeifferi* Thélohan
 - Marne: *Myxobolus pfeifferi* Thélohan
 - Seine: *Myxobolus pfeifferi* Thélohan
 - Paris: *Chloromyxum fluviatile* Thélohan
 - Wimereux: *Myxidium giardi* Cépède
- 2) From Rivers opening into Mediterranean Sea
 - Dauphiné: *Myxobolus mülleri* Bütschli
 - Drac River: *Myxobolus mülleri* Bütschli
Myxobolus pfeifferi Thélohan
 - Grenoble: *Chloromyxum cristatum* Léger
 - Isère River: *Myxidium barbatulae* Cépède
Myxobolus oviformis Thélohan
Myxobolus mülleri Bütschli
Myxobolus cycloides Gurley
Henneguya légeri Cépède
 - Lac d'Annecy: *Myxobolus mülleri* Bütschli
 - Lac de Paladru: *Myxobolus cycloides* Gurley
 - Lac du Bourget: *Myxobolus obesus* Gurley
Henneguya peri-intestinalis Cépède
 - Lyon?: *Henneguya tenuis* Vaney et Conte
 - Rhône River: *Myxobolus pfeifferi* Thélohan
 - Saône River: *Myxobolus pfeifferi* Thélohan

B. Myxosporidia of marine fish

- 1) From Atlantic Ocean
 - Arcachon *Sphaeromyxa sabrazezi* Laveran et Mesnil
 - Concarneau: *Ceratomyxa arcuata* Thélohan
Chloromyxum leydigi Mingazzini
Chloromyxum quadratum Thélohan
Sphaerospora divergens Thélohan
Myxidium incurvatum Thélohan
Sphaeromyxa balbianii Thélohan
 - Le Croisic: *Leptotheca elongata* Thélohan
Leptotheca parva Thélohan
Leptotheca renicola Thélohan
Ceratomyxa appendiculata Thélohan
Myxoproteus ambiguus (Thél.) Dofflein
Sphaerospora rostrata Thélohan

- Concarneau:** *Ceratomyxa arcuata* Thélohan
Chloromyxum leydigi Mingazzini
Chloromyxum quadratum Thélohan
Sphaerospora divergens Thélohan
Myxidium incurvatum Thélohan
Sphaeromyxa balbianii Thélohan
- Roscoff:** *Ceratomyxa sphaerulosa* Thélohan
Ceratomyxa arcuata Thélohan
Ceratomyxa appendiculata Thélohan
Chloromyxum leydigi Mingazzini
Chloromyxum quadratum Thélohan
Sphaerospora rostrata Thélohan
Sphaerospora divergens Thélohan
Myxidium incurvatum Thélohan
Myxidium gadi Georgévitch
Sphaeromyxa balbianii Thélohan
Sphaeromyxa sabraesi Laveran et Mesnil
Sphaeromyxa gasterostei Georgévitch
Myxobolus mülleri Bütschli
- Le-Vivier-sur-mer:** *Leptotheca parva* Thélohan
Myxidium sphaericum Thélohan
Myxobolus exiguus Thélohan
- St.-Valery-en-caux:** *Ceratomyxa sphaerulosa* Thélohan
- 2) From Mediterranean coast
- Marseille:** *Leptotheca elongata* Thélohan
Leptotheca parva Thélohan
Leptotheca renicola Thélohan
Leptotheca hepseli Thélohan
Ceratomyxa arcuata Thélohan
Ceratomyxa pallida Thélohan
Ceratomyxa globulifera Thélohan
Ceratomyxa appendiculata Thélohan
Ceratomyxa reticularis Thélohan
Chloromyxum leydigi Mingazzini
Sphaerospora rostrata Thélohan
Myxidium incurvatum Thélohan
Myxidium sphaericum Thélohan
Sphaeromyxa balbianii Thélohan
Myxobolus exiguus Thélohan
- Banyuls:** *Leptotheca elongata* Thélohan
Leptotheca polymorpha (Thél.) Labbé
Ceratomyxa arcuata Thélohan
Ceratomyxa globulifera Thélohan
Ceratomyxa appendiculata Thélohan
Ceratomyxa reticularis Thélohan
Chloromyxum leydigi Mingazzini
Sphaerospora rostrata Thélohan
Myxidium incurvatum Thélohan
Myxidium sphaericum Thélohan
Sphaeromyxa balbianii Thélohan
Myxobolus exiguus Thélohan

- Villefranche: *Ceratomyxa pallida* Thélohan
Ceratomyxa truncata Thélohan
Ceratomyxa coris Georgévitch
Sphaeromyxa balbianii Thélohan
- Locality unknown: *Leptotheca agilis* Thélohan
Leptotheca perlata (Gurley) Labbé
Myxidium lieberkühni Bütschli
Myxidium histophilum Thélohan
Myxosoma dujardini Thélohan
Myxobolus piriformis Thélohan
Myxobolus dispar Thélohan
Myxobolus obesus Thélohan
Henneguya psorospermica Thélohan
Henneguya media Thélohan
Henneguya brevis Thélohan
Hoferellus cyprini Doflein
C. Myxosporidian in a reptile
Myxidium danilewskyi Laveran

IV GERMANY

A. Myxosporidia of fresh-water fish

- 1) From Rivers opening into North Sea
- Throughout country: *Myxobolus cyprini* Doflein
- Berlin: *Henneguya oviperda* (Cohn)
- Bodensee: *Chloromyxum dubium* Auerbach
Myxobolus mülleri Bütschli
- Gutach: *Myxobolus neurobius* Schuberg et Schröder
Henneguya nüsslini Schub. et Schröder
- Karlsruhe and its vicinity: *Chloromyxum mucronatum* Gurley
Sphaerospora elegans Thélohan
Myxidium lieberkühni Bütschli
Myxidium pfeifferi Auerbach
Myxidium macrocapsulare Auerbach
Henneguya oviperda (Cohn)
Henneguya lobosa (Cohn)
Myxobolus gigas Auerbach
- Leipzig: *Myxobolus* sp. Gurley
- Mosel: *Myxobolus pfeifferi* Thélohan
Myxobolus squamae Keysselitz
Myxobolus cordis Keysselitz
Myxobolus musculi Keysselitz
- Neckar: *Myxobolus exiguus* Thélohan (Heidelberg)
Myxobolus mülleri Bütschli
Myxobolus pfeifferi Thélohan
Myxobolus squamae Keysselitz
Myxobolus cordis Keysselitz
Myxobolus musculi Keysselitz
Henneguya psorospermica Thélohan
Henneguya acerinae Schröder (Heidelberg)
- Rhine: *Myxidium lieberkühni* Bütschli
Myxobolus mülleri Bütschli
Henneguya psorospermica Thélohan
Lentospora encephalina Mulsow

2) From Rivers opening into Baltic Sea

- Alle: *Myxobolus mülleri* Bütschli
 Frisches Haff: *Myxidium lieberkühni* Bütschli
Myxosoma dujardini Thélohan
Myxobolus piriformis Thélohan
Myxobolus dispar Thélohan
Myxobolus exiguus Thélohan
Myxobolus oviformis Thélohan
Myxobolus mülleri Bütschli
Myxobolus cycloides Gurley
Myxobolus anurus Cohn
Myxobolus sp. Wegener
Myxobolus permagnus Wegener
Henneguya psorospermica Thélohan
Henneguya texta (Cohn)
Henneguya minuta (Cohn)
Henneguya lobosa (Cohn)
Henneguya creplini (Gurley)
 Kurisches Haff: *Myxosoma dujardini* Thélohan
Myxobolus exiguus Thélohan
Myxobolus oviformis Thélohan
Myxobolus cycloides Gurley
Henneguya psorospermica Thélohan
Henneguya texta (Cohn)
Henneguya creplini (Gurley) Labbé
 Masurische Seen: *Sphaerospora masovica* Cohn
Myxobolus dispar Thélohan
Myxobolus ellipsoides Thélohan
Myxobolus cycloides Gurley
Myxobolus anurus Cohn
Henneguya psorospermica Thélohan
Henneguya texta (Cohn)
 Pregel: *Myxosoma dujardini* Thélohan
Myxobolus piriformis Thélohan
Myxobolus dispar Thélohan
Myxobolus exiguus Thélohan
Myxobolus oviformis Thélohan
Myxobolus mülleri Bütschli
Myxobolus cycloides Gurley
Myxobolus anurus Cohn
Myxobolus permagnus Wegener
Henneguya psorospermica Thélohan
Henneguya texta (Cohn)
Henneguya minuta (Cohn)
Henneguya lobosa (Cohn)
Henneguya creplini (Gurley) Labbé
 Weichsel: *Myxobolus cyprini* Doëlein

3) Localities unknown:

- Chloromyxum leydigi* Mingazzini
Myxidium lieberkühni Bütschli
Myxidium sp. Gurley
Lentospora cerebralis (Hofer) Plehn

Henneguya schizura (Gurley) Labbé
Hojerellus cyprini Doflein
 Gen. et spec. incert. Leydig
 Gen. et spec. incert. Leydig
 Gen. et spec. incert. Leydig
 Gen. et spec. incert. Borne

V NETHERLAND

Myxosporidian of marine fish
 Helder: *Chloromyxum quadratum* Thélohan

VI ENGLAND

Myxosporidia of marine fish
 Firth of Clyde, More-
 camb, etc.: *Myxobolus aeglefini* Auerbach
 Liverpool (?): *Sphaerospora platessae* Woodcock

VII NORWAY

Myxosporidia of marine fish
 Abelvaer: *Myxidium bergense* Auerbach
Myxidium oviforme Parisi
Zschokkella hildae Auerbach
Myxobolus aeglefini Auerbach
 Bergen: *Leptotheca parva* Thélohan
Leptotheca macrospora Auerbach
Leptotheca informis Auerbach
Leptotheca longipes Auerbach
Ceratomyxa sphaerulosa Thélohan
Myxidium incurvatum Thélohan
Myxidium inflatum Auerbach
Myxidium bergense Auerbach
Myxidium procerum Auerbach
Sphaeromyxa hellandi Auerbach
Zschokkella hildae Auerbach
Myxobolus aeglefini Auerbach
 Bergsfjord: *Myxidium bergense* Auerbach
Zschokkella hildae Auerbach
 Boadsfjord: *Zschokkella hildae* Auerbach
 Bodö: *Myxidium bergense* Auerbach
Zschokkella hildae Auerbach
 Finkongkjeilen: *Myxidium bergense* Auerbach
Zschokkella hildae Auerbach
 Grönöy: *Myxidium bergense* Auerbach
Zschokkella hildae Auerbach
 Hammerfest: *Myxidium bergense* Auerbach
Myxidium oviforme Parisi
Zschokkella hildae Auerbach
 Harstad: *Myxidium bergense* Auerbach
Zschokkella hildae Auerbach
 Honnigsvaag: *Myxidium bergense* Auerbach
Zschokkella hildae Auerbach
 Kabelvaag: *Ceratomyxa drepanopsettae* Awerinzew

	<i>Myxidium bergense</i> Auerbach
	<i>Zschokkella hildae</i> Auerbach
Kiberg:	<i>Zschokkella hildae</i> Auerbach
Kirkenes:	<i>Myxidium bergense</i> Auerbach
Kristiana:	<i>Myxidium bergense</i> Auerbach
Kristiansand:	<i>Leptotheca parva</i> Thélohan
	<i>Leptotheca macrospora</i> Auerbach
	<i>Myxidium oviforme</i> Parisi
	<i>Zschokkella hildae</i> Auerbach
Lödingen:	<i>Myxobolus aeglefini</i> Auerbach
Makur:	<i>Myxidium bergense</i> Auerbach
	<i>Zschokkella hildae</i> Auerbach
Mosjöen:	<i>Zschokkella hildae</i> Auerbach
	<i>Myxidium bergense</i> Auerbach
Nusfjord:	<i>Myxidium bergense</i> Auerbach
	<i>Zschokkella hildae</i> Auerbach
Rörvik:	<i>Ceratomyxa drepanopectae</i> Awerinzew
	<i>Myxidium bergense</i> Auerbach
Rossfjord:	<i>Myxidium oviforme</i> Parisi
Skjervö:	<i>Zschokkella hildae</i> Auerbach
Skjöttningsberg:	<i>Zschokkella hildae</i> Auerbach
	<i>Myxidium bergense</i> Auerbach
Smalfjorden:	<i>Sphaerospora divergens</i> Thélohan
	<i>Zschokkella hildae</i> Auerbach
Stavanger:	<i>Leptotheca parva</i> Thélohan
	<i>Myxidium bergense</i> Auerbach
Svolvaer:	<i>Myxidium bergense</i> Auerbach
	<i>Zschokkella hildae</i> Auerbach
Tjömö:	<i>Leptotheca informis</i> Thélohan
	<i>Ceratomyxa drepanopectae</i> Awerinzew
	<i>Myxidium bergense</i> Auerbach
Torghatten:	<i>Sphaeromyxa hellandi</i> Auerbach
Trondhjem:	<i>Myxidium bergense</i> Auerbach
	<i>Myxidium oviforme</i> Parisi
	<i>Zschokkella hildae</i> Auerbach
Vikholmen:	<i>Zschokkella hildae</i> Auerbach
Vardö:	<i>Myxidium bergense</i> Auerbach
	<i>Myxidium oviforme</i> Parisi
	<i>Zschokkella hildae</i> Auerbach
	<i>Myxobolus aeglefini</i> Auerbach

VIII SWITZERLAND

Myxosporidia of fresh-water fish

1) From Lakes connected with North Sea	
Neuchatel:	<i>Myxobolus fuhrmanni</i> Auerbach
	<i>Myxobolus mülleri</i> Bütschli
	<i>Henneguya oviperda</i> (Cohn)
	<i>Henneguya zschokkei</i> (Gurley) Doflein
Thun:	<i>Henneguya zschokkei</i> (Gurley) Doflein
Zurich:	<i>Henneguya zschokkei</i> (Gurley) Doflein
Lucerne:	<i>Myxosoma dujardini</i> Thélohan
	<i>Myxobolus ellipsoides</i> Thélohan

- Myxobolus oviformis* Thélohan
Myxobolus mülleri Bütschli
Henneguya psorospermica Thélohan
Henneguya texta (Cohn)
Henneguya zschokkei (Gurley) Doflein
 Gen. et spec. incert. Nufer
 Wallen: *Henneguya zschokkei* (Gurley) Doflein
 2) From Lake connected with Mediterranean Sea
 Geneva: *Myxobolus sphaeralis* Gurley
Henneguya zschokkei (Gurley) Doflein

IX AUSTRIA

- A. Myxosporidia of fresh-water fish
 1) From Rivers opening into Black Sea
 Danube tributaries and Neusiedler: *Chloromyxum thymalli* Lebzelter
Myxosoma (?) *lobatum* Nemeček
Myxobolus aeglefini Auerbach
Myxobolus cyprini Doflein
Myxobolus rotundus Nemeček
Myxobolus minutus Nemeček
Myxobolus sp. Lebzelter
Henneguya acerinae Schröder
Henneguya gigantea Nemeček
 2) From Rivers opening into North Sea
 Prag: *Myxosoma dujardini* Thélohan
Myxobolus ellipsoides Thélohan
Myxobolus oculi-leucisci Trojan
 Krakau: *Myxobolus cyprini* Doflein
 B. Myxosporidia of marine fish (Adriatic Sea)
 Rovigno: *Leptotheca agilis* Thélohan
Ceratomyxa pallida Thélohan
Ceratomyxa appendiculata Thélohan
Myxoproteus ambiguus (Thél.) Doflein
Chloromyxum leydigi Mingazzini
Sphaeromyxa sabrazesi Laveran et Mesnil
 Locality unknown: Gen. et spec. incert. Heckel et Kner
 C. Myxosporidian of Amphibia
 Vienna: *Chloromyxum protei* Joseph

X SERBIA

- Pergrad (Danube): *Henneguya gigantea* Nemeček

XI RUSSIA

- A. Myxosporidia of fresh-water fish
 Volga (to Caspian Sea): *Lentospora multiplicata* Reuss
Myxobolus volgensis Reuss
Myxobolus scardinii Reuss
Myxobolus physophilus Reuss
Myxobolus macrocapsularis Reuss
Myxobolus sandrae Reuss
Myxobolus bramae Reuss
Myxobolus cyprinicola Reuss
Myxobolus balleri Reuss

- Don (to Black Sea): *Myxobolus* sp. Gurley
 Locality unknown: *Zschokkella nova* Klokacewa
Myxobolus magnus Awerinzew
Myxobolus carassii Klokacewa
Henneguya kolesnikovi (Gurley) Labbé

B. Myxosporidia of marine fish from Arctic Ocean

- Murman coast: *Ceratomyxa ramosa* Awerinzew
Ceratomyxa drepanopseltae Awerinzew
Myxidium sp. Awerinzew
Leptotheca sp. Awerinzew
Chloromyxum sp. Awerinzew

B. DISTRIBUTION OF MYXOSPORIDIA IN ANIMALS

The number of host species that harbor Myxosporidia is 237, as will be seen from List III.

The two incompletely studied forms are found in Annelida and Insecta, Myxosporidia are the parasites of Vertebrata, especially of Pisces, only few being found infecting Amphibia and Reptilia. They are distributed among these groups of animals as follows:

	Number of host species
Annelida	1
Insecta	1
Pisces	223
Amphibia.....	8
Reptilia	4

Gurley (1894:101-105), Wasielewsky (1896:132-148), Labbé (1899:133-161) and Auerbach (1910:36-45; 1911:471-494) gave lists in which they recorded the names of host species. Wasielewsky arranged the names alphabetically while others listed them according to their systematic order. In the following pages, the writer followed Wasielewsky, i.e., the names of the host species are arranged alphabetically as is supposed to be more convenient in referring to the host than any form presented otherwise.

LIST III. LIST OF HOST SPECIES

Host	Organ Infected	Myxosporidian	Locality
Annelida <i>Nais lacustris</i> (<i>N. proboscidea</i>).....	Unknown	<i>Myxobolus</i> sp.	Germany
Insecta <i>Tortrix viridana</i> L. (imago)	Abdominal cavity	<i>Chloromyxum diploxys</i>	France
Pisces <i>Abramis ballerus</i> L.....	Branchiae	<i>Myxobolus balleri</i>	Russia
<i>A. brama</i> L.....	“	<i>bramae</i>	“

Host	Organ Infected	Myxosporidian	Locality
	Branchiae	<i>Myxobolus cycloides</i>	Germany
	"	<i>ellipsoides</i>	" (?)
	"	<i>exiguus</i>	France
	"	<i>oviformis</i>	France
	"	<i>rotundus</i>	Austria
	Gall-bladder	<i>Sphaerospora masovica</i>	Germany
	Kidney	<i>Myxobolus cyprini</i>	" , Hungary
	Subcut. conn. tissue of oper- culum	<i>gigas</i>	Germany, Italy
<i>A. vimba</i> L.....	Branchiae	<i>cycloides</i>	Germany
	"	<i>ellipsoides</i>	" (?)
	"	<i>oviformis</i>	Germany
<i>Acanthias acanthias</i> L.....	Gall-bladder	<i>Chloromyxum leydigi</i>	France
<i>A. blainvillei</i>	"	<i>magnum</i>	Africa
<i>Acerina cernua</i> L.....	Branchiae	<i>Henneguya acerinae</i>	Germany
	" , Muscle	<i>creplini</i>	"
	Conn. tissue of aliment. canal	<i>tenuis</i>	France
	Eye	<i>Myxobolus magnus</i>	Russia
	Muscle	<i>Leptotheca perlata</i>	France?
<i>Acheilognathus lanceolatum</i> Temm. et Schl.....	Gall-bladder, Gall-duct	<i>Zschokkella acheilognathi</i>	Nippon
<i>Alburnus alburnus</i> L..... (<i>A. lucidus</i> Heck)	Branchiae	<i>Myxobolus cycloides</i>	Germany
	"	<i>dispar</i>	"
	"	<i>ellipsoides</i>	Germany ?
	" , kidney	<i>oviformis</i>	France
	Muscle and spleen	<i>obesus</i>	"
	Eye	<i>dispar</i>	"
		<i>mülleri</i>	Switzerland
<i>Alosa finta</i> Cuv. var. <i>lacustris</i> Fatio.....	Kidney	<i>Mitraspora caudata</i>	Italy
<i>Ameiurus melas</i> Raf.....	Base of spines of 2nd dorsal fin	<i>Henneguya gurleyi</i>	U. S. A.
<i>Ancylosetta quadrocellata</i> Gill.....	Gall-bladder	<i>Ceratomyxa undulata</i>	"
<i>Anguilla japonica</i> Temm. et Schl.....	Integument	<i>Myxidium anguillae</i>	Nippon
	"	<i>Lentospora dermatobia</i>	"
<i>A. vulgaris</i> Flemm.....	Kidney	<i>Myxidium giardi</i>	France
<i>Aphredoderus sayanus</i> Gill.	Subcutaneous intermusc. tiss.	<i>Henneguya monura</i>	U. S. A.
<i>Apogon rex-mullorum</i> Cuv.	Gall-bladder	<i>Myxidium oviforme</i>	Italy
<i>Argentina silus</i> Nilss.....	"	<i>procerum</i>	Norway
<i>Ariodes polystaphylodon</i>	Muscle	<i>Chloromyxum quadratum</i>	Africa
<i>Aspius rapax</i> Ag.....	Branchiae	<i>Myxosoma(?) lobatum</i>	Austria
<i>Aspro asper</i> L.....	"	<i>Myxobolus mülleri</i>	France
<i>A. zingel</i> Cuv.....	"	<i>Henneguya acerinae</i>	"

Host	Organ Infected	Myxosporidian	Locality	
<i>Atherina hepsetus</i> L.....	Gall-bladder	<i>Leptotheca hepseti</i>	France	
<i>Bairdiella chrysura</i>	Urin. bladder	<i>Myxoproteus cornutus</i>	U. S. A.	
<i>B. rouchus</i> Cuv. et Val.....	Gall-bladder	<i>Myxidium striatum</i>	Brazil	
<i>Barbus barbatus</i> L.....	Kidney, spleen,	<i>Myxobolus pfeifferi</i>	France, Germany	
(<i>B. fluviatilis</i>)	intestine, ovary, etc.			
	Inner surface of scale			<i>squamae</i>
	Muscle of ven- tricle			<i>cordis</i>
	Muscle, kidney	<i>musculi</i>	"	
<i>B. plebejus</i> Val.....	"	<i>pfeifferi</i>	Italy	
<i>B. vulgaris</i> Flem.....	Branchiae	<i>mülleri</i>	Germany	
<i>Barilius barna</i>	Under scales	<i>Sphaerospora</i> sp.	Burma	
<i>Belone acus</i> Risso.....	Gall-bladder	<i>Myxidium sphaericum</i>	France	
<i>B. belone</i> L.....	"	<i>Myxidium sphaericum</i>	"	
<i>Blennius ocellatus</i>	"	<i>Sphaeromyxa incurvata</i>	Italy	
<i>B. gattorugine</i> Brunn.....	Kidney	<i>Chloromyxum quadratum</i>	"	
<i>B. pholis</i> L.....	Gall-bladder	<i>Myxidium incurvatum</i>	France	
	Renal tubules	<i>Sphaerospora divergens</i>	"	
<i>Blicca björkna</i> L.....	Branchiae	<i>Myxobolus cycloides</i>	Germany	
	"	<i>ellipsoides</i>	" ?	
	"	<i>macrocapsularis</i>	Russia	
	"	<i>oviformis</i>	Germany	
<i>Box boops</i> L.....	Gall-bladder	<i>Ceratomyxa pallida</i>	France, Italy	
<i>B. salpa</i> L.....	Gall-bladder	<i>Ceratomyxa herouardi</i>	Monaco	
	"	<i>pallida</i>	France, Italy	
	Kidney	<i>Henneguya neapolitana</i>	Italy	
<i>Brevoortia tyrannus</i> Latr....	Muscle	<i>Chloromyxum clupeiidae</i>	U. S. A.	
<i>Brosimius brosmo</i> Ascanius..	Gall-bladder	<i>Leptotheca longipes</i>	Norway	
	"	<i>Sphaeromyxa hellandi</i>	"	
<i>Callionymus lyra</i> L.....	"	<i>Myxidium incurvatum</i>	France, Norway	
	Muscle	<i>Chloromyxum quadratum</i>	France	
<i>Carassius auratus</i> L.....	Branchiae	<i>Lenospora acuta</i>	Nippon	
	Kidney	<i>Sphaerospora angulata</i>	"	
	"	<i>Mitraspora cyprini</i>	"	
	Subcutaneous tiss. of head	<i>Henneguya miyairii</i>	"	
(<i>C. carassius</i> L.).....	Body cavity	<i>Myxobolus</i> sp.	Germany	
	Branchiae	<i>dispar</i>	"	
	"	<i>Sphaerospora carassii</i>	Nippon	
(<i>C. vulgaris</i> L.).....	Body cavity, liver, intestine	<i>Myxobolus carassii</i>	Russia	
<i>Carcharhinus limbatus</i>	Gall-bladder	<i>Zschokkella nova</i>	"	
<i>C. sp.</i>	"	<i>Chloromyxum leydigi</i>	U. S. A.	
	"	<i>Ceratomyxa flagellifera</i>	"	

Host	Organ Infected	Myxosporidian	Locality
<i>Carpiodes difformis</i>	Branchiae	<i>Myxobolus discrepans</i>	U. S. A.
<i>Catostomus commersonii</i> Lac.....	Gall-bladder	<i>Chloromyxum catostomi</i>	"
<i>Centronotus gunellus</i>	"	<i>Sphaeromyxa hellandi</i>	Norway
<i>Cepola rubescens</i> L.....	"	<i>balbianii</i>	France
<i>Cestracion tiburo</i>	"	<i>Chloromyxum leydigi</i>	U. S. A.
<i>C. zygaena</i>	"	<i>Ceratomyxa mesospora</i>	"
	"	<i>recurvata</i>	"
	"	<i>Chloromyxum leydigi</i>	"
	"	<i>Leptotheca fusiformis</i>	"
<i>Chaetodipterus faber</i>	Gall-bladder	<i>Ceratomyxa streptospora</i>	"
	Urin. bladder	<i>Myxoproteus cordiformis</i>	"
<i>Chondrostoma nasus</i> L.....	Branchiae	<i>Myxobolus exiguus</i>	Germany
	"	Gen. et spec. incert.	Switzerland
	Tongue	Gen. et spec. incert.	?
<i>Citharus linguata</i> Gthr.....	Gall-bladder	<i>Myxidium depressum</i>	Italy
<i>Clupea harengus</i> Young.....	"	<i>Ceratomyxa sphaerulosa</i>	Norway
	Muscle	<i>Chloromyxum clupeidae</i>	U. S. A.
<i>C. pilchardus</i> Walb. (<i>Alosa sardina</i>).....	Gall-bladder	<i>Ceratomyxa truncata</i>	France, Italy
	"	<i>Sphaeromyxa balbianii</i>	Italy
<i>Cobitis barbatula</i> L.....	Kidney	<i>Myxidium barbatulae</i>	France
	Urin. bladder	<i>Henneguya legeri</i>	"
<i>C. fossilis</i> L.....	Branchiae,		
<i>Conger conger</i> L. (<i>Leptocephalus c.</i>).....	kidney, spleen	<i>Myxobolus piriformis</i>	Germany
<i>Coregonus lavaretus</i> L. (<i>C. fera</i>).....	Gall-bladder	Gen. incert. <i>congru</i>	Italy
	Branchiae	<i>Henneguya</i> sp.	France?
	" (mucosa)	<i>Myxobolus sphaeralis</i>	Switzerland
	Muscle	<i>Henneguya kolesnikovi</i>	Russia
	"	<i>zschokkei</i>	Switzerland
<i>C. exiguus albellus</i>	" , branchia	"	"
<i>C. wartmanni nobilis</i>	" "	"	"
<i>Coris giofredi</i> Risso.....	Gall-bladder	<i>Ceratomyxa coris</i>	France
<i>C. julis</i> L.....	"	"	"
	Muscle	<i>Chloromyxum quadratum</i>	"
	Gall-bladder	<i>Myxidium oviforme</i>	Italy
<i>Cottus gobis</i> L.....	Branchiae	<i>Myxobolus mülleri</i>	France
<i>C. scorpius</i>	"	<i>Myxidium</i> sp.	Russia
<i>Crenilabrus mediterraneus</i> ..	"	<i>Ceratomyxa inaequalis</i>	Italy
<i>C. melops</i> L.....	Eye	<i>Myxobolus mülleri</i>	Germany
	"		France
	Gall-bladder	<i>Ceratomyxa arcuata</i>	France
	Kidney	<i>Sphaerospora divergens</i>	"
<i>C. pavo</i> Cuv. et Val.....	Gall-bladder	<i>Ceratomyxa inaequalis</i>	Italy
	Kidney	<i>Lentospora asymmetrica</i>	Italy
	"	<i>Sphaerospora divergens</i>	France, Italy
<i>Cyclopterus lumpus</i> L.....	Gall-bladder	<i>Myxidium inflatum</i>	Norway

Host	Organ Infected	Myxosporidian	Locality
<i>Cynoscion regalis</i>	Gall-bladder Urin. bladder, ureters	<i>Myxidium glutinosum</i> <i>Sinuolinea dimorpha</i>	U. S. A. “
<i>Cyprinodon variegatus</i>	Subcutaneous tissue Visceral conn. tissue	<i>Myxobolus lintoni</i> <i>capsulatus</i>	“ “
<i>Cyprinus carpio</i> L.....	Branchiae “ “ “ “ Brain Gall-bladder Kidney “ liver, spl. Kidney “	<i>cyprinicola</i> <i>dispar</i> <i>oviformis</i> <i>toyamai</i> <i>Myxosoma dujardini</i> <i>Myxobolus koi</i> <i>Lentospora encephalina</i> <i>Chloromyxum koi</i> <i>Hoferellus cyprini</i> <i>Myxobolus cyprini</i> <i>Mitraspora cyprini</i> <i>Sphaerospora angulata</i>	Russia France, Germany France ? Nippon France Nippon Germany Nippon Germany, France Germany, Hungary Nippon “
<i>C. (Rasbora) daniconius</i> Day.....	Subcutaneous intermuscular tissue Muscles	<i>Myxobolus</i> sp. <i>Myxobolus nodularis</i>	India India
<i>Dasyatis hastatus</i>	Gall-bladder “	<i>Chloromyxum leydigi</i> <i>Leptotheca scissura</i>	U. S. A. “
<i>D. sabina</i>	“	<i>Chloromyxum leydigi</i>	“
<i>Drepanopsetta platessoides</i> Fabr.....	“	<i>Ceratomyxa drepanopsettae</i>	Russia
<i>Erimyzon sucetta oblongus</i> Lac (<i>Catostomus</i> <i>tuberculatus</i>)	Branchiae Integument	<i>Myxobolus globosus</i> <i>oblongus</i>	U. S. A. “
<i>Esox lucius</i> L.....	Branchiae “ “ Intestinal wall Eye muscle, etc. Ovary Urin. bladder	<i>anurus</i> <i>Henneguya lobosa</i> <i>Henneguya psorospermica</i> <i>Henneguya peri-intestinalis</i> <i>Henneguya schizura</i> <i>Henneguya ovi-perda</i> <i>Myxidium lieberkühni</i>	Germany “ France, Germany France, Italy U. S. A. Germany, Switzerland France, Ita- ly, Canada, U. S. A., Germany
<i>Fundulus</i> sp.....	Muscle	<i>Chloromyxum funduli</i>	U. S. A.

Host	Organ Infected	Myxosporidian	Locality
<i>F. diaphanus</i>	Muscle	<i>Myxobolus funduli</i>	U. S. A.
<i>F. heteroclitus</i>	Branchiae, Muscle	<i>Myxobolus funduli</i>	"
	Branchiae	<i>Myxosoma funduli</i>	"
<i>F. majalis</i>	"	<i>Myxosoma funduli</i>	"
	" , muscle	<i>Myxobolus funduli</i>	"
	Gall-bladder	<i>Myxidium incurvatum</i>	"
<i>Gadus aeglefinis</i> L.....	Cartilage	<i>Myxobolus aeglefini</i>	Germany
	Urin. bladder	<i>Zschokkella hildae</i>	Norway
<i>G. callarias</i> L.....	Cartilage	<i>Myxobolus aeglefini</i>	Germany
	Gall-bladder	<i>Myxidium oviforme</i>	Norway
	Urin. bladder	<i>Zschokkella hildae</i>	Norway
<i>G. esmarkii</i> Nilss.....	Cartilage, bone of cranium, eye	<i>Myxobolus aeglefini</i>	Germany, England
<i>G. merlangus</i> L.....	Gall-bladder	<i>Leptotheca informis</i>	Norway
	Cartilage	<i>Myxobolus aeglefini</i>	Germany
<i>G. morrhua</i> L.....	Cartilage	<i>Myxobolus aeglefini</i>	Germany
<i>G. pollachius</i> L.....	Gall-bladder	<i>Myxidium gadi</i>	France
<i>G. virens</i> L.....	"	<i>bergense</i>	Norway
	Urin. bladder	<i>Zschokkella hildae</i>	"
<i>Galeocerda tigrinus</i>	Gall-bladder	<i>Ceratomyxa lunata</i>	U. S. A.
<i>Galeus galeus</i> L. (<i>G. canis</i>)..	"	<i>sphaerulosa</i>	France
<i>Gambusia affinis</i>	"	<i>Myxidium incurvatum</i>	U. S. A.
	"	<i>phyllium</i>	"
<i>Gasterosteus aculeatus</i>	Kidney (r. t.), ovary	<i>Henneguya media</i>	France
	"	<i>brevis</i>	"
	"	<i>Sphaerospora elegans</i>	" , Italy
	Kidney	<i>Henneguya gasterostei</i>	Italy
<i>G. pungilius</i> L.....	" (r. t.), ovary	<i>brevis</i>	France
	"	<i>media</i>	"
	"	<i>Sphaerospora elegans</i>	"
<i>G. spinachia</i>	Gall-bladder	<i>Sphaeromyxa gasterostei</i>	"
<i>Gobio gobio</i> L. (<i>G. fluviatilis</i>).....	Fin	<i>Myxobolus mülleri</i>	Germany
	Fin, spleen, kidney, liver	<i>oviformis</i>	France
	Branchiae	<i>cycloides</i>	Germany
<i>Gobius fluviatilis</i> L.....	Body-cavity	Gen. et. spec. incert.	"
<i>Gobius paganellus</i> L.....	Gall-bladder	<i>Ceratomyxa arcuata</i>	Italy
<i>Heliases chromis</i> Gthr.....	"	<i>Ceratomyxa arcuata</i>	Italy, Monaco
<i>Hippocampus brevirostris</i> Cuv.....	"	<i>Myxidium incurvatum</i>	Italy
	"	<i>Sphaeromyxa sabrazesi</i>	France, Italy, Hungary

Host	Organ Infected	Myxosporidian	Locality
<i>H. guttulatus</i> Cuv.....	Gall-bladder	<i>Sphaeromyxa sabrazezi</i>	Fr., Hun., Monaco
<i>Hippoglossoides liman-</i> <i>doides</i>	Urin. bladder	<i>Sphaerospora divergens</i>	Norway
<i>Hippoglossus vulgaris</i> Flemm.....	Gall-bladder	<i>Ceratomyxa drepanopsettae</i>	Norway
	"	<i>ramosa</i>	Russia
<i>Hybognathus nuchalis</i> Ag....	Conn. tissue of the head	<i>Henneguya macrura</i>	U. S. A.
<i>Idus melanotus</i> Heck.....	Branchiae	<i>Myxobolus ellipsoides</i>	Germany?
	Muscle	<i>Lentospora multiplicata</i>	Russia
<i>Labeo rohita</i>	Branchiae	<i>Myxobolus rohitae</i>	India
	Fins	<i>Myxobolus seni</i>	"
<i>Labeo niloticus</i> Forsk.....	?	<i>Myxobolus unicipsulatus</i>	Egypt
<i>Labrus turdus</i>	Gall-bladder	<i>Ceratomyxa linozpora</i>	Italy
<i>Leiostomus xanthurus</i>	"	<i>aggregata</i>	U. S. A.
<i>Lepistosteus platystomus</i>	Urinary bladder	<i>Sphaerospora</i> sp.	"
<i>Lepomis cyanellus</i> Raf.....	Mesentery	<i>Myxobolus mesentericus</i>	"
	Urin. bladder	<i>Henneguya mictospora</i>	"
	Kidney	<i>Mitraspora elongata</i>	"
<i>L. humilis</i> Girard.....	Ovary	<i>Wardia ovinocua</i>	"
	Urinary bladd.	<i>Henneguya mictospora</i>	"
<i>L. megalotis</i> Raf.....	Gall-bladder	<i>Chloromyxum trijugum</i>	"
<i>Leuciscus cephalus</i> (<i>Squalis</i> <i>cephalus</i>).....	Air-bladder,		France,
	branchiae	<i>Myxobolus mülleri</i>	Germany
	Branchiae	<i>ellipsoides</i>	" "
	Gall-bladder	<i>Chloromyxum fluviatile</i>	France
<i>L. lucius</i> L.....	Branchiae	<i>Myxobolus</i> sp.	Germany?
<i>L. phoxinus</i> L. (<i>Phoxinus</i> <i>laevis</i> Ag.).....	Conn. tiss. of kidney; ovary	<i>Myxidium histophilum</i>	France
	"	<i>Myxobolus mülleri</i>	"
<i>L. rutilus</i> L.....	Branchiae	<i>ellipsoides</i>	Germany?
	"	<i>mülleri</i>	"
	"	<i>Myxosoma dujardini</i>	France
	Conn. tiss. under the mouth muc. mem- brane	<i>Myxobolus fuhrmanni</i>	Switzerland
	Opercle, pseudo branchiae,		France,
	kidney	<i>cycloides</i>	Germany
	Vitreous body of eye	<i>oculi-leucisci</i>	Austria
	?	<i>Henneguya</i> sp.	Germany
	Heart	Gen. et spec. incert.	"
<i>Leuciscus</i> sp.....	Branchiae	<i>Myxobolus minutus</i>	Austria
	"	<i>Myxosoma</i> (?) <i>lobatum</i>	"
<i>Lophius budegassa</i> Spin.....	Gall-bladder	<i>Ceratomyxa appendiculata</i>	Italy, France
<i>L. piscatorius</i> L.....	"	<i>Ceratomyxa appendiculata</i>	"

Host	Organ Infected	Myxosporidian	Locality
<i>L. piscatorius</i> L.....	Urin. bladder	<i>Myxoproteus ambiguus</i>	France, Italy, Hungary
<i>Lota lota</i> L. (<i>L. vulgaris</i>).....	Branchiae	<i>Myxobolus mülleri</i>	Germany
	"	<i>Myxobolus cycloides</i>	Germany
	Gall-bladder	<i>Chloromyxum dubium</i>	" , Austria
	Urin. bladder, kidney	<i>mucronatum</i>	France
<i>Lucioperca lucioperca</i> L..... (<i>L. sandra</i> Cuv.)	Urin. bladder	<i>Myxidium lieberkühni</i>	France, Germany
	"	<i>Sphaerospora elegans</i>	Germany
	Branchiae	<i>Henneguya acerinae</i>	" , Austria
	" , head, fin, opercle	<i>Myxobolus</i> sp.	?
<i>L. volgensis</i> Pall.....	"	<i>Henneguya gigantea</i>	Aus., Servia
	"	Gen. et sp. incert.	Austria
	Muscle	<i>Myxobolus sandrae</i>	Russia
	Branchiae, cor- nea, dorsal fin	<i>volgensis</i>	"
<i>Melanogrammus aeglefinis</i> ..	Gall-bladder	<i>Myxidium bergense</i>	Canada
<i>Menticirrhus americanus</i> L.	"	<i>striatum</i>	Brazil
<i>Merluccius merluccius</i> L. (<i>M. vulgaris</i>).....	"	<i>Ceratomyxa globulifera</i>	France
	"	<i>Leptotheca elongata</i>	" , Italy
	"	Gen. incert. <i>merluccii</i>	Italy
<i>Micropogon undulatus</i>	"	<i>Ceratomyxa aggregata</i>	U. S. A.
<i>Micropterus salmoides</i> Lac..	Urin. Bladder	<i>Henneguya mictospora</i>	U. S. A.
<i>Misgurnus anguillicau- datus</i> Cantor.....	Branchiae	<i>Myxobolus</i> sp.	Nippon
	Gall-bladder	<i>Chloromyxum fujitai</i>	"
	"	<i>misgurni</i>	"
	"	<i>Myxidium kagayamai</i>	"
<i>Molva vulgaris</i> Flem.....	"	<i>Myxobolus misgurni</i>	"
	Bone	<i>aeglefini</i>	Austria
	Gall-bladder	<i>Leptotheca informis</i>	Norway
<i>Motella maculata</i> Risso.....	"	<i>Sphaeromyxa hellandi</i>	"
	"	<i>balbianii</i>	France
<i>M. tricirrata</i> Bl.....	"	<i>Ceratomyxa arcuata</i>	"
	"	<i>Leptotheca elongata</i>	" , Monaco
	"	<i>Sphaeromyxa balbianii</i>	France
	"	<i>sabrazesi</i>	Monaco
<i>Mugil auratus</i> Risso.....	Intestine, stom- ach, spleen, pyloric coecum	<i>Myxobolus mülleri</i>	Italy
	Kidney	<i>exiguus</i>	" , France
	"	<i>exiguus</i>	"
<i>M. capito</i> Cuv.....	"		
<i>M. cephalus</i> L.....	Gall-bladder	<i>Myxidium incurvatum</i>	U. S. A.
<i>M. chelo</i> Cuv.....	Stomach, spleen, kidney, etc.	<i>Myxobolus exiguus</i>	Italy, France

Host	Organ Infected	Myxosporidian	Locality
<i>M. sp.</i>	Kidney	<i>Sphaerospora rostrata</i>	Italy, France
<i>Muraena sp.</i>	Gall-bladder	<i>Ceratomyxa sp.</i>	Monaco
<i>Mustelus canis</i> Mitch. (<i>M. vulgaris</i>).....	Gall-bladder	<i>Ceratomyxa sphaerulosa</i>	France
<i>Nerophis aequoreus</i> L.....	Gall-bladder	<i>Myxidium incurvatum</i>	France
	Muscle	<i>Chloromyxum quadratum</i>	"
<i>N. annulatus</i>	Gall-bladder	<i>Myxidium incurvatum</i>	"
	"	<i>Sphaeromyxa sabrazei</i>	Monaco
<i>N. lumbriciformis</i>	"	<i>Myxidium incurvatum</i>	France
<i>Notropis megalops</i> Raf.....	Subcut. tissue	Gen. et spec. incert.	U. S. A.
<i>N. gilberti</i> J. et M.....	Muscle	<i>Myxobolus orbiculatus</i>	U. S. A.
<i>N. blennioides</i>	"	"	"
<i>N. anogenus</i>	Fins	<i>aureatus</i>	"
	"	<i>Henneguya brachyura</i>	"
<i>Oncorhynchus keta</i>	Under the skin	<i>Henneguya salminicola</i>	Kamtschatka
<i>O. kisutch</i>	"	"	"
	Connective tiss. of body muscle	"	Alaska
<i>O. nerka</i>	Gall-bladder	<i>Chloromyxum wardi</i>	"
<i>Ophidium vasalli</i>	Gall-bladder	<i>Ceratomyxa arcuata</i>	Monaco
<i>Opsanus tau</i>	Urin. bladder	<i>Sphaerospora polymorpha</i>	U. S. A.
<i>Pagellus centrodontus</i> Del.....	Gall-bladder	<i>Ceratomyxa arcuata</i>	France, Italy
<i>Paralichthys albiguttus</i> J. et G.....	Urin. bladder	<i>navicularia</i>	U. S. A.
	"	<i>spinosa</i>	"
	"	<i>Leptotheca glomerata</i>	"
	"	<i>Sinuolinea brachiophora</i>	"
	"	<i>capsularis</i>	"
	"	<i>opacita</i>	"
<i>P. dentatus</i>	Gall-bladder	<i>Ceratomyxa drepanopsettae</i>	"
	Urin. bladder	<i>navicularia</i>	"
	"	<i>Leptotheca lobosa</i>	"
	"	<i>Sinuolinea capsularis</i>	"
<i>Parasilurus asotus</i> L.....	Intestinal wall	<i>Myxobolus miyaii</i>	Nippon
<i>Peprilus alepidotus</i>	Gall-bladder	<i>Ceratomyxa monospora</i>	U. S. A.
<i>Perca flavescens</i>	Spleen	<i>Myxobolus sp.</i>	"
	Urin. bladder	<i>Henneguya wisconsinensis</i>	"
<i>P. fluviatilis</i>	Branchiae	<i>texta</i>	Italy, Germany
	"	<i>Henneguya minuta</i>	"
	"	<i>Myxobolus sp.</i>	Germany
	" , operculum	<i>permagnus</i>	"
	"	<i>Myxosoma dujardini</i>	Switzerland
	"	<i>Henneguya psorospermica</i>	"
<i>Phoxinus (Clinostomus)</i> <i>funduloides</i> Girard	Under scales on ext. surf.	<i>Myxobolus transovalis</i>	U. S. A.
<i>P. laevis</i>	Branchiae	<i>mülleri</i>	France

Host	Organ Infected	Myxosporidian	Locality
<i>P. laevis</i>	Kidney, ovary	<i>Myxidium histophilum</i>	France
	Urin. bladder	<i>Sphaerospora elegans</i>	Germany
<i>Phycis blennioides</i> Br.....	Urin. bladder	<i>Zschokkella hildae</i>	Norway
<i>P. mediterraneus</i> (<i>P. phycis</i> L.).....	Gall-bladder	<i>Leptotheca polymorpha</i>	France
<i>Pimelodus sebae</i> Cuv. et Val.....	Membrane lining branchial cavity	<i>Henneguya linearis</i>	S. America
<i>Pimephales notatus</i> Raf.....	Gall-bladder	<i>Myxobolus notatus</i>	Canada
<i>Piramutana blochi</i> C. et V.	?	<i>inaequalis</i>	S. America
<i>Platystoma fasciatum</i> L.....	Branchiae	<i>Henneguya linearis</i>	"
<i>Pleuronectes flesus</i> L.....	Gall-bladder	? <i>Ceratomyxa drepanopsettae</i>	Norway
<i>P. platessa</i> L.....	"	"	"
	Otic-capsule	<i>Sphaerospora platessae</i>	England
<i>Pomolobus aestivalis</i>	Muscle	<i>Chloromyxum clupeiadae</i>	U. S. A.
<i>P. mediocris</i> Mitch.....	"	<i>clupeiadae</i>	"
<i>P. pseudoharengus</i> Young.	"	<i>clupeiadae</i>	"
<i>Pseudopleuronectes americanus</i>	Gall-bladder	<i>Ceratomyxa acadensis</i>	Canada
	"	<i>Myxidium</i> sp.	"
	Subcutaneous muscul. tissue	<i>Myxobolus pleuronectidae</i>	U. S. A.
<i>Pteroplatea maclura</i>			
Le Sueur.....	Gall-bladder	<i>Chloromyxum leydigi</i>	"
	"	<i>Leptotheca scissura</i>	"
<i>Raja asterias</i>	"	<i>Myxidium giganteum</i>	Italy
<i>R. batis</i> L.....	"	<i>Chloromyxum leydigi</i>	France
	Gall-duct	<i>Myxidium</i> sp.	Germany ?
<i>R. radiata</i>	Gall-bladder	<i>Chloromyxum</i> sp.	Murman Coast
	"	<i>Chloromyxum leydigi</i>	France
<i>R. undulata</i> Lac.....	"	<i>Chloromyxum leydigi</i>	"
	"	<i>Chloromyxum leydigi</i>	France,
<i>Rhina squatina</i> L.....	"	<i>Chloromyxum leydigi</i>	Germany
<i>Rhinobathus</i> sp. (?) Awer....	"	<i>Ceratomyxa</i> sp. (?)	Africa
<i>Rhodeus amarus</i> Bl.....	Branchiae	<i>Myxobolus cycloides</i>	Germany
<i>Salmo fontinalis</i> Mitch.....	Cartilage, perichondrium	<i>Lentospora cerebralis</i>	"
<i>Scardinius erythrophthalmus</i>	Branchiae	<i>Myxobolus cycloides</i>	"
	"	<i>scardinii</i>	Russia
	"	<i>Myxosoma dujardini</i>	France,
	Gall-bladder	<i>Myxidium macrocapsulare</i>	Germany
	Muscle, spleen	<i>Myxobolus dispar</i>	"
	Air-bladder	<i>permagnus</i>	France
	"	<i>physophilus</i>	Germany
<i>Scalophagus argus</i>	Gall-bladder	<i>Ceratomyxa</i> sp. (?)	Russia
			Africa

Host	Organ Infected	Myxosporidian	Locality
<i>Scoliodon terrae-novae</i>	Gall-bladder	<i>Ceratomyxa abbreviata</i>	U. S. A.
	"	<i>attenuata</i>	"
	"	<i>sphaerophora</i>	"
	"	<i>taenia</i>	"
	"	<i>Chloromyxum leydigi</i>	"
<i>Scomber scombrus</i> L.....	"	<i>Leptotheca parva</i>	France, Norway
	Kidney	<i>renicola</i>	France
	?	Gen. et sp. inc.	Germany?
<i>Scorpaena porcus</i> L.....	Gall-bladder	<i>Ceratomyxa arcuata</i>	France
<i>S. scrofa</i> L.....	"	<i>Ceratomyxa arcuata</i>	"
	"	<i>Myxidium incurvatum</i>	"
<i>S. sp.</i>	"	<i>Leptotheca agilis</i>	" , Germany
<i>Scyllium canicula</i>	"	<i>Ceratomyxa sphaerulosa</i>	Monaco
	"	<i>Chloromyxum leydigi</i>	" , France Germany
<i>S. asterias</i>	"	"	Italy
<i>Sebastes dactylopterus</i>	"	<i>Leptotheca macrospora</i>	Norway
<i>S. norvegicus</i>	"	<i>Leptotheca</i> sp.	Eastern Finmark
<i>S. viviparus</i> H. Kr.....	"	<i>Leptotheca macrospora</i>	Norway
<i>Siphonostoma rondeletii</i>	"	<i>Sphaeromyxa sabrazesi</i>	Monaco
<i>Siphostoma floridae</i>	"	<i>balbianii</i>	U. S. A.
	Urin. bladder	<i>Sinuolinea arborescens</i>	"
<i>S. louisianae</i>	Gall-bladder	<i>Sphaeromyxa balbianii</i>	"
<i>Solea vulgaris</i>	"	<i>Myxidium gadi</i>	France
<i>Sparus berda</i>	"	<i>Ceratomyxa</i> (?) <i>spari</i>	Africa
<i>Sphaeroides maculatus</i>	Urin. bladder	<i>navicularia</i>	U. S. A.
	"	<i>Sinuolinea capsularis</i>	"
	"	<i>Zschokkella globulosa</i>	"
<i>Spinax spinax</i> L.....	Gall-bladder	<i>Chloromyxum leydigi</i>	Italy, France
<i>Squalis agassizii</i> Heck.....	Branchiae	<i>Myxobolus mülleri</i>	France
<i>S. a. Savigny</i> Bona- parte.....	"	<i>Myxobolus mülleri</i>	"
<i>Stenotomus chrysops</i> L.....	Muscle	<i>Chloromyxum clupeiidae</i>	U. S. A.
<i>Stizostedion vitreum</i> Mitch.	Urin. bladder	Gen. et sp. incert.	Canada
<i>Syngnathus acus</i> L.....	Gall-bladder	<i>Myxidium incurvatum</i>	France
	"	<i>Sphaeromyxa sabrazesi</i>	Italy, Monaco
	Muscle	<i>Chloromyxum quadratum</i>	France
<i>S. typhle</i>	Gall-bladder	<i>Myxidium incurvatum</i>	France
<i>Synodontis schall</i> Bl. Schn..	Integum. of cephalic reg.	<i>Henneguya strongylura</i>	Egypt
	?	<i>Myxobolus inaequalis</i>	S. America
<i>Synodus faetans</i>	Gall-bladder	<i>Ceratomyxa agglomerata</i>	U. S. A.
	"	<i>amorpha</i>	"
<i>Tautogolabrus adspersus</i> Walb.....	Muscle	<i>Chloromyxum clupeiidae</i>	"

Host	Organ Infected	Myxosporidian	Locality
<i>Thymallus thymallus</i> L.....	Gall-bladder	<i>Chloromyxum thymalli</i>	Austria
	"	<i>Myxobolus</i> sp.	"
	Neurilemma (?)	<i>pfeifferi</i>	Germany ?
<i>Thysanophris japonicus</i>	Gall-bladder	<i>Sphaeromyxa exneri</i>	Africa
<i>Tinca tinca</i> L. (<i>T. vulgaris</i>).	Branchiae	<i>Myxobolus piriformis</i>	France, Germany
	Air-bladder, kidney, etc.	<i>ellipsoides</i>	"
	Gall-bladder	<i>Chloromyxum cristatum</i>	France
	"	<i>Myxidium pfeifferi</i>	Germany
	Kidney	<i>Myxobolus cyprini</i>	Germany, Hungary
<i>Torpedo narce</i> Risso.....	Gall-bladder	<i>Chloromyxum leydigi</i>	France
<i>T. ocellata</i>	"	<i>Chloromyxum leydigi</i>	Germany
<i>T. torpedo</i> L.....	"	<i>Chloromyxum leydigi</i>	"
<i>Trachinus draco</i> L.....	"	<i>Ceratomyxa reticularis</i>	"
	"	<i>Myxidium incurvatum</i>	"
<i>Trachurus trachurus</i> L.....	Muscle	<i>Chloromyxum quadratum</i>	France, Germany
<i>Trutta fario</i> L.....	Gall-bladder, Gall-duct	<i>Chloromyxum truttiae</i>	France
	Nervous syst.	<i>Myxobolus neurobius</i>	Germany
	Subcutaneous conn. tiss. at base of fin	<i>Henneguya nüsslini</i>	"
<i>T. iridea</i> Gibb.....	Cartilage, peri- chondrium	<i>Lentospora cerebrealis</i>	"
<i>T. salar</i> L.....	"	"	"
	Gall-bladder	<i>Myxidium oviforme</i>	Norway
<i>Trygon pastinaca</i> L.....	Gall-bladder	<i>Chloromyxum leydigi</i>	France
	"	<i>Leptotheca agilis</i>	" , Italy
<i>Tylosurus marianus</i>	Urin. bladder	<i>Chloromyxum granulosum</i>	U. S. A.
<i>T. schismatorhynchus</i>	Gall-bladder	<i>Ceratomyxa tylosuri</i>	Africa
<i>Urophycis chuss</i>	"	<i>acadiensis</i>	Canada
<i>Zoarces angularis</i>	"	<i>acadiensis</i>	"
Amphibia			
<i>Bufo lentiginosus</i>	Kidney	<i>Wardia ohlmacheri</i>	U. S. A.
<i>B. marinus</i> L.....	Gall-bladder	<i>Sphaeromyxa immersa</i>	Brazil
<i>Hyla aurea</i>	Testis, ovary	<i>Myxobolus hylae</i>	Australia
<i>Leptodactylus ocellatus</i>	Gall-bladder	<i>Sphaeromyxa immersa</i>	Brazil
<i>Molge cristata</i> Laur. (<i>Triton</i> c.).....	"	<i>Chloromyxum caudatum</i>	France
<i>Proteus anguineus</i> L.....	Kidney	<i>Chloromyxum protei</i>	Austria
<i>Rana esculenta</i>	"	? <i>Wardia ohlmacheri</i>	France ?
<i>R. temporaria</i> (<i>R. fusca</i>).....	"	? <i>Wardia ohlmacheri</i>	"
Reptilia			
<i>Emys orbicularis</i> L. (<i>Cistudo europaea</i>).....	Kidney	<i>Myxidium danilewskyi</i>	Russia, France

Host	Organ Infected	Myxosporidian	Locality
<i>Lacerta</i> sp.....	Ovarian egg	Gen. et spec. incert.	Italy
<i>Trionyx (Amyda) ganzeticus</i>	Kidney	<i>Myxidium mackiei</i>	India
<i>T. spinifera</i>	"	<i>americanum</i>	U. S. A.

C. DISTRIBUTION OF MYXOSPORIDIA IN THE ORGANS OF THE HOST

Altho some species are found in various organs of the host animal, the majority has one or two particular seats of infection. Among the various organs which become infected, the gall-bladder is most frequently infected. The kidney, branchia and urinary bladder have less chances of being parasitized. As to the infection of the reproductive organs of the host, little is known. The male reproductive organ becomes rarely infected, being reported only twice. The female reproductive organ, however, is more frequently infected. The infection of the next generation of the host animal thru the infected ovum which is known to occur in some Microsporidian parasites, has not been reported in Myxosporidia as yet.

LIST IV. ORGANS OF HOST INFECTED BY MYXOSPORIDIA

I. PISCES

- 1) Integument.—*Sphaerospora* sp. Southwell et Prashad (under the scales)
Myxobolus seni Southwell et Prashad (fin)
Myxobolus transovalis Gurley (under the scales)
Myxobolus unicus Gurley (head)
Myxobolus cycloides Gurley (opercle)
Myxobolus inaequalis Gurley (head)
Myxobolus sp. Gurley (opercle, head, fin)
Myxobolus squamae Keysselitz (inner surface of the scales)
Myxobolus volgensis Reuss (fin)
Myxobolus permagnus Wegener (operculum)
Myxobolus aureatus Ward (fin)
Henneguya brachyura Ward (fin-ray)
Henneguya linearis (Gurley) Labbé (membrane lining branchial cavity)
Henneguya gurleyi Kudo (base of spines of dorsal fin)
Henneguya strongylura (Gurley) Labbé (cephalic region)
Lentospora dermatobia Ishii
- 2) Connective tissue.—*Myxidium anguillae* Ishii (subcutaneous)
Myxobolus fuhrmanni Auerbach (under the oral mucous membrane)
Myxobolus oviformis Thélohan (subcutaneous)
Myxobolus lintoni Gurley (subcutaneous)
Myxobolus oblongus Gurley (chiefly of the head)
Myxobolus gigas Auerbach (of operculum, sides and fins)
Myxobolus capsulatus Davis (visceral)
Henneguya kolesnikovi (Gurley) Labbé (interstitial)
Henneguya nüsslini Schuberg et Schröder (at the base of dorsal fin)
Henneguya miyairii Kudo (of the head)
Gen. et sp. incert. Linton (subcutaneous)

- 3) Muscle.—*Leptotheca perlata* (Gurley) Labbé
Chloromyxum quadratum Thélohan
Chloromyxum funduli Hahn
Chloromyxum clupeiidae Hahn
Lentospora multiplicata Reuss
Myxobolus notatus Mavor (connective tissue of voluntary muscle)
Myxobolus pfeifferi Thélohan
Myxobolus sandrae Reuss
Myxobolus musculi Keysselitz
Myxobolus funduli Kudo
Myxobolus pleuronectidae Hahn
Myxobolus sp. Southwell (subcutaneous intermuscular tissue)
Myxobolus nodularis Southwell et Prashad
Myxobolus orbiculatus Kudo
Henneguya creplini (Gurley) Labbé
Henneguya monura (Gurley) Labbé
Henneguya zschokkei (Gurley) Doflein
Henneguya salminicola Ward
 Gen. et spec. incert. Leydig
- 4) Eye.—*Sphaerospora platessae* Woodcock (optic capsule)
Myxobolus oculi-leucisci Trojan (vitreous body)
Myxobolus ellipsoides Thélohan
Myxobolus mülleri Bütschli
Myxobolus aeglefini Auerbach
Myxobolus volgensis Reuss
Myxobolus magnus Awerinzew
Henneguya schizura (Gurley) Labbé (intercellular tissue of eye muscle)
- 5) Branchiae.—*Sphaerospora carassii* Kudo
Myxosoma dujardini Thélohan
Myxosoma (?) *lobatum* Nemeček
Myxosoma funduli Kudo
Lentospora acuta (Fujita)
Myxobolus piriformis Thélohan
Myxobolus toyamai Kudo
Myxobolus rohita Southwell et Prashad
Myxobolus dispar Thélohan
Myxobolus ellipsoides Thélohan
Myxobolus exiguus Thélohan
Myxobolus oviformis Thélohan
Myxobolus mülleri Bütschli
Myxobolus globosus Gurley
Myxobolus cycloides Gurley (also pseudobranchiae)
Myxobolus sphaeralis Gurley
Myxobolus anurus Cohn
Myxobolus sp. Gurley
Myxobolus gigas Auerbach
Myxobolus volgensis Reuss
Myxobolus scardinii Reuss
Myxobolus macrocapsularis Reuss
Myxobolus bramae Reuss
Myxobolus cyprinicola Reuss

Myxobolus balleri Reuss
Myxobolus sp. Miyairi
Myxobolus sp. Wegener
Myxobolus permagnus Wegener
Myxobolus rotundus Nemeček
Myxobolus minutus Nemeček
Myxobolus funduli Kudo
Myxobolus koi Kudo
Myxobolus discrepans Kudo
Henneguya psorospermica Thélohan
Henneguya texta (Cohn)
Henneguya minuta (Cohn)
Henneguya lobosa (Cohn)
Henneguya creplini (Gurley) Labbé
Henneguya linearis (Gurley) Labbé
Henneguya acerinae Schröder
Henneguya gigantea Nemeček
 Gen. et spec. incert. Heckel et Kner

- 6) Heart.—*Myxobolus cordis* Keysselitz (muscle of ventricle and bulbus arteriosus)
 Gen. et spec. incert. Leydig (auriculo-ventricular valve)
- 7) Air bladder.—*Myxobolus ellipsoides* Thélohan (conn. tiss.)
Myxobolus mülleri Bütschli (conn. tiss.)
Myxobolus physophilus Reuss (surface)
Myxobolus permagnus Wegener
- 8) Body-cavity (cyst).—*Myxobolus* sp. Gurley
Myxobolus carassii Klokacewa
 Gen. et spec. incert. Leydig
- 9) Nervous tissue.—*Myxobolus neurobius* Schuberg et Schröder
Lentospora encephalina Mulsow (blood vessel of brain)
- 10) Bone, cartilage, perichondrium.—*Lentospora cerebialis* (Hofer) Plehn
Myxobolus aeglegini Auerbach
Henneguya brachyura Ward (of fin)
- 11) Stomach, pyloric cecum.—*Myxobolus exiguus* Thélohan
Myxobolus mesentericus Kudo
Henneguya tenuis Vaney et Conte
- 12) Liver.—*Myxobolus ellipsoides* Thélohan
Myxobolus oviformis Thélohan
Myxobolus cyprini Doflein
Myxobolus cordis Keysselitz
Myxobolus muscoli Keysselitz
Myxobolus carassii Klokacewa
Myxobolus mesentericus Kudo
- 13) Gall-bladder.—*Leptotheca agilis* Thélohan
Leptotheca elongata Thélohan
Leptotheca polymorpha (Thélohan) Labbé
Leptotheca parva Thélohan
Leptotheca hepseti Thélohan
Leptotheca sp. Awerinzew
Leptotheca macrospora Auerbach
Leptotheca informis Auerbach
Leptotheca longipes Auerbach

- Leptotheca fusiformis* Davis
Leptotheca scissura Davis
Ceratomyxa arcuata Thélohan
Ceratomyxa sphaerulosa Thélohan
Ceratomyxa pallida Thélohan
Ceratomyxa globurifera Thélohan
Ceratomyxa appendiculata Thélohan
Ceratomyxa truncata Thélohan
Ceratomyxa reticularis Thélohan
Ceratomyxa inaequalis Doflein
Ceratomyxa lino-spora Doflein
Ceratomyxa ramosa Awerinzew
Ceratomyxa drepano-psettae Awerinzew
Ceratomyxa tylosuri Awerinzew
Ceratomyxa (?) *spari* Awerinzew
Ceratomyxa sp. (?). Awerinzew
Ceratomyxa sp (?). Awerinzew
Ceratomyxa acadensis Mavor
Ceratomyxa sp. Georgévitch
Ceratomyxa coris Georgévitch
Ceratomyxa herouardi Georgévitch
Ceratomyxa meso-spora Davis
Ceratomyxa sphaero-phora Davis
Ceratomyxa taenia Davis
Ceratomyxa attenuata Davis
Ceratomyxa recurvata Davis
Ceratomyxa lunata Davis
Ceratomyxa abbreviata Davis
Ceratomyxa flagellifera Davis
Ceratomyxa agglomerata Davis
Ceratomyxa amorph-a Davis
Ceratomyxa mono-spora Davis
Ceratomyxa strepto-spora Davis
Ceratomyxa aggregata Davis
Ceratomyxa undulata Davis
Chloromyxum leydi-gi Mingazzini
Chloromyxum fluviatile Thélohan
Chloromyxum truttiae Léger (also gall-duct)
Chloromyxum cristatum Léger
Chloromyxum dubium Auerbach
Chloromyxum sp. Awerinzew
Chloromyxum thymalli Lebzelter
Chloromyxum koi Fujita
Chloromyxum magnum Awerinzew
Chloromyxum misgurni Kudo
Chloromyxum fujitai Kudo
Chloromyxum trijugum Kudo
Chloromyxum catostomi Kudo
Chloromyxum wardi Kudo
Sphaerospora masovica Cohn
Myxidium incurvatum Thélohan
Myxidium sphaericum Thélohan

Myxidium sp. Gurley (only in gall-duct)
Myxidium giganteum Doflein
Myxidium pfeifferi Auerbach
Myxidium inflatum Auerbach
Myxidium bergense Auerbach
Myxidium procerum Auerbach
Myxidium macrocapsulare Auerbach
Myxidium sp. Awerinzew
Myxidium oviforme Parisi
Myxidium sp. Mavor
Myxidium kagayamai Kudo
Myxidium gadi Georgévitch
Myxidium glutinosum Davis
Myxidium phyllium Davis
Myxidium striatum Cunha et Fonseca
Myxobolus misgurni Kudo (few spores only)
Myxobolus sp. Lebzelter (spores only)
Sphaeromyxa babianii Thélohan
Sphaeromyxa incurvata Doflein
Sphaeromyxa sabrazesi Laveran et Mesnil
Sphaeromyxa hellandi Auerbach
Sphaeromyxa exneri Awerinzew
Sphaeromyxa gasterostei Georgévitch
Zschokkella nova Klokacewa
Zschokkella acheilognathi Kudo (also in gall-duct)
 Gen. incert. *congru* Perugia
 Gen. incer. *merlucii* Perugia
 Gen. et spec. incert. Mavor

- 14) Spleen.—*Myxobolus piriformis* Thélohan
Myxobolus sp. Kudo
Myxobolus ellipsoides Thélohan
Myxobolus exiguus Thélohan
Myxobolus oviformis Thélohan
Myxobolus pfeifferi Thélohan
Myxobolus cyprini Doflein
Myxobolus cordis Keysselitz
Myxobolus musculi Keysselitz
Myxobolus mesentericus Kudo
- 15) Intestine.—*Myxobolus exiguus* Thélohan
Myxobolus mülleri Bütschli
Myxobolus pfeifferi Thélohan
Myxobolus miyairii Kudo
Myxobolus carassii Klokacewa
Myxobolus mesentericus Kudo
Henneguya peri-intestinalis Cépède
Henneguya tenuis Vaney et Conte
- 16) Ovary.—*Wardia ovinocua* Kudo
Sphaerospora elegans Thélohan
Myxidium histophilum Thélohan
Myxobolus pfeifferi Thélohan
Myxobolus mülleri Bütschli
Myxobolus musculi Keysselitz

Henneguya oviperda (Cohn)

Henneguya media Thélohan

Henneguya brevis Thélohan

- 17) Kidney.—a) Urinary tubules.—*Leptotheca renicola* Thélohan
Mitraspora caudata (Parisi) Kudo
Mitraspora cyprini Fujita (also in ureter)
Mitraspora elongata Kudo
Sphaerospora elegans Thélohan
Sphaerospora divergens Thélohan
Henneguya media Thélohan
Henneguya brevis Thélohan
Henneguya gasterostei Parisi
Hoferellus cyprini Doflein
- b) Tissue.—*Mitraspora elongata* Kudo
Chloromyxum quadratum Thélohan
Sphaerospora rostrata Thélohan (Malpighian bodies)
Myxidium histophilum Thélohan
Lentospora asymmetrica Parisi
Myxobolus pfeifferi Thélohan
Myxobolus cyprini Doflein
Henneguya neapolitana Parisi (conn. tiss. of ren. tubules)
Hoferellus cyprini Doflein
- c) Seat, unstated.—*Chloromyxum mucronatum* Gurley
Sphaerospora angulata Fujita
Myxidium barbatulae Cépède
Myxidium giardi Cépède
Myxobolus piriformis Thélohan
Myxobolus ellipsoides Thélohan
Myxobolus exiguus Thélohan
Myxobolus oviformis Thélohan
Myxobolus mülleri Bütschli
Myxobolus obesus Gurley
Myxobolus cycloides Gurley
Myxobolus cordis Keysselitz
Myxobolus musculi Keysselitz
- 18) Urinary bladder.—*Leptotheca lobosa* Davis
Leptotheca glomerata Davis
Ceratomyxa navicularia Davis
Ceratomyxa spinosa Davis
Chloromyxum mucronatum Gurley
Chloromyxum granulosum Davis
Sphaerospora elegans Thélohan
Sphaerospora divergens Thélohan
Sphaerospora polymorpha Davis
Sphaerospora sp. Davis
Sinuolinea dimorpha Davis (also in ureter)
Sinuolinea capsularis Davis
Sinuolinea arborescens Davis
Sinuolinea opacita Davis
Sinuolinea brachiophora Davis
Myxidium lieberkühni Bütschli
Zschokkella hildae Auerbach
Zschokkella globulosa Davis

Myxoproteus ambiguus (Thélohan) Doflein
Myxoproteus cordiformis Davis
Myxoproteus cornutus Davis
Henneguya legeri Cépède
Henneguya wisconsinensis Mavor et Strasser
Henneguya mictospora Kudo
 Gen. et spec. incert. Mavor

19) Testis.—*Myxobolus pfeifferi* Thélohan (only spores)

20) Mesentery.—*Myxobolus mesentericus* Kudo

21) Seat unknown.—*Henneguya* sp. Gurley (integument?)

Gen. et spec. incert. Borne

II AMPHIBIA

1) Gall-bladder.—*Chloromyxum caudatum* Thélohan

Sphaeromyxa immersa (Lutz) Thélohan

2) Urinary tubules of kidney.—*Wardia ohlmacheri* (Gurley) Kudo

Chloromyxum protei Joseph

TABLE I

	Integument	Conn. tissue	Muscle	Eye	Gill	Heart	Air-bladd.	Nerv. tiss.	Bone, cartilage	Stomach wall	Liver	Gall-bladd.	Spleen	Intestine wall	Testis	Ovary	Kidney	Body cavit.	Urinary bladder	Mesentery	Seat unknown
Leptotheca.....			1									11					1(a)*		2		
Ceratomyxa.....												33							2		
Myxoproteus.....																			3		
Wardia.....																1	1(a)				
Mitraspora.....																	3(a)				
																	1(b)*				
Chloromyxum.....			3									14					1(a)				
																	1(b)				
																	1(c)*	1	2		
Sphaerospora.....	1			1	1							1				1	2(a)				
																	1(b)				
																	1(c)		4		
Sinuolinea.....																	2(a)		5		
Myxidium.....		1										18				1	1(b)				
																	2(c)				
																	3(a)		1		
Sphaeromyxa.....												7									
Zschokkella.....												2							2		
Myxosoma.....					3																
Lentospora.....	1		1	1			1	1									1(b)	1			
Myxobolus.....	10	6	9	6	28	1	4	1	1	2	7	2	10	6	2	4	2(b)				
																	9(c)	1		1	1
Henneguya.....	4	3	4	1	8				1	1				2	3		3(a)		3		1
																	1(b)				
Hoferellus.....																	1(a)				
																	1(b)				
Total.....	16	10	18	8	41	1	4	2	3	3	7	88	10	8	2	10	39	3	24	1	2

* For (a), (b) and (c), see page 42.

3) Testis, oviduct.—*Myxobolus hylae* Johnston et Bancroft

III REPTILIA

1) Kidney (ren. tub.).—*Myxidium danilewskyi* Laveran

Myxidium mackiei Bosanquet

Myxidium americanum Kudo

2) Ovary.—Gen. et spec. incert. Mingazzini

IV INSECTA

1) Abdominal cavity.—*Chloromyxum diploxys* Gurley

V ANNELIDA

Myxobolus sp. Gurley

The data in this section are summarized on the preceding page (Table I).

D. THE EFFECT OF ENVIRONMENT ON THE ORGANAL DISTRIBUTION OF MYXOSPORIDIA IN HOSTS

Myxosporidia are almost equally distributed among marine and fresh-water fishes in regard to the number of species. This is shown in the following table.

TABLE II

Genus	Number of species found in marine fish	Number of species found in fresh-water fish	Other hosts			
			Rept.	Amph.	Insect	Annelida
Leptotheca (15)*....	15
Ceratomyxa (35)....	35
Myxoproteus (3)....	3
Wardia (2).....	1	1
Mitraspora (3).....	3
Chloromyxum (22)	7	12	2	1
Sphaerospora (10)	5	5
Sinuolinea (5).....	5
Myxidium (26).....	17	8	3
	(2 common)	(2 common)				
Sphaeromyxa (7)....	6	1
Zschokkella (4)....	2	2
Myxosoma (3).....	1	2
Lentospora (6).....	2	6
	(2 common)	(2 common)				
Myxobolus (63).....	5	56	1	1
Henneguya (32)....	1	31
Hoferellus (1).....	1
Gen. et spec. incert (12).....	4	7	1
TOTAL 237+12	104+4	134+7	4	5	1	1

* The number in parenthesis denotes the number of species in the corresponding genus.

These genera have certain relations to the organal distribution in the body of the host, which are shown in List IV (page 37) and Table I (on page 43) and which can be put together as follows:

TABLE III

Genus	Number of species found in body-cavity	Number of species found in tissue	Number of species found in both	Seat unknown
Leptotheca (15).....	14	1
Ceratomyxa (35).....	35
Myxoproteus (3).....	3
Wardia (2).....	1	1
Mitraspora (3).....	2	..	1	..
Chloromyxum (22).....	18	4
Sphaerospora (10).....	4	4	1	1
Sinuolinea (5).....	5
Myxidium (26).....	22	4
Sphaeromyxa (7).....	6	1
Zschokkella (4).....	4
Myxosoma (3).....	..	3
Lentospora (6).....	1	5
Myxobolus (63).....	2	59	..	2
Henneguya (32).....	4	28
Hoferellus (1).....	1	..
Gen. et spec. inct (12)....	4	5	..	3
TOTAL 237+12.....	121+4	109+5	3	4+3

From the facts shown in the above tables, the following conclusions can be drawn.

1) The genera *Leptotheca* (one species in tissue), *Ceratomyxa*, *Myxoproteus*, *Sinuolinea* and *Sphaeromyxa* (one species in Amphibia) include parasites from the body cavity of marine fish.

2) The majority of the genera *Lentospora*, *Myxosoma*, *Myxobolus* (one species in Amphibia), *Henneguya* and *Hoferellus* include parasites in tissues of fresh-water fish.

3) The genera *Chloromyxum*, *Sphaerospora*, *Myxidium* and *Zschokkella* include forms that infect the body-cavity as well as the tissue of marine and fresh-water fishes.


4) The genus *Mitraspora* includes three species that parasitize the fresh-water fish.

5) The genus *Myxidium* has three species found in the kidney of reptiles.

- 6) The genera *Wardia*, *Chloromyxum*, *Sphaeromyxa* and *Myxobolus* include species parasitic in Amphibia.
- 7) The genus *Chloromyxum* has one species found in an insect.
- 8) The genus *Myxobolus* has one species found in an annelid, which was not normally recorded.

THE SPORE

As will be shown later (page 52-55), the spore stage is still the only constant character by which various forms of Myxosporidia are identified from each other. For this reason it is necessary to have a clear conception of the form and structure of the spore and at the same time to define the terms used in the present paper, even tho they have commonly been used heretofore.



TEXTFIGURE

DIAGRAMMATICAL FRONT [A] AND SIDE [B] VIEWS OF A MYXOBOLUS SPORE.

(For further explanation see following pages.)

The spore of Myxosporidia is covered by a shell, which is composed of two valves usually symmetrical in form and size that come in contact in the sutural plane. The sutural line is straight in most cases, tho sometimes curved like an S. It is more or less thickened, forming the sutural ridge. The sutural ridge is to be made out clearly in fresh as well as in stained preparations and furnishes important data in regard to the classification of the parasite. The thickness of the shell-valve is usually uniform; in some species (*Myxobolus*), however, it may differ slightly in different parts of the shell. Besides, in many species of *Myxobolus*, the shell differentiates a small triangular intercapsular appendix on the inside at the anterior end directed posteriad between two polar capsules.

The form of the spore varies greatly owing to the shape of the shell together with its variously developed appendages; 1) lateral appendages as in *Ceratomyxa*, 2) anterior processes as in *Myxoproteus*, 3) posterior processes as in *Wardia* (fringe-like), *Mitraspora* (filiform), *Hoferellus* (spinous), *Henneguya* (tail-form), etc.

The surface of the shell may be smooth or exhibit various markings. More or less conspicuous ridges varying in form and number in different species, may run parallel to the sutural line, may show a network-like structure or may exhibit short tooth-like processes arising from the sutural ridge and radiating toward the center of each valve. When the ridges are fine, they form delicate striations, arranged usually parallel to the sutural line. Tho these markings are usually easily seen in vivo, they are very often more readily studied in stained preparations.

Inside of the shell are present the polar capsules and sporoplasm. Gurley (1894:120) and Davis (1917:210) used the term "capsules" instead of polar capsules because of the facts that "the situation implied by the the latter (polar capsule) is not constant" (Gurley) and that "they are often not in the position indicated by the term polar capsule" (Davis). The present writer, however, does not agree with these authors and retains the commonly used term, polar capsule, thruout the present paper on the basis of the fact that these polar capsules are situated at or near the more or less attenuated anterior end in the great majority of species or at each end (in Myxidiidae) of the spore, except in the few cases as in *Wardia* in which they are situated in the central portion and have the foramina at the anterior end of the spore.

The polar capsules may be pyriform or spherical. They are located at or near one end (anterior end) of the spore. In Myxidiidae, one polar capsule is situated at each end, in which case no distinction can be made between the anterior and posterior ends. The end or side opposite to the anterior, is the posterior end of the spore. The number of polar capsules in a spore varies according to the different genera. There is only one polar capsule in the spore of unicapsular *Myxobolus*, four in *Chloromyxum*, two in all the other genera. They may be equal or unequal in form and size. When two polar capsules are located at the anterior end, they may be convergent or divergent. Each has a foramen to the outside of the spore thru the shell in or near the sutural line, thru which the polar filament is extruded. The foramen is observable in the fresh condition. Staining will very often show clearly the canals thru the shell. Each polar capsule has an independent foramen.

In the polar capsule exists a coiled polar filament, which in most cases can be recognized without difficulty in the fresh condition. The polar filament is as a rule a more or less extended, probably hollow thread connected with the polar capsule, which is extruded from the spore thru the foramen under the action of the stimulants such as the digestive fluid of the host or certain chemicals. In *Sphaeromyxa* it is rather short and thick, tapering to a point. The polar filament is coiled around the longest axis of the polar capsule, except in *Sphaeromyxa* in which it is coiled around an axis perpendicular to the longest axis of the polar capsule.

The sporoplasm occupies the extracapsular cavity at the posterior region of the spore. It is of granular structure with almost always two nuclei. Besides, it has an iodophilous vacuole mostly round or oval in the spores of the family Myxobolidae. It occurs thruout the spore stage and is the important character of the said family. The contents of the vacuole is probably of glycogenous nature and is stained deeply with iodine. Small refringent fat globules have also been observed in the spore.

Davis (1917:212) proposed to use capsular and postcapsular sides in place of anterior and posterior ends which have most frequently been used and are also used in the present paper. The latter terms can be employed as properly as Davis' terms except in the case of the Myxidiidae, where both terms, strictly speaking, are inapplicable.

Tho various abnormal spores are very often encountered in several species, the majority of the spores are of typical form, structure and size. In *Myxosoma* and *Myxobolus*, the spore sometimes develops a short posterior process, which is highly developed in the spore of the genus *Henneguya*.

Young spores, generally speaking, are more rounded in form than the mature form, while the mature spores, as a rule, are of definite form, structure and size characteristic to the species. It should, however, be kept in mind that there is a certain amount of variation among these characters.

As is generally recognized, one must mention whether the spores were measured in fresh condition or in fixed and stained state. The fresh spore is generally more or less larger than the mounted one.

DEFINITION OF TERMS USED FOR DESCRIPTIONS

- Anterior end.*—The end of the spore where the polar capsules open; in most cases the polar capsules are situated at this end.
- Anterior process.*—The spinous process of the shell at the anterior end of the spore of the genus *Myxoproteus*.
- Breadth.*—The larger diameter of the spore measured at right angles to the length or sutural diameter; the shorter diameter thus measured being the thickness.
- Capsulogenous cell.*—A small island of protoplasm with a nucleus, in which polar capsule becomes differentiated.
- Cyst.*—The vegetative form of more or less conspicuous size in tissues of the host, surrounded usually by a membranous structure composed of the host issue.
- Disporous.*—The character of a trophozoite of forming only two spores.
- Foramen.*—Opening of the polar capsule thru which the polar filament is extruded.
- Front view.*—The view in which length and breadth of the spore are laid horizontally.
- Gemmules.*—A small mass of trophozoite separated from the mother body by plasmotomy. Used by Davis (1917). (See page 105.)
- Iodinophilous vacuole.*—The vacuole in the sporoplasm of the spore of the family *Myxobolidae*, the contents of which are stained brownish with iodine.
- Lateral process.*—The lateral prolongation of the shell-valve at right angles to the sutural plane.
- Length.*—Antero-posterior diameter of the spore in the sutural plane; equivalent to sutural diameter.
- Longitudinal striations.*—Fine ridges or thickenings marked longitudinally on the shell of the spore.
- Mesoplasm.*—An intermediate layer between ectoplasm and endoplasm, coined by Cohn in the case of *Myxidium lieberkühni* (see page 107).
- Mictosporous.*—The character of the trophozoite of forming a variable number of spores in an individual.
- Monosporous.*—The character of the trophozoite of forming a single spore.
- Pansporoblast.*—Coined by Gurley (1893:408) used here in the same meaning, an enclosed area in the endoplasm of the vegetative form, in which two sporoblasts become differentiated.

Plasmogamy. Fusion of two trophozoites, coined by Doflein (1898).

Plasmotomy.—Division of trophozoite into daughter individuals, coined by Doflein (1898).

Polar capsule.—The pyriform or spherical, hollow body in the spore which forms a polar filament.

Polar filament.—The filament which is coiled inside the polar capsule.

Polysporous.—The character of the trophozoite of forming spores, more than two.

Posterior filament.—Fine posterior appendage of the spore.

Posterior processes.—Posterior differentiations of the shell.

Ridge.—Linear or network-like elevation of the shell of the spore.

Shell.—The envelope of the spore.

Shell-valves.—Two valves which compose the shell of the spore.

Sporoplasm.—The protoplasmic mass found inside of the spore (amebula or sporozoite), usually situated in the posterior portion of the spore.

Sutural diameter.—Same as length.

Sutural edge.—The edge of the shell-valves cut by the sutural plane.

Sutural line.—The line on the shell of the spore marked by the sutural plane.

Sutural plane.—The plane on which two shell-valves meet together.

Sutural ridge.—The ridge marking the sutural line.

Tail.—The posterior prolongation of the valves from the median posterior end; it may be a single process or bifurcated.

Thickness.—See breadth.

Trophozoite.—The vegetative or multiplicative stage of a Myxosporidian.

Vegetative form.—Same as trophozoite.

CLASSIFICATION OF MYXOSPORIDIA

The classification of Myxosporidia, was first carried out by Thélohan as early as 1892, who considered rightly that the spore was the only reliable means for the purpose. In 1899 and 1901, Doflein introduced into the classification two Legions, Disporea and Polysporea, and a new family. This plan has generally been followed by various authors in dealing with these protozoa.*

The classification of the said author, however, no longer agrees with our present knowledge of the animals. In the first place, as was pointed out by some authors, for instance Davis (1917:217), it is far from being correct to divide the Myxosporidia into two Legions, Disporea and Polysporea, on the basis of the number of spores formed in each vegetative form, since this differs even in one and the same species as was observed by Léger, Auerbach, Awerinzew, Parisi, Georgévitch, Davis, Kudo and others (see Table IV on page 53).

Auerbach who had observed numerous interesting facts in this group, had adopted Doflein's classification in his splendid work (1910) by simply adding two genera, Zschokkella and Lentospora, to the family Myxidiidae. In the following year (1911), he tried a new classification, on the same basis as Doflein did, by introducing two new Legions besides these two already existing, and by discarding all the families. Thus:

I Monosporea	a) Genus	Coccomyxa
II Mictosporea	a) Genus	Zschokkella
	b) Genus	Myxoproteus
	c) Genus	Myxidium
	d) Genus	Sphaeromyxa
	e) Genus	Chloromyxum
	f) Genus	Sphaerospora
III Disporea	a) Genus	Ceratomyxa
	b) Genus	Leptotheca
IV Polysporea	a) Genus	Myxosoma
	b) Genus	Lentospora
	c) Genus	Myxobolus
	d) Genus	Henneguya
	e) Genus	Hoferellus

As will be distinctly seen from Table IV, the classification not only fails to improve Doflein's classification in bringing together the genera, Myxo-

* Doflein still uses the same classification in his recent work (1916).

proteus, Myxidium and Sphaeromyxa into Mictosporaea, and Lentospora and Henneguya into Polysporaea, but increases the confusion concerning relationship among the genera.

TABLE IV

Genus	Mono- and di- sporous	Mono- and poly- sporous	Di- sporous	Mono, di- and poly- sporous	Di- and poly- sporous	Poly- sporous	Unknown
Leptotheca 15 species.....	8	7
Ceratomyxa 35 species.....	3	..	23	..	4	..	5
Myxoproteus 3 species.....	2	1
Wardia* 2 species.....	1	1
Mitraspora* 3 species.....	2	1	..
Chloromyxum 22 species.....	1	2	..	2	6	4	7
Sphaerospora 10 species.....	1	..	2	1	2	2	2
Sinuolinea* 5 species.....	2	..	2	1	..
Myxidium 26 species.....	1	..	2	2	2	9	10
Sphaeromyxa 7 species.....	5	2
Zschokkella 4 species.....	1	1	..	1	1
Myxosoma 3 species.....	3	..
Lentospora 6 species.....	2	2	2
Myxobolus 63 species.....	54	9
Henneguya 32 species.....	1	1	25	5
Hoferellus 1 species.....	1	..

* These three genera are unknown to Auerbach, except two species which were formerly placed in Leptotheca and Sphaerospora.

Parisi (1912) followed Auerbach in his paper dealing with Myxosporidia from Italian waters. Poche (1913) put Auerbach's classification in better form as follows:

- Order: Myxosporidia
 - 2 Superfamily Mictosporea
 - 2 Family Myxidiidae
 - 2 Genus Zschokkella
 - 3 Genus Myxoproteus
 - 4 Genus Myxidium
 - 5 Genus Sphaeromyxa
 - 6 Genus Sphaerospora
 - 3 Family Chloromyxidae
 - 7 Genus Chloromyxum
 - 3 Superfamily Disporea
 - 4 Family Ceratomyxidae
 - 8 Genus Ceratomyxa
 - 9 Genus Leptotheca
 - 4 Superfamily Polysporea
 - 5 Family Myxosomatidae (Poche)
 - 10 Genus Myxosoma
 - 11 Genus Lentospora
 - 6 Family Myxobolidae
 - 12 Genus Myxobolus
 - 13 Genus Henneguya
 - 14 Genus Hoferellus

For the same reason given in discussing Auerbach, this, however, is not conformable with the present state of knowledge regarding these protozoa.

It was not until 1917 that the classification of the Myxosporidia approached to a more natural state in the valuable work by Davis (1917: 219-221). He pointed out sharply the unsatisfactory features in Doflein's classification and proposed a different system as follows:

- Order: Myxosporidia.
 - Suborder I Myxosporea Davis
 - Family 1 Ceratomyxidae
 - Genus 1 Leptotheca
 - Genus 2 Ceratomyxa
 - Family 2 Sphaerosporidae Davis
 - Genus 1 Myxoproteus
 - Genus 2 Sphaerospora
 - Genus 3 Sinuolinea
 - Family 3 Myxidiidae
 - Genus 1 Myxidium
 - Genus 2 Sphaeromyxa
 - Genus 3 Zschokkella
 - Family 4 Chloromyxidae
 - Genus 1 Chloromyxum

Suborder II Cystosporea Davis

Family 1 Myxosomidae* Davis

Genus 1 Myxosoma

Genus 2 Lentospora

Family 2 Myxobolidae

Genus 1 Myxobolus

Genus 2 Henneguya

Genus 3 Hoferellus

Thus, Davis selected the form of the spore for the establishment of two suborders and further rearranged the genera into closer positions to show relationship to each other better than any one of the previous authors. He, however, named the suborders according to a secondary character, i.e., the seat of the parasites in the host. According to his definition the trophozoites of the species belonging to Myxosporea are "with few exceptions free living in the body-cavity," while those of Cystosporea "with few exceptions" are tissue parasites.

From TABLE III on page 45, are taken the following data regarding this point:

	Total number of species known	Number of species found in body cavity	Number of species found in tissue	Number of species found in both places	Seat unknown
Myxosporea.....	132	114	14	2	2
Cystosporea.....	105	7	95	1	2

Thus it appears that the terms Myxosporea and Cystosporea do not seem to be properly used. These may be replaced by terms that denote the first and common character of the suborders.

The suggestions as to the adoption of other characters than the spore for the divisions of Myxosporidia, proposed by Awerinzew (1907:831; 1908:64), Auerbach (1910:161) and Davis (1917:217) can only be applied in the future. At the present time, the characters concerning the vegetative form do not appear to afford a better and more natural basis for the classification of Myxosporidia than those of the spore. Thus from the taxonomic point of view the present situation does not seem to be much improved as compared with that at the end of the last century.

The writer proposes in the following pages a new classification based on the characters of the spore.

* Davis did not notice the establishment of the family Myxosomatidae by F. Poche (1913), including exactly the same genera. See page 54.

Order MYXOSPORIDIA Bütschli 1881

Suborder EURYSPOREA nom. nov.

Largest diameter of the spore at right angles to the sutural plane. One polar capsule on each side of the plane. Sporoplasm with no iodophilous vacuole. Vegetative form found in body cavity (except 2 species). Great majority parasites of marine fish. Monosporous, disporous and polysporous.

Family CERATOMYXIDAE Doflein 1899

With the characters of the suborder.

Genus LEPTOTHECA Thélohan 1895

Shell-valves of spore hemispherical or shortly rounded. 15 species. Disporous (7 unknown). 14 species in body-cavity; 1 in tissue; all in marine fish. Type species: *Leptotheca agilis* Thélohan.

Genus CERATOMYXA Thélohan 1892

Shell-valves, conical and hollow, attached on the bases; free ends extended, tapering to more or less sharply pointed or rounded ends. Sporoplasm usually does not fill the cavity, but is located asymmetrically in it. 35 species. Disporous (23 species), monosporous and disporous (3 species), disporous and polysporous (4 species) and unknown (5 species). All (except 2 species in urinary bladder) in the gall-bladder of marine fish. Type species: *Ceratomyxa arcuata* Thélohan.

Genus MYXOPROTEUS Doflein 1898 emend. Davis 1917

Spores roughly pyramidal; with or without distinct processes from the base of the pyramid. 3 species. Disporous (one species unknown). All in urinary bladder of marine fish. Type species: *Myxoproteus ambiguus* (Thélohan) Doflein.

Genus WARDIA nov. gen.

Spore form of isosceles triangle with two convex sides. Oval in profile. Surface of shell with fine ridges which turn into fringe-like processes at the posterior end. The polar capsules, large and perfectly spherical, situated at the central portion of the spore, opening at the anterior tip. Two species. Polysporous (one species unknown). Tissue parasite (one species) of fresh-water fish and amphibia, both found in Illinois, U. S. A. Type species: *Wardia ovinocua* nov. spec.

Genus MITRASPORA Fujita 1912 emend. Kudo

Spores spherical or ovoidal. Two polar capsules pyriform, one situated on each side of the sutural plane. Shell longitudinally striated; with or without long and fine filaments projecting posteriorly in a row at right

angles to the sutural plane at the posterior side. 3 species. Disporous and polysporous. All found in kidney of fresh-water fish. Type species: *Mitraspora cyprini* Fujita.

Suborder SPHAEROSPORA nom. nov.

Spores spherical or subspherical, with two to four polar capsules. Sporoplasm without iodophilous vacuole. Vegetative form found in body-cavity and tissue. Monosporous, disporous and polysporous. Parasites of marine and fresh-water fish and amphibia.

Family CHLOROMYXIDAE Thélohan 1892*

Spores with four polar capsules. Monosporous, disporous and polysporous.

Genus CHLOROMYXUM Mingazzini 1890

With the characters of the family. 22 species. 18 in body cavity; 4 in tissue. 7 from marine and 12 from fresh-water fish, 2 in amphibia, 1 in insect. Type species: *Chloromyxum leydigi* Mingazzini.

Family SPHAEROSPORIDAE Davis 1917

Spores with two polar capsules. Monosporous, disporous and polysporous.

Genus SPHAEROSPORA Thélohan 1892

Spores with two polar capsules. Monosporous, disporous and polysporous. 10 species. Body-cavity and tissue. 5 from fresh-water and 5 marine fish. Type species: *Sphaerospora divergens* Thélohan.

Genus SINUOLINEA Davis 1917

Spores with or without lateral processes. Two polar capsules spherical. Sutural line sinuous. 5 species. Disporous and polysporous. In the urinary bladder of marine fish. Type species: *Sinuolinea dimorpha* Davis.

Suborder PLATYSPORA nom. nov.

Sutural plane of the spore coincides with or at an acute angle to the longest diameter. One or two polar capsules. Sporoplasm with or without an iodophilous vacuole.

Family MYXIDIIDAE Thélohan 1892

Two polar capsules, one at each end. Sporoplasm without any iodophilous vacuole. Spores fusiform.

* Thélohan (1892) used the terms: Chloromyxidées, Myxididées, Myxobolidées, which Gurley (1893) made over into Chloromyxidae, Myxidiidae, Myxobolidae, so that the credit of recognizing and establishing these families belongs to Thélohan.

Genus MYXIDIUM Bütschli 1882

Spores more or less regularly fusiform, with pointed or rounded ends. Polar filaments long and fine. 26 species. Monosporous, disporous and polysporous. 22 in body-cavity; 4 in tissue. 15 in marine and 6 in fresh-water fish, 2 in fishes from both waters and 3 in reptilia. Type species: *Myxidium lieberkühni* Bütschli.

Genus SPHAEROMYXA Thélohan 1892

Spores fusiform, with truncated ends. Polar filament short and thick. Trophozoites large and disc shaped. 7 species. Polysporous (2 unknown). 6 in body-cavity; 1 unknown. 6 in marine fish; 1 in amphibia. Type species: *Sphaeromyxa balbianii* Thélohan.

Genus ZSCHOKKELLA Auerbach 1910

Spores, semicircular in front view; pointed at ends. Polar capsules large and spherical, opening on the flat edge near the tips. Sutural line usually curved in S-form. 4 species. Monosporous, disporous and polysporous. Body-cavity. 2 from marine and 2 from fresh-water fish. Type species: *Zschokkella hildae* Auerbach.

Family MYXOSOMATIDAE Poche 1913

Two polar capsules at the anterior end. Sporoplasm without iodophilous vacuole.

Genus MYXOSOMA Thélohan 1892

Spores ovoidal, flattened and more or less elongated. 3 species. Polysporous. Tissue parasites. 2 in fresh-water and 1 in marine fish. Type species: *Myxosoma dujardini* Thélohan.

Genus LENTOSPORA Plehn 1905

Spores similar to Myxobolus in form. Sporoplasm without any iodophilous vacuole. 6 species. Disporous and polysporous (2 unknown). 1 in marine and 3 in fresh-water fish, 2 from fishes in both waters. Type species: *Lentospora cerebralis* (Hofer) Plehn.

Family MYXOBOLIDAE Thélohan 1892

Spores with one or two polar capsules at the anterior end, with or without posterior processes. Sporoplasm with an iodophilous vacuole. Majority polysporous in fresh-water fishes.

Genus MYXOBOLUS Bütschli 1882

Spores ovoidal or ellipsoidal; flattened. One or two polar capsules at the anterior end. Shell without posterior process. 63 species. Polysporous (9 species unknown). 59 species in tissue; 4 unknown. 5 in marine and 56 in fresh-water fish, 1 in annelid and 1 in amphibia. Type species: *Myxobolus mülleri* Bütschli.

Genus HENNEGUYA Thélohan 1892

Spores more or less globular or ovoidal. Two polar capsules at the anterior end. Posterior end of the shell-valves prolonged into more or less extended processes, which unite and form a tail in the median line. 32 species. Polysporous, disporous and monosporous. 28 species in tissue and 4 in body-cavity. In fresh-water fish, except one. Type species: *Henneguya psorospermica* Thélohan.

Genus HOFERELLUS Berg 1898

Spores pyramidal, with two posterior processes from the lateral faces. 1 species. Polysporous. Tissue and body-cavity of fresh-water fish. Type and only species: *Hoferellus cyprini* Doflein.

DESCRIPTIONS OF GENERA AND SPECIES

Suborder EURYSPOREA nom. nov.

The definition of the suborder is recorded on page 56.

Family CERATOMYXIDAE Doflein

1899	<i>Ceratomyxidea</i>	Doflein	1899 : 378
1901	<i>Ceratomyxidae</i>	Doflein	1901 : 182

The characters of the family are described on page 56.

Genus LEPTOTHECA Thélohan

1895	<i>Leptotheca</i>	Thélohan	1895 : 331
------	-------------------	----------	------------

The characters of the genus are described on page 56.

Type species: *Leptotheca agilis* Thélohan.

LEPTOTHECA AGILIS Thélohan

[Figs. 1 to 5]

1892	<i>Ceratomyxa agilis</i>	Thélohan	1892 : 962
1895	<i>Leptotheca agilis</i>	Thélohan	1895 : 332
1898	<i>Leptotheca agilis</i>	Doflein	1898 : 294, 297

Habitat: Gall-bladder of *Trygon pastinaca* L. and *Scorpaena* sp.; France, Rovigno, Napoli.

Vegetative form: Form generally elongated. Anterior end rounded where a mass of fat globules is found, while the posterior end terminates in a point. Size not exceeding 85μ by 20 to 25μ . The posterior part is sometimes divided into a certain number of lobes. In the protoplasm, the globules are clearly seen. Pseudopodia are localized at the anterior portion of the body. They are long, 40 to 50μ in length, filiform and very active in moving from back toward front, just like the motion of oars. Disporous.

Spore: Slightly elongated. Dimensions: sutural diameter 6 to 7μ , breadth 11 to 12μ .

LEPTOTHECA ELONGATA Thélohan

[Fig. 6]

1895	<i>Leptotheca elongata</i>	Thélohan	1895 : 332
1898	<i>Leptotheca elongata</i>	Doflein	1898 : 312
1917	<i>Leptotheca elongata</i>	Georgévitch	1917b : 99-106

Habitat: Gallbladder of *Merluccius merluccius* L. (*M. vulgaris*) and *Motella tricirrata*; Marseille, Banyuls, Le Croisic, Napoli, Monaco.

Vegetative form: Form variable. Many individuals show, however, a very characteristic form. It is elongated and has the length of about 120μ . The anterior end is enlarged into a disc-shaped depression, on the edge of which, the branched pseudopodia are formed. The body gradually narrows itself toward the posterior end. Also club-shaped, etc. The short lobose pseudopodia show no movement like that of oars.

Georgévitch's form: Young forms, oval or rounded, are attached to the epithelial cells of the bladder with a long filiform pseudopodium at the free end. Such forms often agglomerate in great number.

Spore: Form similar to the spore of *Leptotheca agilis*. Dimensions on the average: Sutural diameter 12 to 15μ , breadth 18 to 20μ .

LEPTOTHECA POLYMORPHA (Thélohan) Labbé

1895	<i>Leptotheca elongata</i>	Thélohan	1895 : 332-333
1899	<i>Leptotheca polymorpha</i>	Labbé	1899 : 88

Habitat: Gall-bladder of *Phycis mediterraneus* (*P. phycis* L.); Banyuls.

Vegetative form: Form extremely polymorphous, with three main types. 1) Somewhat regularly club-shaped, with lobose pseudopodia, sometimes filiform at one end. 2) Irregular as is the case with *Ceratomyxa truncata*, with long (25μ) ectoplasmic processes, which are motionless or very slow in motion. Lobose pseudopodia are formed actively. 3) More or less rounded with bristle-like filose pseudopodia. Intermediary forms are also found. Often many individuals unite together. The protoplasm is much different from other forms, i.e., more homogeneous and compact. Granules are hardly visible on account of vacuolar appearance.

Spore: Dimensions: sutural diameter 10 to 12μ , breadth 18 to 20μ , length of polar filament 40μ .

LEPTOTHECA PARVA Thélohan

[Fig. 7]

1895	<i>Leptotheca parva</i>	Thélohan	1895 : 333
1912	<i>Leptotheca parva</i>	Auerbach	1912 : 42-43

Habitat: Gall-bladder of *Scomber scombrus* L.; Marseille, Le Croisic, Le Vivier-sur-mer, Kristiansund, Stavanger, Bergen.

Vegetative form: Form ordinarily rounded, spherical or subspherical. Often club-shaped. Size not larger than 12 to 15μ in diameter. Protoplasm finely granular. Pseudopodia lobose.

Spore: Small, more or less elongated, curved in arch-form. Dimensions: sutural diameter 3 to 4μ , breadth 8 to 10μ .

LEPTOTHECA RENICOLA Thélohan

1895	<i>Leptotheca renicola</i>	Thélohan	1895 : 333
------	----------------------------	----------	------------

Habitat: Urinary tubules of the kidney of *Scomber scombrus* L.; Marseille, Le Croisic.

Vegetative form: Small. No marked character.

Spore: Globular. Form similar to the spore of *Sphaerospora*. Dimensions: sutural diameter 8μ , breadth 10μ .

LEPTOTHECA HEPSETI Thélohan

1895	<i>Leptotheca hepseti</i>	Thélohan	1895 : 334
------	---------------------------	----------	------------

Habitat: Gall-bladder of *Atherina hepsetus* L.; Marseille. Of rare occurrence; Thélohan observed it but once.

Vegetative form: Not described.

Spore: Form triangular with rounded angles. Dimensions: sutural diameter 7 to 8μ , breadth 12 to 15μ .

LEPTOTHECA PERLATA (Gurley) Labbé

[Fig. 8]

1883		Balbani 1883 : 201, 204
1894	<i>Chloromyxum (Sphaerospora) perlatum</i>	Gurley 1894 : 272
1899	<i>Leptotheca perlata</i>	Labbé 1899 : 88

Habitat: *Acerina cernua* L.; France (?).

Vegetative form: Not described.

Spore: Elliptic. Two small polar capsules converging. Dimensions not given.

LEPTOTHECA sp. Awerinzew

[Figs. 16, 17]

1908	<i>Leptotheca</i> sp.	Awerinzew	1908 : 51, 52
------	-----------------------	-----------	---------------

Habitat: Gall-bladder of *Sebastes norvegicus*; Eastern Finmark?

Vegetative form: Rounded form with clear differentiation of protoplasm into ectoplasm and endoplasm. Plasmotomy occurs.

Spore: Undescribed. No figure.

LEPTOTHECA MACROSPORA Auerbach

[Fig. 9]

1909	<i>Leptotheca macrospora</i>	Auerbach	1909 : 70-71
1910	<i>Leptotheca macrospora</i>	Auerbach	1910b : 768-769
1910	<i>Leptotheca macrospora</i>	Auerbach	1910c : 167
1912	<i>Leptotheca macrospora</i>	Auerbach	1912 : 42-43

Habitat: Gall-bladder of *Sebastes viviparus* H. Kr. and *S. dactylopterus*; Bergen, Kristiansund (May, September).

Vegetative form: Trophozoites spherical with the average diameter of 26 to 30μ . Homogeneous ectoplasm layer exhibits somewhat active ameboid movements. Endoplasm, granular in living specimen, is rather sharply distinguishable from the ectoplasm and contains large nuclei.

Spore: Size large. Form resembles to that of *Leptotheca parva*. Dimensions: sutural diameter and thickness 13μ , breadth 26μ . Polar capsules short oval, with a length of 5.2μ , length of polar filament about 130μ (KOH). In the second host, a few normal and numerous abnormal spores with three or four polar capsules were observed.

LEPTOTHECA INFORMIS Auerbach

[Fig. 10]

1910	<i>Leptotheca informis</i>	Auerbach	1910b : 770-771
1912	<i>Leptotheca informis</i>	Auerbach	1912 : 42-44

Habitat: Gall-bladder of *Molva vulgaris* Flem. and *Gadus merlangus*; Bergen, Tjomo.

Vegetative form: Young trophozoites with somewhat long and narrow pseudopodia formed of hyaline ectoplasm; movements active. The protoplasm is differentiated into ectoplasm and endoplasm. When stained, two large (7 to 9μ) and two small (3 to 4μ) nuclei were observed in an individual, 27μ long excluding the pseudopodia. Sporulating trophozoites are generally round and each forms two spores, which are developed independently to each other (i.e., not in ordinary pansporoblast). Auerbach observed centrosomes in the nuclei of large type in division. Disporous.

Spore: Large and heavily built. Greatly curved. Sutural line fairly well marked. Polar capsules round. Dimensions: sutural diameter 10μ , breadth 18 to 20μ , thickness 9μ , diameter of polar capsules 3 to 4μ . Sporeplasm contains two nuclei, 3.5 to 4μ in diameter.

LEPTOTHECA LONGIPES Auerbach

[Fig. 11]

1910	<i>Leptotheca longipes</i>	Auerbach	1910b : 771
1912	<i>Leptotheca longipes</i>	Auerbach	1912 : 42-43

Habitat: Gall-bladder of *Brosmius brosmæ* Asc.; Bergen (May).

Vegetative form: Trophozoites elongated or rounded. Only few pseudopodia which are very long. Small forms with a very long process, were observed in large numbers; length of the body being 10μ , while the process was 60μ long. Endoplasm contains nuclei of various sizes. Disporous.

Spore: Form similar to that of *Leptotheca informis*, though smaller. Dimensions: sutural diameter 8 to 9μ , breadth 12 to 14μ , thickness 8μ , diameter of polar capsule 2.5μ .

LEPTOTHECA FUSIFORMIS Davis

[Fig. 12]

1917	<i>Leptotheca fusiformis</i>	Davis	1917 : 222
------	------------------------------	-------	------------

Habitat: Gall-bladder of *Cestracion zygaena*; Beaufort (July).

Vegetative form: Pyriform, tapering gradually toward the posterior end, which usually terminates in a long, slender process; colorless and transparent. Progressive movements rapid. Endoplasm granular, the granules being more abundant at the anterior end. The average size of full-grown individuals: 50μ by 13μ . Disporous.

Spore: Elliptical in front view; fusiform in side view. Sutural plane slightly oblique to the longest diameter, the line forming a marked ridge. Polar capsules open on opposite sides of the spore. Sporoplasm finely granular, confined to the central part of spore. Dimensions: sutural diameter 9μ , breadth 16μ , polar capsule 4.5μ long, length of polar filament 30μ .

LEPTOTHECA SCISSURA Davis

[Fig. 13]

1917 *Leptotheca scissura* Davis 1917 : 222

Habitat: Gall-bladder of *Dasybatis hastatus* and *Pteroplatea maclura* Le Sue; Beaufort (July, August).

Vegetative form: Young form elongated, with long attenuated posterior process; usually slightly constricted just posterior to rounded anterior end, which bears numerous long, filiform pseudopodia. Progressive movement rapid. Ectoplasm distinguishable at the anterior end. Endoplasm usually filled with small, clear, colorless spherules, which become larger and yellowish as the body increases in size. Each spherule contains one to several dark-brown granules, which increase in size and number and finally collect in an irregular clump at the centre of spherule. The larger individuals are usually flattened dorso-ventrally. The posterior end is divided into long slender processes, presenting sometimes a network caused by the fusion of two or more adjacent processes. Full-grown forms: length 125 to 150μ , breadth 20 to 25μ . The longest observed, 195μ by 16μ . Disporous.

Spore: Elliptical in front view; somewhat flattened along the posterior side. Sutural line distinct and at right angles to the longest diameter. Polar capsules have foramina at some distance from the capsular margin. Sporoplasm finely granular, nearly filling both valves. Dimensions: sutural diameter 11μ , breadth 22μ , diameter of polar capsule 4μ .

LEPTOTHECA LOBOSA Davis

[Fig. 14]

1917 *Leptotheca lobosa* Davis 1917 : 223

Habitat: Urinary bladder of *Paralichthys dentatus* L.; Beaufort (July).

Vegetative form: Usually spherical which may form a large rounded pseudopodium composed of ectoplasm. Body colorless and transparent to translucent. Ameboid movements very slow. Ectoplasm contains coarse granules, which are of uniform size and very distinct. Endoplasm

less granular and more transparent than ectoplasm, containing numerous large, yellow, fat globules, which are abundant in large forms. Diameter up to 24μ . Disporous.

Spore: Elliptical in front view; valves slightly tapering but rarely alike. Sutural line forming a sinuous ridge. Polar capsules open at some distance from the anterior margin. Sporoplasm nearly filling both valves. Free spores are often seen to remain united at the sutural line. Dimensions: sutural diameter 9 to 10μ , breadth 16 to 18μ ; diameter of polar capsule 3μ .

LEPTOTHECA GLOMEROSA Davis

[Fig. 15]

1917 *Leptotheca glomerosa* Davis 1917 : 223

Habitat: Urinary bladder of *Paralichthys albiguttus*; Beaufort.

Vegetative form: Rounded or somewhat irregular in shape, with short lobose pseudopodia. Body transparent and colorless. Ameboid movements slow. Ectoplasm hyaline, forming a distinct outer layer. Endoplasm finely granular, with numerous small fat globules varying in size. Almost entire body is used for spore formation. Diameter of rounded sporulating trophozoite about 11μ . Disporous.

Spore: Approximately cylindrical; valves rounded at ends. The coiled polar filament not visible in the polar capsule. Sutural line at right angles to the longest diameter. Sporoplasm finely granular, fills the extracapsular cavity of spore. Dimensions: sutural diameter 4.5μ , breadth 9μ , diameter of polar capsule 2μ .

Genus CERATOMYXA Thélohan

1892 *Ceratomyxa* Thélohan 1892 : 169, 171, 175
1895 *Ceratomyxa* Thélohan 1895 : 334

The characters of the genus described on page 56.

Type species: *Ceratomyxa arcuata* Thélohan.

CERATOMYXA ARCUATA Thélohan

[Figs. 18 to 22]

1892 *Ceratomyxa arcuata* Thélohan 1892a : 1091
1895 *Ceratomyxa arcuata* Thélohan 1895 : 335-336
1899 *Ceratomyxa arcuata* Labbé 1899 : 90
1912 *Ceratomyxa arcuata* Parisi 1912 : 290-291
1913 *Ceratomyxa arcuata* Jameson 1913 : 2
1916 *Ceratomyxa arcuata* Georgévitch 1916a : 3

Habitat: Gall-bladder of *Pagellus centrodontus* C. et V., *Crenilabrus melops* L., *Motella tricirrata* Bl., *Ophidium vasalli*, *Gobius paganellus* L., *Heliases chromis* Gthr.; *Scorpaena scrofa* L., *S. porcus* L.; France (Marseille, Banyuls, Concarneau, Roscoff), Italy (Napoli, summer), Monaco (May).

Vegetative form: Polymorphous; generally club-shape, pseudopodia localized at the broad end; the other end cylindrical or terminating in a sharp point. Some other different forms. Pseudopodia, always localized, lobose pointed at the extremities. Ectoplasm hyaline and thin. Endoplasm contains fat globules and particular elements, mostly large refractive globules, which seem to disappear in the sporulating individuals. Dimensions (maximum): length 35 to 40 μ , breadth 12 to 15 μ , pseudopodia about 10 μ long. Disporous.

Spore: Arch form. Shell valves equal. Sporoplasm occupies the extracapsular cavity of the spore. The length varies rather considerably according to the development of the lateral processes, which are occasionally acuminate or very short. Often extremities are rounded. Dimensions (Thélohan); breadth 20 to 30 μ , sutural diameter 5 to 8 μ . Parisi's measurements: breadth 25 to 31 μ , sutural diameter 5.5 to 6 μ , length of polar capsules 3.5 to 4 μ , length of polar filaments 25 μ .

Remarks: The writer agrees with Parisi in eliminating Labbé's two subspecies (1899:90), as they are too arbitrary.

CERATOMYXA SPHAERULOSA Thélohan

[Figs. 23 to 24]

1892	<i>Ceratomyxa sphaerulosa</i>	Thélohan	1892 : 171
1895	<i>Ceratomyxa sphaerulosa</i>	Thélohan	1895 : 334-335
1909	? <i>Ceratomyxa sphaerulosa</i>	Auerbach	1909 : 80
1912	<i>Ceratomyxa sphaerulosa</i>	Auerbach	1912 : 4, 45
1916	<i>Ceratomyxa sphaerulosa</i>	Georgévitch	1916a : 3

Habitat: Gall-bladder of *Mustelus canis* Mitch. (*M. vulgaris*), *Galeus galeus* L. (*G. canis*), *Clupea harengus*, *Scullium canicula* Cuv.; St-Valery-en-Caux, Roscoff, Bergen, Monaco (May).

Vegetative form: Form more or less definite among the adults. Generally elongated. Both ends slightly attenuated. Wide in the middle part of the body. Lobose pseudopodia at one of the extremities. Others more massive or more or less regularly spherical, in which case the pseudopodia are formed from the whole surface. Spherical form does not exceed 50 to 60 μ in diameter. Other forms 90 to 100 μ by 30 to 40 μ (largest). Young forms colorless and are more variable than the adults. Protoplasm homogeneous and finely granular. Adult form, on the contrary, yellowish or greenish yellow. The endoplasm is filled with small (5 μ in diameter) spheres, in the centre of which 5 to 6 small granules, yellowish brown or greenish in color, are present. Disporous.

Spore: Remarkably large. Polar filament can be seen *in vivo* (easily extruded by KOH, ether, etc.) Sporoplasm occupies one of the shell-valves, while a small mass of very pale looking substance is seen in the other. Dimensions: sutural diameter 10 to 12 μ , breadth 90 to 100 μ , subspherical polar capsule 6 to 7 by 5 μ , sporoplasm 12 to 15 by 8 to 9 μ .

CERATOMYXA PALLIDA Thélohan

1895	<i>Ceratomyxa pallida</i>	Thélohan	1895 : 336-337
1898	<i>Ceratomyxa pallida</i>	Doflein	1898 : 341
1916	<i>Ceratomyxa pallida</i>	Georgévitch	1916b : 2, 3

Habitat: Gall-bladder of *Box boops* L. and *B. salpa* L.; Marseille, Villefranche, Rovigno, Monaco (May).

Vegetative form: Ordinarily spherical not exceeding 16 to 20 μ in diameter. Many individuals often found in massive groups. Pseudopodia lobose and mostly short. Protoplasm extremely pale with fine granules.

Spore: Dimensions: sutural diameter 5 μ , breadth 25 to 30 μ .

CERATOMYXA GLOBULIFERA Thélohan

[Fig. 25]

1895	<i>Ceratomyxa globulifera</i>	Thélohan	1895 : 338
------	-------------------------------	----------	------------

Habitat: Gall-bladder of *Merluccius merluccius* L. (*M. vulgaris*); Marseille, Banyuls.

Vegetative form: Polymorphous. Elongated into long branches, including endoplasm. Endoplasm contains small refractive globules.

Spore: Elongated. Shell-valves unequal, one being longer and finer than the other. Dimensions: sutural diameter 10 μ , breadth 50 μ .

CERATOMYXA APPENDICULATA Thélohan

[Fig. 26]

1892	<i>Ceratomyxa appendiculata</i>	Thélohan	1892a : 963-964
1895	<i>Ceratomyxa appendiculata</i>	Thélohan	1895 : 337
1898	<i>Ceratomyxa appendiculata</i>	Doflein	1898 : 300, 311

Habitat: Gall-bladder of *Lophius piscatorius* L., *L. budegassa* Spin.; Roscoff, Le Croisic, Marseille, Banyuls, Napoli, Rovigno.

Vegetative form: Extremely polymorphous. Young form spherical, spatulaform, club-shape, etc. In adult form, the main thick part of the body, in which spore formation takes place, forms 1 to 6 long prolongations, twice or three times longer than the main part of the body. Pseudopodia lobose, filiform or elongated with enlargements. Disporous.

Spore: Lateral prolongations of shell-valves well developed. Dimensions: sutural diameter 5 to 7 μ , breadth 50 μ .

CERATOMYXA TRUNCATA Thélohan

[Fig. 27]

1895	<i>Ceratomyxa truncata</i>	Thélohan	1895 : 336
1912	<i>Ceratomyxa truncata</i>	Parisi	1912 : 289-290

Habitat: Gall-bladder of *Clupea pilchardus* Walb. (*Alosa sardina*); Marseille, Villefranche, Napoli (August, September).

Vegetative form: Polymorphous. Ordinarily more or less rounded, with lobose pseudopodia. Pseudopodia long and often shows very active movements. Endoplasm very finely granular, contains small fat globules which are found in irregular mass or in a circular form. Disporous.

Spore: Valves are short and truncate. Sporoplasm occupies the whole cavity. Spores with three valves are frequently encountered. Dimensions: breadth 25μ , 5μ in sutural diameter. According to Parisi, spores with two shell-valves are rather few (10%), while those with three (70%) and four shell-valves (20%) prevail in number! Dimensions: breadth 20 to 30μ , length of the polar filament 45μ .

CERATOMYXA RETICULARIS Thélohan

[Fig. 28]

1895 *Ceratomyxa reticularis* Thélohan 1895 : 337-338

Habitat: Gall-bladder of *Trachinus draco* L.; Banyuls.

Vegetative form: Extremely polymorphous. Generally spherical or club-shaped. Well developed trophozoites have the similar form as in *C. appendiculata*. Endoplasm highly reticular, with refringent fluid.

Spore: Shell valves are short and truncate, one of which is curved to the rear. Dimensions: sutural diameter 12 to 15μ , breadth 45 to 50μ .

CERATOMYXA INAEQUALIS Doflein

[Fig. 29]

1898 *Ceratomyxa inaequalis* Doflein 1898 : 284-285

Habitat: Gall-bladder of *Crenilabrus mediterraneus* and *C. pavo*; Napoli.

Vegetative form: Form usually club-shaped. Protoplasm in active motion, is differentiated distinctly into ectoplasm and endoplasm. Body yellowish brown by the presence of granules in endoplasm. Inactive formation of pseudopodia. Ameboid movements or progressive movements by means of the posterior process. Size: 20 to 40μ by 5 to 10μ in average. Length of the posterior process up to 30μ . After spore formation, only two nuclei remain in protoplasm, which apparently degenerate later. Disporous.

Spore: Elliptical, somewhat flattened. Massive. Very transparent. Both ends round, but unequally built, i.e., one end is club-shaped. Polar capsules are somewhat round and are bound to the shell by protoplasmic bridges. The polar filament is not seen in fresh spores. Dimensions: sutural diameter 6μ , breadth 31μ , diameter of the polar capsule 2.5 to 3μ , length of polar filament is half breadth of the spore (diluted nitric acid).

CERATOMYXA LINOSPORA Doflein

[Figs. 30 to 31]

1898 *Ceratomyxa linospora* Doflein 1898 : 285Habitat: Gall-bladder of *Labrus turdus*; Napoli.

Vegetative form: Club- or spindleshape. Protoplasm highly granulated. Body whitish grey, though very transparent. Pseudopodia very fine and only formed at the anterior end of the body. Size: 30 to 35 μ by 16 to 18 μ . Disporous.

Spore: Form symmetrical with long thread-like lateral processes. In sporoblast, the processes are wound around the spore. It is twice as long as the breadth of the spore. Polar capsules large and spherical pyriform. Dimensions: total breadth 50 μ , breadth of the main part of the spore 10 to 12 μ , sutural diameter 5 μ , length of lateral process 20 μ . "Polar filament was too fine to be measured."

CERATOMYXA RAMOSA Awerinzew

[Figs. 32 and 33]

1907	<i>Ceratomyxa ramosa</i>	Awerinzew	1907 : 831-834
*1908	<i>Ceratomyxa ramosa</i>	Awerinzew	1908 : 60-66

Habitat: Gall-bladder of *Hippoglossus vulgaris* Flemm.; Kjellebjord, Murman coast.

Vegetative form: Form irregular ameboid, owing to the presence of peculiar pseudopodia. The middle part of the body is enlarged into an ellipsoidal form, where nuclei and sporoblasts are present. From this part two, rarely one or three processes are formed, which branch out several pseudopodia of different length. The finer portions of pseudopodia anastomose each other and form a characteristic and remarkable network. Differentiation of protoplasm is not very distinct. Ectoplasm is not well developed, tho covering the entire surface of the body as a thin layer. Endoplasm slightly vacuolated and granular, forms the greater part of the body. Disporous and polysporous.

Spore: Form and size (?) resemble *C. arcuata*. Slightly curved toward the posterior side. Valves usually unequally built, one being longer than the other. Sporoplasm almost always asymmetrically situated in the shell. Polar capsules on each side of the sutural plane and of the plane perpendicular to the sutural plane, cutting the spore into two equal parts. Young spores in development ellipsoidal to kidney bean shape. Dimensions: sutural diameter 12 to 20 μ , breadth 50 to 80 μ .

*Professor J. Zeitlin has kindly translated some part of the paper, for which the writer expresses his thanks.

CERATOMYXA DREPANOPSETTAE Awerinzew

[Figs. 34 to 39]

1907	<i>Ceratomyxa</i> sp.	Awerinzew	1907 : 832-833
1908	<i>Ceratomyxa drepanopsettae</i>	Awerinzew	1908 : 1-41, 45-47
1909	<i>Ceratomyxa drepanopsettae</i>	Awerinzew	1909 : 74-112
1912	<i>Ceratomyxa drepanopsettae</i>	Auerbach	1912 : 44-45
1918	<i>Ceratomyxa drepanopsettae</i>	Kudo	1918 : 14-15

Habitat: Gall-bladder of *Pleuronectes platessa*, *P. flesus*, *Drepanopsetta plattessoides*, *Hippoglossus vulgaris*, *Hippoglossoides limandoides* and *Paralichthys dentatus*; Murmankuste, Kabelvaag, Rorvik, Tjomo, Woods Hole (August, September).

Vegetative form: Polymorphous. Usually very much elongated and slender forms. Protoplasm differentiated. Endoplasm coarsely granular. Pseudopodia lobose and filiform (2 to 3 μ), with which the trophozoites attach themselves to the epithelium of the bladder. Disporous.

Spore: Curved toward the posterior side. Shell with rounded ends. Valves almost always unequally built. Dimensions: breadth 50 to 80 μ . Auerbach's form: Form variable. Size: sutural diameter about 12 to 14 μ , breadth about 56 μ , diameter of polar capsule about 4 to 6 μ , length of the cavity in which the sporoplasm is located about 34 μ . Kudo's form: Variable. Sutural diameter 8 to 10 μ , average breadth 64 μ , diameter of polar capsule 6 μ .

CERATOMYXA TYLOSURI Awerinzew

[Figs. 40 and 41]

1913	<i>Ceratomyxa tylosuri</i>	Awerinzew	1913a : 153
------	----------------------------	-----------	-------------

Habitat: Gall-bladder of *Tylosurus schismatorhynchus*; Lorenço Marques, Delagoa Bay (Africa).

Vegetative form: Large, irregular, disc-like or large ameboid, with blunt lobose pseudopodia and highly granular protoplasm.

Spore: Large. The anterior edge arch-shape, while the posterior edge has two small horns which are located symmetrically to the sutural line. Polar capsules elongated and are separated from binuclear sporoplasm by a special membrane. Rarely spore with three polar capsules. Dimensions breadth 124 to 140 μ , sutural diameter 40 to 45 μ , thickness 25 to 30 μ .

CERATOMYXA (?) SPARI Awerinzew

[Figs. 42 and 43]

1913	<i>Ceratomyxa</i> (?) <i>spari</i>	Awerinzew	1913a : 153-154
------	------------------------------------	-----------	-----------------

Habitat: Gall-bladder of *Sparus berda*; Lorenço Marques, Delagoa Bay (Africa).

Vegetative form: Large (100 to 120 μ), disc-form ameboid, containing a large number of enclosures and granules of different size. In one case, a number of this form, carrying no spore, underwent budding, which resulted in forming spherical forms of various size, some of which divided again into 2 to 6 parts (Plasmotomy?). Monosporous and disporous.

Spore: More or less curved. Two polar capsules lie closely together on each side of the sutural plane. Ends of shell-valves are rounded. Dimensions: breadth 50 to 60 μ , sutural diameter 12 to 15 μ , thickness 12 to 15 μ , polar filament very long (length not given).

Remarks: Awerinzew thinks this is the intermediate form between *Leptotheca* and *Ceratomyxa*.

CERATOMYXA sp. (?) Awerinzew

1913 *Ceratomyxa* sp. (?) Awerinzew 1913a : 154

Habitat: Gall-bladder of *Scatophagus argus*; Delagoa Bay (Africa).

Vegetative form: Small, disc-form ameboid (25 to 35 μ), containing two spores of indistinct contour, on account of incomplete formation of the shell. Two spores, apparently, developed in one pansporoblast. Disporous.

Spore: Form could not exactly be made out. Polar capsules were arranged like those of other *Ceratomyxa*.

CERATOMYXA sp. (?) Awerinzew

1913 *Ceratomyxa* sp. (?) Awerinzew 1913a : 154-155

Habitat: Gall-bladder of *Rhinobathus* Awer. (?); Lorenço Marques (Africa).

Vegetative form: Irregular shape. Endoplasm highly granular. In the epithelial layer of the gall-bladder numerous, spherical cysts (30 to 35 μ) were found. Two spores are formed in one pansporoblast. Disporous.

Spore: Cylindrical with broad and slightly rounded ends. Dimensions: sutural diameter 16 to 19 μ , breadth 70 to 80 μ , thickness 16 to 19 μ .

CERATOMYXA ACADIENSIS Mavor

[Figs. 44 to 47]

1915 *Ceratomyxa acadiensis* Mavor 1915 : 27-30
1916 *Ceratomyxa acadiensis* Mavor 1916 : 551-574

Habitat: Gall-bladder of *Urophycis chuss* (trophozoites are attached to undetermined Myxosporidia, see p. 176), *Zoarces angularis*, *Pseudopleuronectes americanus*; New Brunswick (Canada) (July to September).

Vegetative form: Polymorphous. Typically club-shaped with very long tail, or irregularly stellate. Pseudopodia show rigidity. Sometimes

clumps of protoplasm along their length, which are connected by thin hyaline filaments of ectoplasm. Differentiation of protoplasm is usually observable at the anterior region. Dimensions: length, excluding tail, 12 to 15 μ , breadth 10 to 20 μ , tail up to 60 μ . Disporous.

Spore: Wide, short and slightly compressed dorso-ventrally, with very long fine lateral filaments. Polar capsules spherical. Polar filament invisible *in vivo*. Dimensions: breadth 40 to 50 μ , sutural diameter 7 to 8 μ , diameter of polar capsule 3 to 4 μ , length of polar filament 70 μ , length of lateral filaments 250 to 300 μ .

CERATOMYXA sp. Georgévitch

1916	<i>Ceratomyxa</i> sp.	Georgévitch	1916a : 3
------	-----------------------	-------------	-----------

Habitat: Gall-bladder of *Muraena* sp.; Monaco (May).

Vegetative form: No description.

Spore: No description. No figure.

CERATOMYXA CORIS Georgévitch

[Fig. 48]

1916	<i>Ceratomyxa coris</i>	Georgévitch	1916a : 4-5
1917	<i>Ceratomyxa coris</i>	Georgévitch	1917a : 1-20

Habitat: Gall-bladder of *Coris julius*, *C. giofredi*; Villefranche (March, June).

Vegetative form: Various forms, club-shape, spherical or elongated, with lobose or filiform pseudopodia. Disporous and rarely Polysporous.

Spore: More or less ellipsoidal. Lateral prolongations of the shell-valves short and truncate. Sutural line straight. Sporoplasm, elongate, rounded, elliptical, fills a part of the extracapsular cavity of the spore. Polar capsules rounded, almost spherical, not converging. Dimensions not given.

Remarks: Georgévitch observed (1917: Fig. 30) that spores of *Glugea marionis* occurred in disporous trophozoite of *Ceratomyxa coris*, which he thought to have happened accidentally by plasmogamy of these two Cnidosporidia. The above mentioned figure, however, strongly suggests that *G. marionis* may be leading parasitic life in the trophozoite of *C. coris*.

CERATOMYXA HEROUARDI Georgévitch

[Fig. 49]

1913	<i>Leptotheca</i> (?) sp.	Jameson	1913 : 2
1916	<i>Ceratomyxa herouardi</i>	Georgévitch	1916a : 5-8
1916	<i>Ceratomyxa herouardi</i>	Georgévitch	1916b : 717-19, 983-985
1917	<i>Ceratomyxa herouardi</i>	Georgévitch	1917 : 375-399

Habitat: Gall-bladder of *Box Salpa* L.; Monaco (May).

Vegetative form: Polymorphous. Elongated with same breadth or tapering to one end; club-shaped with roundish enlargements. Young trophozoites spherical or pyriform. Pseudopodia long and narrow or broad and bi- or multi-lobate. Body colorless both in the young and the adult. Protoplasm homogeneous and finely granular. Disporous and polysporous. Spores are found inside of the endoplasm and in the roundish buds, ordinarily two spores being formed in each bud. Number of buds on one trophozoite varies. Plasmotomy by budding and division.

Spore: Elongated elliptic. Polar capsules spherical and large. Sutural plane cuts the spore into exactly equal two parts. Two nuclei in sporoplasm are rather small and are always in one of the shell-valves. Dimensions not given.

Remark: The form, mentioned by Jameson in the same seat, host and locality, that "has something of the appearance of a *Leptotheca*" and that is also "almost certainly neither of the two Myxosporidia—*Ceratomyxa pallida* and *Henneguya neapolitana* . . .," is probably identical with the present form.

CERATOMYXA MESOSPORA Davis

[Fig. 50]

1917 *Ceratomyxa mesospora* Davis 1917 : 223-224

Habitat: Gall-bladder of *Cestracion zygaena*, *C. tiburo*; Beaufort (July).

Vegetative form: Pyriform, elongate, with long, slender posterior process. Numerous filiform pseudopodia at anterior end. Progressive movements rapid. Body colorless. No sharp demarcation between ectoplasm and endoplasm. Endoplasm finely granular and filled with small, colorless, homogeneous spherules. Spherules absent at anterior end. Size: total length 70 to 85 μ , length exclusive of posterior process 50 to 75 μ , breadth 20 to 25 μ . Disporous.

Spore: Greatly elongate, each valve forming a slightly tapering cone, rounded at the apex. Valves not compressed. Sutural plane forming an acute angle with the longest diameter. Polar capsules conspicuous. Coiled polar filaments very distinct. Polar capsules are remarkable in that they are asymmetrically situated, one being always located in the widest part of the spore, while the other being a little to one side. Sporoplasm asymmetrically situated, sometimes being entirely confined to the larger valve. Dimensions: breadth 50 to 65 μ , sutural diameter about 8 μ , diameter of polar capsule 4.5 μ , length of polar filament 90 μ .

Remarks: Similar to *C. sphaerulosa* Thél. and occurs with *C. recurvata* Davis in the same organ.

CERATOMYXA SPHAIROPHORA Davis

[Fig. 51]

1917 *Ceratomyxa sphaerophora* Davis 1917 : 224

Habitat: Gall-bladder of *Scoliodon terrae-novae*; Beaufort.

Vegetative form: Pyriform, elongate. Numerous fine filiform pseudopodia at anterior end. Progressive movements rapid. Body colorless and transparent. Ectoplasm clear and homogeneous. Structure of endoplasm highly variable, in majority of trophozoites filled with transparent homogeneous spherules. Small fat globules at the anterior end. In some sporulating individuals, the endoplasm shows vacuolated structure without any spherules, usually, however, sporulating trophozoites exhibit well-defined spherules. The spherules or vacuoles, as the case may be, are separated by a thin layer of distinctly granular endoplasm containing numerous rod-shaped or rounded, colorless bodies, which in their appearance are strikingly like small bacteria tho they are not bacteria, as they fail to take Giemsa stain. Size of sporulating trophozoites 100 to 110 μ by 25 μ . Disporous.

Spore: Shell-valves greatly elongated, tapering gradually toward the ends. Long, attenuated ends of valves hollow and so fragile that it is almost impossible to find an example in which they are not more or less distorted. Sutural plane perpendicular or only slightly oblique to the longest diameter. Polar capsules are spherical and large; slightly convergent, opening some distance apart on the anterior side. Coiled polar filament distinct. Sporoplasm confined to large, central part of spores, but extending farther into one valve than the other. Dimensions: total breadth 115 to 140 μ , sutural diameter about 12 μ , diameter of polar capsules 6 μ , length of polar filament 75 μ .

CERATOMYXA TAENIA Davis

[Figs. 52 and 53]

1917 *Ceratomyxa taenia*

Davis

1917 : 224-225

Habitat: Gall-bladder of *Scoliodon terrae-novae*; Beaufort.

Vegetative form: Similar to those of *C. sphairophora* Davis, and no character has been found by which they may be distinguished. Sporulating trophozoites can be easily distinguished on account of the very different appearance of the spore and their different arrangement within the trophozoites. The spores of this species are situated, as is usually the case in *Ceratomyxa*, with the greater part of the spore parallel to the long axis of the trophozoite, only a part of one valve being bent back along the rest of the spore. Size: sporulating trophozoites length 80 μ , breadth 25 μ . Disporous.

Spore: Valves greatly elongated. Shell very thin, the membrane on opposite sides of each valve being in contact for about two-thirds of its length, forming a thin ribbonlike structure; basal third of each valve only slightly compressed; terminal ribbonlike portion of each valve usually twisted so that plane of ribbon is at right angles to the main part of the spore. Polar capsules small, pyriform to spherical and convergent. Coiled polar filament indistinct. Sutural plane perpendicular to the longest

diameter. Sporoplasm finely granular, filling the basal third of each valve, sometimes extending farther into one valve than the other. Dimensions: breadth 140 to 150 μ , breadth of central portion 45 μ , sutural diameter 6 μ , diameter of the polar capsules 3 μ .

CERATOMYXA ATTENUATA Davis

[Fig. 54]

1917 *Ceratomyxa attenuata* Davis 1917 : 225

Habitat: Gall-bladder of *Scoliodon terrae-novae*; Beaufort (July).

Vegetative form: Elongate, pyriform, with long, tapering posterior process; at anterior end numerous long filiform pseudopodia. Progressive movements rapid. Ectoplasm distinct only at anterior end. Endoplasm filled with small, refractive, yellowish or brownish granules, which are uniformly distributed throughout the trophozoite. Between the brownish granules, the endoplasm is clear and colorless, not granular, except at extreme anterior end where it contains a clump of small fat globules. Size of full-grown trophozoites 100 to 120 by 27 μ . Disporous.

Spore: Valves greatly elongated; a symmetrical, one valve being about 15 μ shorter than the other and ending abruptly; the longer valve tapering gradually to a point. About midway of each valve, is a thin septum; external to the septum the valves are empty. Polar capsules are large, opening on the anterior margin. Coiled polar filaments distinct. Sutural plane oblique to longitudinal axis, usually forming a ridge. Sporoplasm asymmetrically situated in central part of the spore. Dimensions: breadth 115 μ , sutural diameter 9 μ , diameter of polar capsules 4.5 μ , length of polar filament 60 μ .

CERATOMYXA RECURVATA Davis

[Figs. 55 and 56]

1917 *Ceratomyxa recurvata* Davis 1917 : 225-226

Habitat: Gall-bladder of *Cestracion zygaena*; Beaufort (July).

Vegetative form: Pyriform with long, slender posterior process. Body colorless. Actively motile, forming filiform pseudopodia of ectoplasm at anterior end. Endoplasm colorless and granular, filled with large, homogeneous spherules. Full-grown trophozoites 130 to 175 μ , length of the main body about 100 μ . Spores are developed singly from distinct sporoplasts and not necessarily in pairs. Disporous and polysporous (up to 10 spores, 6 and 8 are common numbers).

Spore: Valves greatly curved toward the posterior side, usually symmetrical, but occasionally one may be much more incurved than the other. Valves circular in cross section at the base but toward the ends greatly

flattened. Ends of valves sharply pointed. Polar capsules large, opening at some distance from the anterior margin. Coiled polar filaments distinct. Sporoplasm finely granular usually extending farther into one valve than the other. Dimensions: breadth between points of greatest curvature about 16μ , sutural diameter 8 to 9μ , diameter of polar capsules 4.5μ .

CERATOMYXA LUNATA Davis

[Figs. 57 to 60]

1917 *Ceratomyxa lunata*

Davis

1917 : 226-227

Habitat: Gall-bladder of *Galeocerca tigrinus*; Beaufort (August).

Vegetative form: Pyriform, rounded after being on the slide for some time. Progressive movements slow. Endoplasm filled with large, homogeneous spherules, which are usually colorless, sometimes light yellow. At extreme anterior end, the endoplasm contains numerous small fat globules. Disporous.

Spore: Considerably variable in size and form. The larger and more typical are more or less crescent-shaped; symmetrical; valves curved toward rear, terminating in more or less rounded ends. Polar capsules large and open on opposite sides of spore. Coiled polar filament distinct. Sporoplasm finely granular, symmetrically situated in spore. Smaller spores differ from large ones chiefly in size; valves are much shortened and have a greater curvature, with more distinctly rounded ends. Dimensions: breadth 30μ (longest 38μ), sutural diameter 9μ , diameter of polar capsules 4μ , length of polar filament 37μ . Small forms: breadth 15μ , sutural diameter 7μ , diameter of polar capsules 3μ .

CERATOMYXA ABBREVIATA Davis

[Fig. 61]

1917 *Ceratomyxa abbreviata*

Davis

1917 : 227

Habitat: Gall-bladder of *Scoliodon terrae-novae*; Beaufort (August).

Vegetative form: Elongate, pyriform, with usually a very long, slender posterior process. Body colorless. Progressive movements rapid. Distinct differentiation of protoplasm, posterior process usually composed of ectoplasm (rarely endoplasm may extend into it for a short distance). Pseudopodia, short, tapering or filiform at anterior end. Dimensions: length up to 90μ , breadth 10 to 12μ , diameter of rounded sporulating trophozoites about 27μ . Disporous.

Spore: Roughly crescent-shaped; sutural diameter exceptionally great in comparison with the breadth. Ends of valves rounded, slightly asymmetrical. Shell exceptionally tough and resistant to reagents. Polar capsules large, prominent and open on opposite sides of spore. Sporoplasm finely granular, confined entirely to one valve. Dimensions: breadth 17μ , sutural diameter 14μ , diameter of polar capsules 4.5μ .

CERATOMYXA FLAGELLIFERA Davis

[Fig. 62]

1917 *Ceratomyxa flagellifera* Davis 1917 : 227

Habitat: Gall-bladder of *Carcharhinus* sp?; Beaufort (July).

Vegetative form: Pyriform, short, tapering toward the posterior end, sometimes dividing into a number of long, slender, transparent processes. Extremely long filiform pseudopodia, developed at anterior end, can be seen to sweep slowly back like a whiplash until they come to lie by the side of the body. Progressive movements slow. Ectoplasm clear, transparent, forming a distinct layer at anterior end. Endoplasm in large trophozoites filled with large numbers of rod-shaped, bacteria-like bodies, which are more abundant in the anterior half than in the posterior. Endoplasm in younger trophozoites, with much less or without any bacteria-like bodies, shows a vacuolated structure. Size up to 115 to 120 μ in length and 40 to 45 μ in breadth. Disporous.

Spore: Valves greatly elongated, conical, with rounded ends. Sutural ridge well marked. Polar capsules large, opening on opposite sides of spore. Coiled polar filament very distinct. Sporoplasm granular, symmetrically situated, but extending only a short distance into each valve. Dimensions: breadth 118 μ , sutural diameter 12 μ , diameter of polar capsules 6 μ .

CERATOMYXA AGGLOMERATA Davis

[Fig. 63]

1917 *Ceratomyxa agglomerata* Davis 1917 : 228

Habitat: Gall-bladder of *Synodus foetans*; Beaufort.

Vegetative form: Pyriform, usually with long, slender, posterior process. Body colorless and transparent. Actively motile, moving by means of characteristic wavelike movements of the ectoplasm, from which are projected numerous short, conical to filiform pseudopodia. Pseudopodia travel back along sides of body for about one-third its length and then disappear, new ones being continually formed at the anterior end. Ectoplasm distinguishable at anterior end. Endoplasm clear, very transparent, usually homogeneous, sometimes finely granular. Large numbers of fat globules usually present. Size of sporulating trophozoites 38 μ by 12 μ . Disporous

Spore: Asymmetrical, one valve being smaller and more attenuated than the other; larger valve compressed. Polar capsules spherical. Coiled polar filaments indistinct. Sporoplasm filling nearly entire smaller valve, but only extending a short distance into the larger one. Dimensions: breadth 24 to 28 μ , sutural diameter 5 μ , diameter of polar capsules 3 μ .

CERATOMYXA AMORPHA Davis

[Fig. 64]

1917 *Ceratomyxa amorpha* Davis 1917 : 228

Habitat: Gall-bladder of *Synodus foetans*; Beaufort.

Vegetative form: Rounded or irregular in shape, with short lobose pseudopodia; not pyriform; slowly ameboid. Body colorless. Ectoplasm well developed, forming a distinct layer; transparent, finely granular. Endoplasm granular, with large numbers of small fat globules scattered through it or aggregated into one or two large clumps (difference between the present form and *C. agglomerata*). Disporous.

Spore: Asymmetrical; crescent-shaped; valves short, conical, somewhat compressed. One valve distinctly smaller and more conical than the other. Sutural ridge perpendicular to longitudinal axis. Polar capsules large, opening at some distance from the anterior side. Coiled polar filaments distinct. Sporoplasm granular, asymmetrically situated, being chiefly confined to smaller valve. Dimensions: breadth 27μ , sutural diameter 11μ , diameter of polar capsules 4μ .

CERATOMYXA MONOSPORA Davis

[Figs. 65 to 67]

1917 *Ceratomyxa monospora* Davis 1917 : 228-229

Habitat: Gall-bladder of *Peprilus alepidotus*; Beaufort. Abundantly present in June, much less in July, being entirely absent in the bladder at the end of the month.

Vegetative form: Pyriform, with a slender posterior process and one to several filiform pseudopodia at anterior end. Body colorless and transparent. Movements very slow. No clear differentiation between ectoplasm and endoplasm, the entire body being composed of a clear, finely granular protoplasm. Fat globules more abundant in larger individual, which are aggregated into small clumps. Size of vegetative trophozoites up to 24μ in length and 15μ in width. Monosporous form much smaller than disporous. Monosporous and disporous. Nearly entire substance of trophozoite is used up in spore formation.

Spore: Crescent-shaped. Valves cylindrical, tapering toward the end, which is rounded and compressed. Curvature of valves varies. One valve is more attenuated than the other. Sutural ridge perpendicular to the longest diameter. Polar capsules large. Sporoplasm usually asymmetrically situated. Dimensions: breadth 18 to 25μ , sutural diameter 5 to 6μ , diameter of polar capsules 3μ .

Remarks: This species is evidently very close to *C. pallida* Thél. Similar form was found in *Prionotus evolans* (gall-bladder), which showed somewhat larger trophozoites and spores than *C. monospora*.

CERATOMYXA STREPTOSPORA Davis

[Figs. 68 and 69]

1917 *Ceratomyxa streptospora* Davis 1917 : 229

Habitat: Gall-bladder of *Chaetodipterus faber*; Beaufort (June, but not in July).

Vegetative form: Pyriform, colorless and transparent. A few conical, filiform, wavelike pseudopodia at anterior end. Ectoplasm recognizable at anterior end. Endoplasm finely granular, with a few, small, fat globules, filled with transparent, homogeneous spherules. Size: 48 by 12 μ to 60 by 9 μ . Disporous.

Spore: Compressed valves greatly elongated, with rounded ends. Sutural ridge. Polar capsules spherical. Coiled polar filament indistinct. Sporoplasm finely granular, entirely filling both valves. Dimensions: breadth 34 to 39 μ , sutural diameter 4 μ , diameter of polar capsules 3 μ .

CERATOMYXA AGGREGATA Davis

[Fig. 70]

1917 *Ceratomyxa aggregata* Davis 1917 : 229

Habitat: Gall-bladder of *Leostomus xanthurus*, *Micropogon undulatus*; Beaufort (July).

Vegetative form: Form rounded to somewhat irregular in shape, rarely pyriform; slowly ameboid. Body colorless and transparent. No clear differentiation of protoplasm. Endoplasm finely granular, containing numbers of small fat globules. Sporulating trophozoites show a tendency to collect in groups composed of a large number of individuals so closely associated that it is often impossible to make out the individual outlines. Size of full-grown form 18 μ by 14 μ . Disporous.

Spore: Crescent-shaped; valves much elongated, tapering toward the ends, which are compressed. Polar capsules spherical and opaque. Sporoplasm granular, situated symmetrically in the spore cavity. Dimensions: breadth about 50 μ , sutural diameter 6 to 7 μ , diameter of polar capsule 3.5 μ .

CERATOMYXA UNDULATA Davis

[Fig. 71]

1917 *Ceratomyxa undulata* Davis 1917 : 230

Habitat: Gall-bladder of *Ancylopesetta quadrocellata* Gill.; Beaufort (June to August).

Vegetative form: Pyriform, sometimes fusiform, tapering toward posterior end. Movements rapid. Body colorless. Ectoplasm observable at anterior part, constantly undergoes rapid, wavelike undulating movements and extrudes fine conical or filiform pseudopodia. Pseudopodia

are formed very rapidly and vary in length. After reaching a considerable length the pseudopodia usually travel posteriorly along sides of body and then disappear. Endoplasm very transparent, often vacuolated, containing numerous small fat globules. Size of full-grown trophozoite: 25μ by 10 to 12μ in average. Disporous.

Spore: Crescent-shaped. Valves cylindrical, not compressed, ends rounded, one valve being somewhat longer and more conical than the other. Polar capsules convergent. Coiled polar filaments distinct. Sporoplasm granular, asymmetrically situated, sometimes being almost confined to more conical valve. Dimensions: breadth 22 to 24μ , sutural diameter 6μ , diameter of polar capsules 3μ .

CERATOMYXA NAVICULARIA Davis

[Figs. 72 and 73]

1917 *Ceratomyxa navicularia* Davis 1917 : 230

Habitat: Urinary bladder of *Paralichthys dentatus*, *P. albiguttus*, *Sphaeroides maculatus*; Beaufort (June to August).

Vegetative form: Rounded or slightly irregular in shape, never pyriform. Body colorless. Very slow ameboid. No distinct ectoplasm. Entire trophozoite finely granular, containing a few small fat globules. Nearly entire body is used up in the formation of spores. Diameter about 17μ . Disporous.

Spore: Variable in shape and size. Symmetrical or asymmetrical, often boat-shaped, slightly compressed dorso-ventrally, with rounded ends. Polar capsules convergent, shows polar filament indistinctly. Sporoplasm finely granular, extending into both valves, but usually somewhat farther into one than the other. Dimensions: breadth 14 to 22μ (average 16μ), sutural diameter 5 to 7.5μ (average 6μ), diameter of polar capsules 2μ .

CERATOMYXA SPINOSA Davis

[Fig. 74]

1917 *Ceratomyxa spinosa* Davis 1917 : 230

Habitat: Urinary bladder of *Paralichthys albiguttus*; Beaufort.

Vegetative form: Rounded or slightly irregular in shape, with short, lobose pseudopodia; slowly ameboid. Body colorless and transparent. Distinct differentiation of protoplasm along the entire surface, ectoplasm forming outer layer. Endoplasm faintly granular, with numerous small fat globules. Monosporous and disporous.

Spore: Central portion greatly enlarged; ovoid, with very long tapering process extending out from each end. Sutural line perpendicular to the longest diameter. Polar capsules large and spherical. Sporoplasm finely

granular, chiefly located in one valve, extending into the other only a short distance beyond the capsule. Dimensions: breadth 80μ , breadth of enlarged central portion 13μ , sutural diameter 7μ , diameter of polar capsules 4μ .

Genus MYXOPROTEUS Doflein emend. Davis

1898	<i>Myxoproteus</i>	Doflein	1898 : 287
1917	<i>Myxoproteus</i>	Davis	1917 : 219

The characters of the genus are described on page 56.

Type species: *Myxoproteus ambiguus* (Thélohan) Doflein.

MYXOPROTEUS AMBIGUUS (Thélohan) Doflein

[Figs. 75 to 80]

1895	<i>Myxosoma ambiguus</i>	Thélohan	1895 : 344
1898	<i>Myxoproteus ambiguus</i>	Doflein	1898 : 287-288

Habitat: Urinary bladder of *Lophius piscatorius*; Le Croisic, Rovigno, Napoli.

Vegetative form: Polymorphous. Color milky white. Protoplasm is filled with numerous granules and fat globules. Pseudopodia, short, pointed lobose. Plasmogamy and plasmotomy take place. Many small individuals formed apparently by plasmotomy, often, make up groups. Disporous, polysporous?

Spore: Almost pyramidal, with anterior processes. Two very large polar capsules at the anterior end. The distance between these capsules is equal to or greater than the diameter of the capsules. Sporoplasm with two nuclei. Dimensions: length 25μ , breadth 18 to 20μ , diameter of polar capsules 7μ .

MYXOPROTEUS CORDIFORMIS Davis

[Figs. 81 to 83]

1917	<i>Myxoproteus cordiformis</i>	Davis	1917 : 231
------	--------------------------------	-------	------------

Habitat: Urinary bladder of *Chaetodipterus faber*; Beaufort (June, July).

Vegetative form: Rounded; very slowly ameboid, usually forming a single, short, lobose pseudopodium. Body colorless and transparent. Ectoplasm not distinct. Entire trophozoite finely granular, with a few fat globules. Rarely vacuolar. Small trophozoites often show a single large, central vacuole. Rounded sporulating trophozoites 18μ in diameter. Disporous.

Spore: Heart-shaped in front view, with peculiar wing-like expansions on each side which contain remains of parietal cells. Sutural plane oblique in position. Capsulogenous cells distinct. Sporoplasm finely granular, fills the extracapsular cavity of the spore. Dimensions: length 12μ , breadth 10 to 11μ , thickness 6μ , polar capsules 3 to 4μ in diameter.

MYXOPROTEUS CORNUTUS Davis

[Fig. 84]

1917 *Myxoproteus cornutus*

Davis

1917 : 231

Habitat: Urinary bladder of *Bairdiella chrysura*; Beaufort.

Vegetative form: Somewhat elongated or irregular in shape, with short lobose pseudopodia; slowly ameboid. Differentiation of protoplasm clear. Ectoplasm well developed, hyaline; in rounded individuals forming a distinct layer around the body. Endoplasm opaque, contains coarse refringent granules varying in shape, and a few fat globules. In contracted rounded resting condition, endoplasm becomes condensed, while ectoplasm appears more abundant. Rounded trophozoites up to 27μ in diameter. Disporous.

Spore: Heart-shaped, with two anterior processes. Shell very thick. Polar capsules large, opening some distance apart. Coiled polar filament distinct. Sporoplasm finely granular, with a few small fat globules, fills the extracapsular cavity of the spore. Dimensions: sutural diameter exclusive of the processes 9μ , breadth 12μ , length of processes 5μ , diameter of polar capsules 3μ .

Genus WARDIA nov. gen.

The characters of the genus are described on page 56.

Type species: *Wardia ovinocua* nov. spec.

WARDIA OVINOCUA nov. spec.

[Figs. 85 to 95]

Habitat: Ovum and connective tissue of ovary of *Lepomis humilis* Girard;* Salt Fork, Ill. (November). Only one fish, 6.5 cm. long with normal appearance, was found to be infected.

Vegetative form: Trophozoites form cysts visible to the naked eyes as white spherical spots in the pink-colored ovary. Four cysts present. The cyst (Figs. 85 and 86), in section, shows a circular form surrounded by several layers of hypertrophied nurse cells and connective tissue, in which many large blood vessels are present. Protoplasm is not clearly differ-

*Professor F. Smith of the Department kindly identified all the fish that were collected in the vicinity of Urbana and mentioned in this paper as hosts, for which the writer wishes to express his appreciation.

entiated into ectoplasm and endoplasm, the whole protoplasm showing granulated reticular structure. Cysts contained numerous fully developed spores and a small number of spores in developmental stages, which suggested the fact that two spores rise from each pansporoblast. The parasite is also found in the state of diffuse infiltration in the connective tissue around the cyst. Diameter of cysts 316 to 445 μ in sections. Polysporous.

Spore: In front view, isosceles triangular form, two sides of which usually convex, with more or less attenuated anterior end (Figs. 87, 90, 92); in profile, ellipsoidal (Fig. 88); and oval viewed from the anterior end (Fig. 89). Sutural plane at right angles to the longest diameter (Figs. 87 and 89). Shell comparatively thin except at the anterior end and has many fine network-like ridges on the surface. These ridges are hardly observable on fresh spores on account of their fine form and the conspicuously large polar capsules lying in the spore. When stained, however, they are not only made distinctly visible, but the prolongation of each ridge from the posterior edge which forms about 1 μ long fringe-like structure is also more clearly recognized (Figs. 90–95). Two large and spherical polar capsules located in the central portion of the spore. Coiled (5 to 6 times) polar filament extremely distinct. The openings of polar capsules at the anterior end. Sporoplasm comparatively small, finely granular, without any vacuole, contains two small nuclei, when stained. Dimensions in vivo: sutural diameter 9 to 10 μ , breadth 10 to 12 μ , thickness 6 μ , diameter of the polar capsule 4 μ , length of polar filament 35 to 45 μ .

WARDIA OHLMACHERI (Gurley) Kudo

[Figs. 96 and 97]

1893	<i>Myxosporidian</i>	Ohlmacher	1893 : 561–567
1893	<i>Chloromyxum ohlmacheri</i>	Whinery	1893 : 660–662
1894	<i>Chloromyxum (Sphaerospora)</i> <i>ohlmacheri</i>	Gurley	1894 : 267–272
1895	? <i>Leptotheca ranae</i>	Thélohan	1895 : 383
1899	<i>Leptotheca ohlmacheri</i>	Labbé	1899 : 87

Habitat: Urinary tubules of kidney of *Bufo lentiginosus* Shaw and kidney of *Rana esculenta* and *R. temporaria (R. fusca)*; Sycamore, De Kalb county, Ill.

Vegetative form: Not found.

Spore: Transversely elliptic. Sutural plane perpendicular to the longer axis of the spore. A well defined undulate-parallel longitudinal striation on the shell. Sutural ridge comparatively well marked. Two polar capsules lying side by side, occasionally only one. Dimensions: sutural diameter 6 μ , breadth 8 μ , diameter of polar capsule 3 to 3.5 μ , length of polar filament 6 to 8 times the breadth of spore (48 to 64 μ).

Remarks: This form is apparently very much different from any species of genus *Leptotheca*, in the general form, form of polar capsules,

striations on the shell and the habitat. Tho the form of the spore is different from the type species of the genus *Wardia* and nothing is known about the vegetative form, the presence of large spherical polar capsules in the central portion of the spore, the striations on the shell and the occurrence of the same nature, i.e., from fresh waters in the close-by localities, show its nearer relationship to the genus *Wardia* than to the genus *Leptotheca*. Hence it is placed here provisionally.

Genus MITRASPORA Fujita emend. Kudo

1912 *Mitraspora* Fujita 1912 : 259-260

The characters of the genus are described on page 56.

Type species: *Mitraspora cyprini* Fujita.

MITRASPORA CYPRINI Fujita

[Figs. 98 to 104]

1912 *Mitraspora cyprini* Fujita 1912 : 259-260

Habitat: Renal tubules of kidney and ureters of *Cyprinus carpio* L. and *Carassius auratus* L.; Sapporo (winter), Tokio (March).

Vegetative form: Fujita's only description is as follows: "The sporoblast contains generally three or four spores." The present writer observed a similar form in the ureter and the renal tubule of the kidney of *Cyprinus carpio* L., in Tokio. The observations are as follows: Trophozoites small ameboid (Fig. 98). Body colorless. Movements tardy. Differentiation of protoplasm imperfect. The hyaline ectoplasm recognizable at one side of the body, where lobose pseudopodia are formed (Figs. 98-99). Endoplasm granular with vacuoles and brownish granules, which become larger as the body grows. Size 10 to 40 μ . Disporous (Kudo) and polysporous (? , Fujita).

Spore: Fujita's descriptions are as follows: Form resembles monk's hood, slightly attenuated at its anterior end. Shell uniformly thin, except at two points of the truncated posterior end. Each shell valve has eight distinct striations which run longitudinally and turn into long cilia up to 5.8 μ long, planted in a single row at the posterior end of the spore. Two polar capsules at the anterior end. The nucleus is obscure and no vacuole is present. Dimensions: length 10 to 13 μ , breadth 5 μ , polar capsules 3.8 μ by 2 μ , length of polar filament 15 μ (weak glycerine). The writer observed the following facts: More rounded with rounded anterior end in front and side views. Shell more or less thick along the entire posterior margin. Striations on shell, variable in number. Sporoplasm granular, without any vacuole, exhibits two nuclei when stained. Posterior filaments 5 to 6 in number and 5 to 6 μ long, being absent in some spores. Dimensions in vivo: length 10 μ , breadth 8 to 9 μ , thickness 6 to 8 μ , polar capsule 4 μ by 1.5 to 2 μ , length of polar filament 35 to 40 μ .

Remarks: Tho Fujita does not describe the vegetative form and there are some differences in the form and size of the spore between the forms, the writer does not find out any objection against the union of the above mentioned two forms.

MITRASPORA CAUDATA (Parisi) Kudo

[Figs. 105 to 107]

1910	<i>Sphaerospora caudata</i>	Parisi	1910 : 253-254
1912	<i>Sphaerospora caudata</i>	Parisi	1912 : 289
1913	<i>Sphaerospora caudata</i>	Parisi	1913 : 396-402

Habitat: Renal tubules of kidney of *Alosa finta* Cuv. var. *lacustris* Fatio; Lake Como.

Vegetative form: Rounded or variously elongated owing to the movements. Protoplasm is distinctly differentiated into ectoplasm and endoplasm. Ectoplasm, hyaline and homogeneous, forms slowly moving lobose pseudopodia. Endoplasm granular, contains yellow globules and fat granules. Disporous and polysporous.

Spore: Subspherical in front view; oval in profile; anterior end being more rounded than the posterior end. Shell rather thick, longitudinally striated. In front view, the posterior end enlarged into a quadrangular form, which appears as a small spine in side-view and which projects backward long and fine filaments, usually six in number. Two well developed polar capsules open on each side of the sutural plane. Polar filament coiled 5 to 6 times. Sporoplasm without any iodophilous vacuole. Dimensions: external length 10 to 11 μ , internal length 7 to 9 μ , length of polar capsules 4 to 4.5 μ , length of polar filament up to 48 μ , length of posterior filaments up to 28 μ .

MITRASPORA ELONGATA nov. spec.

[Figs. 602 to 621]

Habitat: In the urinary tubules and tissue of kidney of *Lepomis cyanellus*; Crystal Lake, Urbana, Ill. From June to July, all the fish examined, 36 in number and 10 cm. in average length, were found to be infected. Other fish such as *Lepomis pallidus* and *Lepomis humilis*, caught at the same time, were free from the infection. Early in June, the number and size of the parasites in a host body were rather small and only a small number of spores could be recognized in the fresh state with the addition of potassium hydrate solution. The growth of the parasite was rather remarkable during the hot weeks in the latter part of June and July so that every fish caught on July 17th showed a heavy infection, exhibiting small whitish pustules over the surface of the organ. During June, vegetative forms and spores were found in the lumen of the urinary tubules,

altho some contained the parasitic masses in the tissue. About the middle of July, the parasite forms conspicuous cysts in the tissue thruout the organ. The cyst may or may not be covered by a thick layer of connective tissue from the host. Aside from this hypertrophy, the host did not show any pathological change which could be recognized.

Vegetative form: Youngest trophozoite found in the urinary tubule is multinucleate, rounded, and of from 20 to 50 μ in diameter. The protoplasm is not differentiated, the entire body is finely granular or coarsely reticular in structure. In the protoplasm are to be seen nuclei and sporoblasts at different stages of development. The union of two propagative cells similar to that of *Myxobolus toyamai* produces a small body which develops into a single sporoblast and ultimately into a single spore (Figs. 605-613). In later stages, the trophozoite reaches a size of 200 μ in diameter showing many stages of spore formation and mature spores, surrounded by thick layers of connective tissue from the host. Polysporous.

Spore: Elongated oblong with pointed anterior and truncated posterior extremities. The width is often greatest at the middle of the polar capsules, the posterior portion is much narrower than the anterior. Nearly circular in the cross-section thru the polar capsules. The shell is thin, the sutural line being faintly marked in fresh state. It generally is obliquely located in relation to the capsules. The shell also shows fine longitudinal striations, 14 to 16 in number, on each valve. The sutural line as well as the striations are best seen in spores stained with Heidenhain's iron hematoxylin. Two polar capsules elongated pyriform, mostly equal in size, occupy the anterior half of the spore. Abnormal situations of the polar capsules are sometimes observed (Fig. 619). The coiled polar filament is faintly visible in fresh spores. It is spirally coiled along the wall of the polar capsule without any central axis. This fact was clearly observed in stained section as is shown in Figs. 620 and 621. The filament has seven or eight windings, thus agreeing with the actual length of the extruded polar filament. The polar filament was extruded under the action of potassium hydrate solution. The extrusion takes place even in some spores which were treated with Schaudinn's fixative and kept in 95 per cent alcohol for three months (see the similar observations on *Myxobolus discrepans* on page 157). The sporoplasm is finely granular and transparent. When stained, it shows two nuclei in the center or near the posterior part of the body. Dimensions of preserved spores: length 15 to 17 μ , breadth 5 to 6 μ , thickness 4.5 to 5.5 μ , polar capsule 7.5 μ by 2 μ , length of polar filament 40 to 50 μ .

Suborder SPHAEROSPOREA nom. nov.

The definition of the suborder is recorded on page 57.

Family CHLOROMYXIDAE Thélohan

1892	<i>Chloromyxidées</i>	Thélohan	1892 : 173
1895	<i>Chloromyxidées</i>	Thélohan	1895 : 344

The characters of the family are described on page 57.

Genus CHLOROMYXUM Mingazzini

1890	<i>Chloromyxum</i>	Mingazzini	1890 : 160
1892	<i>Chloromyxum</i>	Thélohan	1892 : 173-176
1895	<i>Chloromyxum</i>	Thélohan	1895 : 344

The characters of the genus are described on page 57.

Type species: *Chloromyxum leydigi* Mingazzini.

CHLOROMYXUM LEYDIGI Mingazzini

[Figs. 108 to 113]

1851		Leydig	1851 : 225-234
1852		Leuckart	1852 : 435
1854		Lieberkühn	1854 : 352
1890	<i>Chloromyxum leydigi</i>	Mingazzini	1890 : 160-164
1892	<i>Chloromyxum leydigi</i>	Thélohan	1892 : 166, 169-170
1894	<i>Chloromyxum leydigi</i>		
	<i>Chloromyxum incisum</i>	Gurley	1894 : 259-260
1895	<i>Chloromyxum leydigi</i>		
	<i>Chloromyxum incisum</i>	Thélohan	1895 : 345-346
1898	<i>Chloromyxum leydigi</i>	Doflein	1898 : 292, 310, etc.
1912	<i>Chloromyxum leydigi</i>	Erdmann	1912 : 149-162
1916	<i>Chloromyxum leydigi</i>	Georgévitch	1916a : 3
1917	<i>Chloromyxum leydigi</i>	Davis	1917 : 236-237
1917	<i>Chloromyxum leydigi</i>	Erdmann	1917 : 276-321
1918	<i>Chloromyxum leydigi</i>	Georgévitch	1918 : 182-189

Habitat: Gall-bladder of *Rhina squatina* L., *Spinax spinax* L., *Scyllium canicula*, *S. asterias*, *Raja batis* L., *R. clavata* L., *R. undulata* Lac., *Torpedo narce* Ris., *T. marmorata*, *T. ocellata*, *T. torpedo* L., *Acanthias acanthias* L., *Trygon pastinaca* L., *Dasybatis hastatus*, *D. sabina*, *Pteroplatea maclura* Le Sueur, *Scoliodon terrae-novae*, *Cestracion zygaena*, *C. tiburo*, *Carcharhinus limbatus*; Roscoff, Concarneau, Marseille, Banyuls, Rovigno, Heligoland, Beaufort, Monaco (May), Napoli, Genova. Erdmann observed the species at Naples from March to August. She noticed mixed infection with *Ceratomyxa reticulata* and especially with *Leptotheca parva*. Georgévitch studied the parasite at Monaco from February to April.

Vegetative form: Polymorphous, being spherical, oval or irregular. The change of the form rather rapid under favourable conditions. Differentiation of protoplasm distinct. Ectoplasm with pseudopodia of various form, i.e., lobose, filiform or intermediary; short, pointed or branched. Endoplasm alveolar, filled with yellowish granules. Doflein observed the plasmotomic multiplication of young trophozoites. Polysporous. Erdmann's observations (1917) may be summarized as follows: Ameboid.

Color of the body greenish to dark green. The protoplasm is clearly differentiated into ectoplasm and endoplasm. The ectoplasm is hyaline and covers the entire surface of the body. It appears as a fine fibrous structure when fixed with Bouin's solution. The endoplasm contains besides nuclei, two kinds of spherules; one smaller and yellowish "color-carriers" and the other larger and light to dark greenish reserve bodies. The color-carrier is in part composed of myelin, while the reserve body is of glycogenous nature. The infection was studied experimentally *per os*: young trophozoites appeared in 3 to 5 days which continued to 10th day, various trophozoites were seen in 13 to 19 days, and sporulating individuals were first recognized in 39 days after the infection. The trophozoite multiplies in number either by fission or by budding. It usually contains enclosures which seem to be degenerating erythrocytes. Mictosporous.

Spore: Ovoidal. Shell-valves show wide edge at sutural plane, which is attenuated at the anterior end and forms a quadrilateral process at the posterior end, from which a row of cilia grows. Shell-valves have ridges (6 to 7), which run parallel to the posterior margin. The striations may vary considerably. Four polar capsules at the anterior end. Dimensions: length 8μ . Erdmann gave the following dimensions: Spores from *Torpedo marmorata* and *T. ocellata*: length 6 to 9μ , breadth 5μ , polar capsule 3μ by 2μ . Those from *Scyllium asterias*: length 8 to 9μ , breadth 6μ , polar capsule 2μ by 1μ . Those from *Raja batis*: length 7 to 8μ , breadth 5μ , polar capsule 2μ by 1μ . Length of polar filament 20 to 30μ (absolute alcohol warmed up to 40°C .).

CHLOROMYXUM CAUDATUM Thélohan

[Fig. 114]

1895 *Chloromyxum caudatum* Thélohan 1895 : 346

Habitat: Gall-bladder of *Molge cristata* Laur.; Vicinity of Rennes.

Vegetative form: Body yellowish with lobose pseudopodia. Protoplasm finely granular.

Spore: Oval or spheroidal. Shell enlarged at the anterior part, having a simple or bifurcated tail-like process, as in Henneguya, at the posterior end. Dimensions: total length 18μ , length 8μ , breadth 6 to 7μ , length of tail 10μ .

CHLOROMYXUM QUADRATUM Thélohan

[Figs. 115 to 117]

1891		Pfeiffer	1891 : 111
1893		Pfeiffer	1893 : 81
1895	<i>Chloromyxum quadratum</i>	Thélohan	1895 : 347
1912	<i>Chloromyxum quadratum</i>	Parisi	1912 : 289
1913	<i>Chloromyxum quadratum</i>	Awerinzew	1913a : 155
1913	<i>Chloromyxum quadratum</i>	Fermor	1913 : 199

Habitat: Muscle of *Syngnathus acus* L., *Trachurus trachurus* L., *Nerophis aequorens* L., *Callionymus lyra* L., *Coris julis* L., *Ariodes polystaphylodon*, kidney of *Blennius gattorugine* Brunn; Helder, Roscoff, Concarneau, Marseille, Beira (Africa), Napoli (summer).

Vegetative form: Not described by any of these authors.

Spore: Quadrangular pyramid with curved edges and roundish angles. Four polar capsules at the anterior end. Dimensions: length 6μ , breadth 5μ , length of polar filament 8 to 10μ .

CHLOROMYXUM FLUVIATILE Thélohan

[Fig. 118]

1892	<i>Chloromyxum fluviatile</i>	Thélohan	1892 : 173-176
1895	<i>Chloromyxum fluviatile</i>	Thélohan	1895 : 346

Habitat: Gall-bladder of *Leuciscus cephalus* L.; Paris.

Vegetative form: Young trophozoites colorless; adults yellowish. Form highly variable. Active change of the form of body. Clear differentiation between ectoplasm and endoplasm. Ectoplasm usually recognizable at one end of the body where lobose pseudopodia are formed. Size reaches 25 to 30μ . Polysporous.

Spore: Spherical, generally small. Sutural ridge fairly well marked. Dimensions: 7 to 8μ in diameter.

CHLOROMYXUM MUCRONATUM Gurley

[Figs. 119 to 122]

1854		Lieberkühn	1854 : 352-353
1879		Leuckart	1879 : 248
1882		Bütschli	1882 : Pl. 38 : 17
1883		Balbani	1883 : 201, 203
1893	<i>Chloromyxum mucronatum</i>	Gurley	1893 : 419
1894	<i>Chloromyxum mucronatum</i>	Gurley	1894 : 264, 265
1908	<i>Chloromyxum mucronatum</i>	Auerbach	1908 : 456
1909	<i>Chloromyxum mucronatum</i>	Auerbach	1909a : 71

Habitat: Urinary-bladder and kidney of *Lota lota* L.; Bodensee, other localities not mentioned.

Vegetative form: Spherical, elliptical or irregular. Size up to 75μ in diameter. Protoplasm containing irregularly scattered fat-like globules.

Spore: Sharp-contoured; subglobular, mucronate anteriorly. Dimensions: length 8μ .

CHLOROMYXUM DIPLOXYS (Gurley) Thélohan

[Figs. 123 to 125]

1866		Balbiani	1866 : 600-602
1867		Balbiani	1867 : 275, 276, 335
1882		Bütschli	1882 : 590
1890		Pfeiffer	1890 : 559
1892		Henneguy et Thélohan	1892 : 587
1893	<i>Cystodiscus ? diploxys</i>	Gurley	1893 : 411
1894	<i>Cystodiscus ? diploxys</i>	Gurley	1894 : 281
1895	<i>Chloromyxum diploxys</i>	Thélohan	1895 : 347

Habitat: Abdominal cavity of *Tortrix viridana* L. (imago);

Vegetative form: Trophozoites form spherical cysts, 230 to 400 μ in diameter. Cyst membrane rather thick. Protoplasm containing brownish granules, and fat-like globules (red with iodine).

Spore: Elliptic or slightly flattened. Sutural line straight, forming a ridge. Two polar capsules at each end.

CHLOROMYXUM PROTEI Joseph

1905	<i>Chloromyxum protei</i>	Joseph	1905 : 450-451
1907	<i>Chloromyxum protei</i>	Joseph	1907 : 398-412

Habitat: Renal tubules of kidney of *Proteus anguineus* L.; Vienna.

Vegetative form: Generally rounded or sausage form. No clear differentiation between ectoplasm and endoplasm. Movements slow. Probable occurrence of plasmotomy by budding and division. Size: 40 to 45 μ by 28 to 40 μ .

Spore: Spherical. Shell finely striated parallel to the sutural line. Four polar capsules each with an independent opening. Dimensions: 10 to 13 μ in diameter, polar capsules 4 to 6 μ long. The polar filament appears to be rather short.

CHLOROMYXUM TRUTTAE Léger

[Fig. 126]

1906	<i>Chloromyxum truttæ</i>	Léger	1906 : 267-270
------	---------------------------	-------	----------------

Habitat: Gall-bladder and gall-duct of *Trutta fario* L.; Dauphiné.

Vegetative form: Ameboid form. Elongated. Form resembles an *Amoeba limax* of about 40 μ in length. Roundish or irregularly contoured, with small pseudopodia. Ovoidal or spherical, 25 to 40 μ in diameter without any visible pseudopodia (resting state). Body colorless, clear and hyaline. Very active movements which last for several hours after the death of the host. Broad and obtuse pseudopodia well developed at the anterior end of the body. Endoplasm alveolar, contains variable numbers of nuclei, which are seen *in vivo*, refractive bodies and chromatic granules. Monosporous(?) and polysporous.

Spore: Spherical. Four polar capsules of different size. Shell-valves marked with parallel ridges. Dimensions: 8 to 9 μ in diameter.

CHLOROMYXUM CRISTATUM Léger

[Figs. 127 and 128]

1906	<i>Chloromyxum cristatum</i>	Léger	1906 : 270-272
------	------------------------------	-------	----------------

Habitat: Gall-bladder of *Tinca vulgaris* Cuv.; Grenoble.

Vegetative form: Ordinarily massive, with oval or round contours, without noticeable pseudopodium. Ectoplasm hyaline. Endoplasm granular and colorless. Average diameter of the adults about 20 μ . Monosporous, rarely disporous.

Spore: Spherical or subspherical. Ten marked ridges run antero-posteriorly on each shell-valve, so that it presents a cog-wheel form in cross section. Four polar capsules at the anterior end, one pair being smaller than the other. Sporoplasm with two nuclei. Dimension: 10 to 11 μ .

CHLOROMYXUM DUBIUM Auerbach

[Figs. 129 to 133]

1908	<i>Chloromyxum dubium</i>	Auerbach	1908 : 456-459
1910	<i>Chloromyxum dubium</i>	Auerbach	1910c : 177

Habitat: Gall-bladder of *Lota vulgaris* Cuv.; Bodensee (April to September).

Vegetative form: Spherical or rounded. Rarely irregular forms. Protoplasm is differentiated distinctly into ectoplasm and endoplasm. Ectoplasm very thin, forms pseudopodia which move slowly. Endoplasm granular, contains fat globules. Majority of the trophozoites appear to live floating in the bile, while some are attached to the epithelium of the bladder. Disporous and polysporous.

Spore: Spherical, with four polar capsules. Each shell valve has longitudinal ridges, variable in number (6 ridges are found on the drawing), which run parallel to the sutural line. Four polar capsules of nearly same size and convergent. Sporoplasm finely granular with two nuclei. Dimensions: diameter 10.8 μ , length of polar capsule 3.6 μ .

CHLOROMYXUM sp. Awerinzew

[Fig. 134]

1908	<i>Chloromyxum</i> sp.	Awerinzew	1908 : 43, 47, 48
------	------------------------	-----------	-------------------

Habitat: Gall-bladder of *Raja radiata*; Murman coast?.

Vegetative form: Form rounded. The protoplasm is distinctly differentiated into ectoplasm and endoplasm. Ectoplasm hyaline and comparatively abundant in quantity compared with endoplasm, forms lobose pseudopodia of active movements. Endoplasm vacuolated, contains enclosures. Between the two layers, a thin layer of protoplasm, reticular in structure and stained deeply with hematoxylin, is present.

Spore: No figure.

CHLOROMYXUM THYMALLI Lebzelter

1912 *Chloromyxum thymalli* Lebzelter 1912 : 295-296Habitat: Gall-bladder of *Thymallus thymallus* L.; Vienna?

Vegetative form: Irregular form, 33 to 35 μ long in average. Endoplasm contains fat globules which stain brown with carmine. Trophozoites attached to the epithelium. In average, 6 spores formed in each individual. Intracellular stage in the epithelial cell is supposed. Polysporous.

Spore: Spherical. Shell structure similar to *C. protei*, but ridges are more developed and exhibit somewhat wavy courses. Polar capsules of equal size. Dimensions: 9 to 9.5 μ in diameter, polar capsules 3 μ .

CHLOROMYXUM KOI Fujita

[Fig. 135]

1913 *Chloromyxum koi* Fujita 1913 : 257-259Habitat: Gall-bladder of *Cyprinus carpio* L.; Sapporo (Nippon).

Vegetative form: Spherical, with greatest diameter up to 50 μ , containing 1 to 3 spores. Each spore is situated in a clear space surrounded by a membranous envelope (sporoblast?), around which there is some finely granular matter (endoplasm?).

Spore: Spherical, exhibiting a somewhat angular contour at the anterior end. Shell thick and has well marked ridges on the surface, i.e., 4 to 5 circular ridges and on both sides of these ridges, two more ridges each bent in a loop-like manner, so that the outline of spore in cross section, is very much like of a toothed wheel with nearly equidistant teeth, 16 to 18 in number. Four polar capsules, two slightly larger than the other two. Dimensions: length 16 μ , breadth 10 μ , length of polar capsule 4 μ , length of polar filament 64 μ .

CHLOROMYXUM MAGNUM Awerinzew

[Figs. 136 to 138]

1913 *Chloromyxum magnum* Awerinzew 1913 : 155-156

Habitat: Gall-bladder of *Acanthias blainvillei*;* Algoa Bay, East London, Lüderitzbucht (Africa).

Vegetative form: Ameboid. Body yellowish by the presence of large yellowish granules in endoplasm. Often round or rosary form. Pseudopodia sometimes absent, so that the trophozoites move like *Amoeba limax* with a cluster of small, hairy pseudopodia at the posterior end. In larger form, small round pseudopodia, composed of homogeneous ectoplasm, are formed. Plasmotomy by budding, was often observed. Usually polysporous, rarely monosporous.

Spore: Elongated spherical form. Four polar capsules at the narrow, anterior end. Sporoplasm with two nuclei. Dimensions: length 40 to 48 μ , breadth 30 to 38 μ , length of polar capsules 12 to 15 μ .

* Misprinted in Awerinzew's paper as *blainvillei*.

CHLOROMYXUM FUNDULI Hahn

[Figs. 139 and 140]

1915 *Chloromyxum funduli* Hahn 1915 : 205-206

Habitat: Muscle of *Fundulus* sp.; Woods Hole. In one fish.

Vegetative form: Hahn made observations on few fresh and stained smears. According to him, it is clear that the staining was abnormal. It is hard to quote this here as he used different terms without giving any definition. The reader is advised to consult Hahn's paper.

Spore: Form slightly resembles that of *Chloromyxum quadratum*. Posterior end rounded, the anterior portion narrow and truncated at the tip; optical cross-section thru the posterior part of the polar capsules, circular. Four polar capsules at the anterior end. Dimensions: height (length) 6μ , breadth and thickness 7.5μ respectively.

Remarks: As to the comparison of the present species with *Chloromyxum clupeiidae* Hahn, see p. 94.

CHLOROMYXUM MISGURNI Kudo

[Figs. 141 to 146]

1916 *Chloromyxum misgurni* Kudo 1916 : 6-7

Habitat: Gall-bladder of *Misgurnus anguillicaudatus* Cantor; Tokio (September).

Vegetative form: Round or irregular. Semicircular when viewed from side. From the flat surface, many fine root-like, filiform pseudopodia are extruded. No clear differentiation between ectoplasm and endoplasm. Endoplasm alveolar. Trophozoites always found attached to the lining epithelial cells. Size up to 50μ by 20μ . Polysporous (6 to 8 spores), rarely disporous.

Spore: Spherical, slightly attenuated at the anterior end. Sutural line straight and forms a ridge. Fine longitudinal striations run parallel to the sutural line. Four polar capsules at the anterior end. Sporoplasm finely granular, has two nuclei of equal size. Dimensions: length 8 to 9μ , breadth 6 to 7μ , thickness 5 to 6μ , length of polar capsule 2 to 3μ , of polar filament 28 to 35μ (KOH).

Remarks: The host is often infected at the same time by *Chloromyxum fujitai*, the trophozoites of which can be distinguished from the present form by the structure and the floating habit in the bile. Spores in the two species are decidedly different in form, structure and size.

CHLOROMYXUM FUJITAI Kudo

[Figs. 147 to 152]

1916 *Chloromyxum fujitai* Kudo 1916 : 7-9

Habitat: Gall-bladder of *Misgurnus anguillicaudatus* Cant.; Tokio, (5% of the fish examined in September, found infected).

Vegetative form: Round or irregular. No clear differentiation of protoplasm. Endoplasm highly vacuolated. Ectoplasm being hardly distinguishable. Size up to 40μ in diameter. Trophozoites float in the bile in almost all cases. Disporous and polysporous (up to 8 spores).

Spore: Spherical, often attenuated at the anterior end. Sutural line not straight. Shell very thick, shows thick ridges running longitudinally on the surface. In optical cross section, the spore presents an outline like a cog-wheel with 20 to 22 ridges. The thickness of ridges varies regularly; the thickest ones being located on two lines where a plane perpendicular to sutural plane cuts the shell longitudinally, others decreasing in thickness as they approach the sutural line. Four polar capsules at the anterior end. Sporoplasm with two nuclei. Dimensions: length 10 to 12μ , breadth 8 to 10μ , polar capsules 2 to 3μ , length of polar filament 23 to 30μ (KOH).

CHLOROMYXUM CLUPEIDAE Hahn

[Figs. 153 to 156 and 562 to 565]

1900	Sporozoa	Tyzzar	1900 : 66-68
1901	Sporozoa	Linton	1901 : 438
1910	<i>Chloromyxum</i> sp.	Auerbach	1910 : 178
1917	<i>Chloromyxum clupeiidae</i>	Hahn	1917 : 13-19

Habitat: Body musculature of *Clupea harengus*, *Pomolobus pseudoharengus*, *P. aestivalis*, *P. mediocris*, *Brevoortia tyrannus*, *Stenotomus chrysops*, *Tautoglabrus adspersus*; Woods Hole.

Tyzzar mentioned in his paper and also in a letter to the writer that he collected the material in August of 1900 and that he found the infection occurred only among young fish. Hahn also called attention on the latter fact.

Vegetative form: Hahn's observations are as follows:

Clusters of spores ("pseudocysts") are spindle-shaped, especially when young, usually lying between the bundles of muscle fibres. Color white or creamy. Larger ones usually "in pocket just beneath the integument." Schizogonic multiplication probably exists. Parasites hard to stain, anilin dyes being unable to stain at all. Large form (probably composed of many individuals) 890μ by 30μ .

Tyzzar described as follows: Cysts up to 1 to 2 mm. in length, lying between the muscle fibres of the myotomes, surrounded at times by membranous connective tissue. The parasites also occur in diffused infiltration.

Linton found two cysts, 1.74mm. by 1.16mm. and 1.16mm. by 0.58mm. and also diffused state between the fibrillae.

The writer's observation on slides prepared by Dr. Tyzzar* is as follows: Two cysts in sections; one almost spherical, 480μ by 430μ , sur-

*The writer had recently the opportunity of examining the slides of the parasites prepared by Dr. Tyzzar, which occasion he appreciated very much. As a result of this, the writer became convinced of the identity of forms observed by Tyzzar, Linton and Hahn, tho he could not examine the latter authors' specimens.

rounded by several layers (about 10μ thick) of connective tissue of the host, the other oval, 120μ by 110μ . The staining sufficed to reveal only indistinct structure of the parasites. The homogeneous ectoplasm surrounds the entire surface of the body as a uniform, but very thin layer. Endoplasm granular, filled with spores of remarkably identical stages of development. Isolated spores, also, occur in the muscle bundle in the state of diffused infiltration. Polysporous.

Spore: Hahn describes it as follows: Low conical pyramid with round base; square with bulging sides. No indication of valves in the spore shell. Dimensions: height (length) 5μ , breadth and thickness 7μ , polar capsule 2μ by 1μ .

Linton's form: squarish in outline with rounded corners, 7μ in diameter.

Tyzzler describes his form as follows: Quadrilateral in anterior end view; oval in side view. The four corners are a little protuberant and are directed slightly forward. Shape varies considerably in different species of host. The corners of the spore from *Stenotomus chrysops*, are greatly drawn out, exhibiting stellate form. Four polar capsules radiating from the anterior extremity toward the four corners. Shell shows four furrows radiating from the anterior extremity outwards to the side. Sporoplasm occupies extracapsular cavity. Polar filaments are extruded under the action of acetic acid. Dimension: breadth 7 to 7.5μ .

The writer's observations are as follows:

Spores in fixed and decolorized smears. In smear, most of the spores are seen lying on the base exposing the anterior end view toward the observer's eyes, a few lying with the sutural diameter parallel to the surface of the slide. Form quadrilateral with corners more or less drawn out in anterior end view; oval, with concave posterior side in front view (Figs. 562 to 564). Shell apparently thin but was not clearly separated from the sporoplasm which is finely granular and fills the extracapsular cavity of the spore. Four polar capsules of nearly same size and pyriform. Coiled polar filament indistinct. When stained, the polar capsules stained deeply. It is remarkable to see almost all of the spores exhibit four deeply stained nuclei of capsulogenous cells, which in ordinary case disappear as the spore matures. Dimensions: height (length) 4 to 4.75μ , breadth and thickness 5.4 to 6.5μ , polar capsule about 1.5μ by 0.75μ .

Remarks: Thus the forms of Tyzzler, Linton and Hahn had better be treated as one and the same species. As to the distinction of *Chloromyxum funduli* and the present species, the writer is unable to make it clear as he could not examine the preparation of the former species and especially as he observed some intermediate forms between these two forms in Dr. Tyzzler's preparations of the present species.

CHLOROMYXUM GRANULOSUM Davis

[Figs. 157 and 158]

1917 *Chloromyxum granulorum* Davis 1917 : 237

Habitat: Urinary bladder of *Tylosurus marianus*; Beaufort (July, August).

Vegetative form: Elongated when first placed on the slide, but soon becomes contracted and motionless; progressing by very slow ameboid movements. Ectoplasm usually undistinguishable, being noticed only in a few individuals which had formed one or two short, lobose pseudopodia of hyaline ectoplasm. Body colorless to light yellow. After being on the slide for some time rounded trophozoites often became surrounded by a distinct ectoplasmic layer. Entire body usually coarsely granular, the granules varying greatly in size and shape; sometimes indistinctly vacuolated. Fat globules also present. Size of rounded trophozoites about 30 μ . Disporous and polysporous.

Spore: Spherical, with four distinct ridges on the posterior half of each valve converging toward the anterior end. Sutural ridge distinct. Polar capsules pyriform and convergent. Dimensions: diameter 7 μ , polar capsules 2 μ .

Remarks: Trophozoites from some fish were all colorless, while the larger trophozoites from others were distinctly yellow.

CHLOROMYXUM TRIJUGUM nov. spec.

[Figs. 159 to 182]

Habitat: Gall-bladder of *Lepomis megalotis* Raf.; Stony Creek, and Homer Park, Ill. (November). The parasite was only found in this species, *Lepomis humilis* and *L. cyanellus* seined at the same time being free from the infection. Six specimens, three from each of the above mentioned localities, harboured abundantly both free spores and trophozoites of various stages of development in the bile. The fish, from 6.5 to 10.5cm long, were normal in external appearance and the bladders did not show any particular abnormality, compared with those of other fish, as is usually the case.

Vegetative form: Trophozoites float usually free in the bile, younger forms are most frequently attached to the epithelium of the bladder. Form extremely polymorphous, manifesting various shapes such as, almost circular, rounded, oval, elongated or irregular, which is chiefly due to the active extrusion and retraction of the pseudopodia from the body surface. Body is highly transparent and colorless in both the young and the adult. The differentiation of protoplasm into ectoplasm and endoplasm, is distinctly visible in vivo as well as in stained preparations, especially in larger forms (Figs. 159 to 165). The endoplasm presents an alveolar structure without

any enclosure except the nuclei and various stages of spore formation (Figs. 161, 165, 168 to 171), the alveolar network being smaller at the periphery than in the center. The ectoplasm is a hyaline, transparent and homogeneous layer, free from any coarse granulation in fresh conditions. It shows, however, a very fine reticular structure in stained preparations. The pseudopodia are of two kinds in form, always, formed of ectoplasm alone: the filose and bristle-like form, sometimes branching and protruding from the entire surface or from a localized part of the body, vary in length from 0.5 to 4 μ according to the size of the individual (Figs. 159, 161, 164). This form developed, sometimes, into a thicker form with two to four branched finer processes. The blunt, lobose pseudopodium formed at a localized part of the body is well recognizable in larger individuals. Frequently the filose and the lobose pseudopodia are formed on a trophozoite at the same time. The movements of the blunt pseudopodia were striking in some specimens. At the beginning of the observation, ten minutes after the bile was removed from the host, two club-shaped pseudopodia (Figs. 161 to 163) which were extruded from a trophozoite, the largest diameter of which being 20 μ , moved very actively in the semicircular area changing their forms, showing maximum length of 20 μ . In about thirty minutes, they were retracted and from the same place, a short, oval-shaped pseudopodium was seen to be extruded, which remained in the same position for some time without great change of form (Fig. 164). In another case, a trophozoite with a very broad and rounded pseudopodium extruded actively two to three rounded smaller processes at its extremity (Figs. 165 to 167). After fifteen minutes the pseudopodium was retracted, the ectoplasm forming a uniformly thick layer around the endoplasm. The observations were done at room temperature in hanging drop preparations, sealed with vaseline and paraffin, by using comp. oc. 12 and apo. imm. ob. 2mm., which caused no mechanical pressure upon the parasites. The change of form and especially that of pseudopodia, was clearly observed for one hour and twenty minutes under the above mentioned conditions after the bile was removed from the host. The trophozoites when kept for sixteen hours at room temperature, underwent degeneration and disintegrated, setting free the spores which were formed in them.

No active multiplication by plasmotomy, was observed *in vivo*. In fixed preparations, however, forms that suggested the occurrence of the process in the present myxosporidian, were recognized. As was stated before, the pseudopodia are always formed of the ectoplasm and as each portion of these dividing forms has many nuclei, the author is inclined to record the presence of plasmotomy in the present form.

Size varies greatly. The monosporous form 10 μ by 14 μ , disporous 15 μ by 25 μ and polysporous 30 μ by 50 μ , the largest individual, developing and containing more than 200 spores, was 300 μ by 50 μ .

Spore: Generally circular in front view; oval in side view. Shell comparatively thick, consequently the coiled polar filament is frequently indistinct. Sutural ridge straight and distinct. Each valve has a thick straight, sometimes slightly zigzag-form ridge that runs parallel to the sutural line, so that in side view, three distinct ridges encircling the spore are recognized (Figs. 177 and 180). From each of these two ridges, eight to twelve short ridges are directed toward the center of each valve, which can distinctly be observed on the spores stained with Heidenhain's iron hematoxylin (Figs. 179 and 180). They can be seen as faint markings rising from the margin directed toward the center of the spore, in front view of fresh spores. Four pyriform polar capsules of slightly different size open their foramina independently at the anterior end of the spore (Figs. 178 and 181). The sporoplasm, granular and finely reticular, shows almost always two nuclei when stained. Dimensions in vivo: length and breadth 8 to 10 μ , thickness 5 to 7 μ , polar capsules 3 to 5 μ by 2 to 3 μ , length of polar filament 32 to 40 μ (H₂O₂, KOH).

Remarks: In carefully made smear of the bile, a number of empty spores which had been seen in fresh hanging drop preparations, and often spores, in which the sporoplasm with two elongated nuclei seemed to leave the shell (Fig. 182), were recognized. As this particular spore was found close to a thicker mass of the wall of the gall-bladder in the smear, it can hardly be thought that the mechanical pressure during the preparation lead to the mission of the sporoplasm from the spore. It is possible, on the other hand, to think that this is one of the cases of the germination of the spore in the host in which they were developed, as was reported by the author in *Nosema bombycis* Nägeli (Kudo, 1916).

CHLOROMYXUM CATOSTOMI nov. spec.

[Figs. 560 and 561]

Habitat: Gall-bladder of *Catostomus commersonii* Lac.; Salt Fork, Urbana, Ill. (October). Four fish, from 8 to 14cm.; apparently normal.

Vegetative form: Form usually rounded, with filiform pseudopodia. Majority attached to the epithelium, a few being free in the bile. Body colorless. Protoplasm is not well differentiated. Endoplasm occupying the entire body is of granular structure with vacuoles and refringent spherules. Size: from 15 to 35 μ . When kept for 16 hours in a refrigerator, the trophozoites liberated the spores. The number of spores in each trophozoite is usually 2 or 3, rarely 5 to 6. Active plasmatomic multiplication observed when examined. Spores were comparatively small in number, while the trophozoites were attached abundantly to the epithelium of the gall-bladder. Disporous and polysporous.

Spore: Form approximately spherical in front view; oval in profile. Shell with very fine striations which run parallel to the sutural ridge that is fairly well marked. Rounded polar capsules almost of same size, have

independent openings at the anterior end. Coiled polar filament indistinct. Abnormal spores with five polar capsules are sometimes seen. Dimensions of fixed spores: length 8μ , breadth 7μ , thickness 5 to 6μ , polar capsules 2 to 2.5μ by 1.5μ .

CHLOROMYXUM WARDI nov. spec.

[Figs. 632 to 642]

Habitat: In the gall-bladder of *Oncorhynchus nerka*: Klutina Lake, Alaska (August). A single gall-bladder collected and preserved in formol by Professor Ward, was found to harbor the present species. The study was done on preserved material and on stained smear preparations.

Vegetative form: Young trophozoites (Fig. 632) show ameboid form, and are mostly multinucleated. The protoplasm is not well differentiated either in unstained material or in stained specimens. It is granulated thruout the body, and is vacuolated at places. The smallest form measured was 18μ in largest diameter. The shape of the body suggests its possession of ameboid movements when alive, altho the writer could not examine fresh specimens. Large trophozoites in which the spore formation had partly been completed are generally rounded with reticular protoplasm. Size varies to some extent. The trophozoite shown in figure 633 contains six mature spores and is 23μ in largest diameter. The largest one found was 38 by 30μ , showing ten spores and nuclei. Each spore appears to develop independently from a single sporoblast. Disporous and polysporous.

Spore: Rounded pyramidal in front view (Figs. 640 and 641); circular in transverse section (Fig. 638). The shell is thickened near the posterior margin (Figs. 640 and 641). Sutural line is not straight, the ridge being fairly distinct. The striations on the shell vary to a considerable extent (Figs. 634 to 637 and 639). Four polar capsules at the anterior end, mostly unequal in size and shape. The coiled polar filament is invisible in formol material. Potassium hydrate solution does not cause its extrusion in the preserved spores. The sporoplasm is finely granular with two nuclei. Dimensions of unstained preserved spores: diameter 7.5 to 9μ , polar capsule 3 by 2.5μ .

Remarks: The writer was able to study forty specimens of gall-bladder of Alaskan fishes, chiefly of salmon, which have been collected by Professor Henry B. Ward, during the summer of 1919, for which he wishes to express his deepest appreciation. The examination of these specimens showed that myxosporidia were found only in one of the gall-bladders, and that specimen presented a fairly heavy infection of the present species.

Family SPHAEROSPORIDAE Davis

1917 *Sphaerosporidae*

Davis

1917 : 219

The characters of the family are described on page 57.

Genus SPHAEROSPORA Thélohan

1892	<i>Sphaerospora</i>	Thélohan	1892 : 167
------	---------------------	----------	------------

The characters of the genus are described on page 57.

Type species: *Sphaerospora divergens* Thélohan.

SPHAEROSPORA DIVERGENS Thélohan

[Figs. 183 to 186]

1895	<i>Sphaerospora divergens</i>	Thélohan	1895 : 339-340
1912	<i>Sphaerospora divergens</i>	Parisi	1912 : 289
1912	<i>Sphaerospora divergens</i>	Auerbach	1912 : 41-42

Habitat: Urinary tubules of kidney of *Blennius pholis* L., *Crenilabrus melops* L., *C. pavo* Cuv. et V, and urinary bladder of *Hippoglossoides limandoides*; Concarneau, Roscoff, Napoli (July), Smalfjorden.

Vegetative form: Rounded discoidal or spherical or more or less elongate. Ectoplasm transparent, without real pseudopodium. Movements extremely slow. Endoplasm, granular, contains fat globules and small yellowish granules. Size of sporulating individuals: 65 μ by 55 μ , 60 μ by 25 μ , 60 μ by 20 μ , etc. Polysporous (Thélohan); monosporous, and disporous (Auerbach).

Spore: Spherical. Shell with fine striations. Two polar capsules divergent; coiled polar filament visible in fresh state. Sporoplasm fills the extracapsular cavity of the spore. Dimensions: 10 μ in diameter, often 10 μ by 12 μ , the larger diameter coinciding with sutural plane, thickness 8 μ (Auerbach), polar capsules about 4 μ long, length of polar filament 20 to 25 μ .

SPHAEROSPORA ELEGANS Thélohan

[Figs. 187 and 188]

1890		Thélohan	1890 : 193-209
1892	<i>Sphaerospora elegans</i>	Thélohan	1892 : 167-175
1894	<i>Chloromyxum (Sphaerospora) elegans</i>	Gurley	1894 : 266
1895	<i>Sphaerospora elegans</i>	Thélohan	1895 : 338-339
1909	<i>Sphaerospora elegans</i> *	Auerbach	1909a : 71
1912	<i>Sphaerospora elegans</i>	Parisi	1912 : 289

Habitat: Renal tubules of kidney, connective tissue of ovary and urinary bladder of *Gasterosteus aculeatus* L., *G. pungitus* L., *Lota vulgaris* Cuv., *Phoxinus laevis* L.; Paris, Bretagne, Karlsruhe, Lake Garda.

Vegetative form: Rounded or slightly elongated, not exceeding 20 to 25 μ in diameter. Protoplasm homogeneous, very finely granular, contains numerous refractive globules, probably of fatty nature. Pseudopodia lobose. Movements slow. Disporous.

*Misprinted as *Sphaeromyxa elegans*.

Spore: Spherical, somewhat attenuated at the anterior end. Sutural ridge present, terminating in a small projection at each end of the spore. Two polar capsules spherical. Coiled polar filament not visible in fresh state. Dimensions: diameter 10μ in average, sutural diameter about 11μ .

SPHAEROSPORA ROSTRATA Thélohan

[Fig. 189]

1895 *Sphaerospora rostrata* Thélohan 1895 : 339

Habitat: Malpighian bodies of kidney of *Mugil* sp.; Roscoff, Le Croisic, Le Vivier-sur-mer, Marseille, Banyuls.

Vegetative form: Not described.

Spore: Subspherical. Shell shows deep longitudinal striations which end in sharp spinous edges at the posterior end. Sutural ridge well marked. Anterior part shows enlargement of quadrangular lamella, which is spinous in side view. Dimensions: 10 to 12μ in diameter, sutural diameter 1 to 2μ longer, length of polar filament 40μ .

Remarks: The parasites cause the degeneration of the Malpighian bodies.

SPHAEROSPORA MASOVICA Cohn

[Figs. 190 to 192]

1902 *Sphaerospora masovica* Cohn 1902 : 628-632

Habitat: Gall-bladder of *Abramis brama* L.; Mauersee.

Vegetative form: Polymorphous, due to active movements. Transparent and colorless, while in motion. Endoplasm highly granular, contains yellowish enclosures. Ectoplasm hyaline, forms a narrow layer around the body, occasionally developing into a blunt lobose pseudopodium. Pseudopodia of two kinds; lobose and filose, also intermediate forms. Filiform pseudopodia are formed and retracted more slowly than the lobose. Plasmotomy is of probable occurrence. Two spores are formed in each pansporoblast. Size variable: 10μ (with no spore), 18μ (with sporoblasts), 29μ (with 4 sporoblasts), 38μ (with 22 sporoblasts). Disporous(?), poly-sporous.

Spore: Spherical. Sutural ridge well marked. Polar capsules and sporoplasm are comparatively small, the former convergent. By warming the spore, polar filament is extruded and at the same time two filaments ("starren Fäden") are made visible at the anterior part of the sutural plane. Sporoplasm with two nuclei, no vacuole being present. Dimensions: diameter 8μ , length of polar filament 38μ , length of sutural filament 14μ .

Remarks: Cohn did not observe free spores in the gall-bladder. He, however, saw many free spores, separated from each other, in the intestine, concluding that the body and pansporoblast membrane of trophozoites, are destroyed in the intestine, setting the spores free.

SPHAEROSPORA PLATESSAE Woodcock

[Figs. 193 and 194]

1904 *Sphaerospora platessae* Woodcock 1904 : 59-60Habitat: Otic-capsule of *Pleuronectes platessa* L.; England.

Vegetative form: Cysts opaque masses about 1mm. in diameter. The cartilage was greatly hypertrophied. Polysporous (presumably).

Spore: Spherical. Shell unornamented. Two polar capsules. Sporoplasm with several refractive granules, but without any vacuole. Dimensions: diameter 8 to 9 μ , length of polar filament about 70 μ .

Remarks: Woodcock placed this species provisionally in the genus as he could not examine any fresh material, but had studied smears only.

SPHAEROSPORA ANGULATA Fujita

[Figs. 195 to 197]

1912 *Sphaerospora angulata* Fujita 1912 : 261-262Habitat: Kidney of *Cyprinus carpio* L., *Carassius auratus* L.; Sapporo (Nippon).

Vegetative form: Only description: "The number of the spore in the sporoblast is in this case always less than in the others, rarely exceeding two."

Spore: Somewhat triangular, with convex sides, oval in sideview. Slightly pointed at the mid-posterior margin of the spore. Shell very thin, faintly marked with concentric striations. Two oblong polar capsules are of unequal size. Dimensions: length 7 to 8 μ , breadth 6 to 7 μ , thickness 5 μ , length of largest polar capsule 3.8 μ , length of polar filament twice as long as that of the spore.

SPHAEROSPORA POLYMORPHA Davis

[Figs. 198 and 199]

1917 *Sphaerospora polymorpha* Davis 1917 : 231-232Habitat: Urinary bladder of *Opsanus tau*; Beaufort (June, July).Vegetative form: Elongate, but never very irregular in shape. Slowly ameboid. Body colorless. Ectoplasm clearly seen in younger forms, forming one to several large lobate pseudopodia, which in turn extrude several short, conical pseudopodia. In larger forms, ectoplasm is, often, recognizable only at ends of pseudopodia, which in such cases are composed chiefly of endoplasm. Endoplasm granular, vacuolated in some smaller forms, but in larger individuals vacuoles are indistinct or absent; small fat globules abundant in large forms; numbers of rounded sporoblast cells can be distinctly seen. Size of large trophozoites 35 μ by 50 μ . Disporous and polysporous (polysporous forms rarely contain many spores at the same time).

Spore: Spherical, sometimes slightly compressed infero-superiorly. Sutural ridge; on each side are a number of concentric striations extending around each valve parallel to sutural line. Polar capsules pyriform and large. Coiled polar filaments indistinct. Sporoplasm finely granular. Dimensions: diameter 7 to 10 μ , (8 μ in average), polar capsules 4 to 5 μ by 2 to 2.5 μ .

SPHAEROSPORA sp. Davis

1917 *Sphaerospora* sp. Davis 1917 : 213

Habitat: Urinary bladder of *Lepisosteus platystomus*; Gainesville, Fla.

Vegetative form: No description.

Spore: Not described.

SPHAEROSPORA sp. Southwell et Prashad

1918 *Sphaerospora* sp. Southwell and Prashad 1918 : 347-348

Habitat: Under the scales of *Barilius barna*; from the vicinity of the Ruby Mines, Burma (June).

Vegetative form: The cysts occurred in very large numbers, one under each scale.

Spore: Authors' description: "The poor condition of the material did not allow of a complete account of its structure, but the bicapsulate, rounded structure of its spores places it undoubtedly in the genus *Sphaerospora* Thélohan."

SPHAEROSPORA CARASSII nov. spec.

[Figs. 200 to 204]

Habitat: Gill filament of *Carassius carassius* L.; Tokio (February).

Vegetative form: Trophozoites small ameboid in groups or in diffused condition in the connective tissue of the gill filament. No cyst formation. The number of trophozoites in groups is generally small. The largest group found in sections was 96 μ by 36 μ , the macroscopical examination always failing to trace the parasites. The trophozoites, 10 to 20 μ long, with poorly differentiated protoplasm and usually reticular endoplasm without any particular enclosure (Fig. 200). Ameboid movements not observed. Schizogonic multiplication rapid, each of the daughter individuals developing into two spores. Disporous. Other sporous characters could not be determined.

Spore: Spherical in front and side views, the form variable to some extent (Figs. 201-203). Shell smooth. Sutural ridge fairly distinct. Two polar capsules, broadly pyriform, of equal size and convergent, located at

the anterior end, one on each side of the sutural plane. Coiled polar filament highly distinct (5 to 6 times) in vivo. Sporoplasm granular, shows two nuclei when stained; no vacuole of any nature. Dimensions in vivo: diameter 8 to 13 μ , polar capsules 4 to 5 μ by 2.5 to 3.5 μ , length of polar filament 35 to 40 μ (KOH or pressure).

Remarks: No species of the genus, has ever been found in the branchiae. The characters of the spore, however, compel the writer to place the form in the present genus.

Genus SINUOLINEA Davis

1917 *Sinuolinea* Davis 1917 : 219

The characters of the genus are described on page 57.

Type species: *Sinuolinea dimorpha* Davis.

SINUOLINEA DIMORPHA Davis

[Figs. 205 to 213]

1916 *Sphaerospora dimorpha* Davis 1916 : 333-377
1917 *Sinuolinea dimorpha* Davis 1917 : 232-233

Habitat: Urinary bladder and ureter of *Cynoscion regalis*; Beaufort.

Vegetative form: Disporous and polysporous trophozoites differ distinctly from each other. Disporous trophozoites irregular, colorless, transparent and show slow movements. When attached to the epithelium, rounded with one to several pseudopodia. Differentiation of protoplasm distinct. Occasionally endoplasm contains one or more erythrocytes. Average diameter of full-grown form 25 to 30 μ .

Polysporous form: when attached to the bladder epithelium, the free end is drawn out into a long, cylindrical process, covered with numerous short, hairlike ectoplasmic processes. While not movable, these processes are readily absorbed and reformed. When the trophozoite is detached from the epithelium, the larger end gives rise to numerous conical or arborescent pseudopodia, by means of which the trophozoite moves slowly. Endoplasm extends into the proximal portion of large pseudopodia. It is granular and vacuolated, contains numerous fat globules, refractive granules, yellowish crystals (hematoidin?) and erythrocytes in various stages of disintegration. Endoplasm also contains gemmules, each composed of outer layer and finely granular central portion. Size varies greatly: up to 575 μ by 90 μ .

Spore: Spherical. Sutural ridge well marked. Polar capsules large and spherical. Sporoplasm forms a rounded granular mass. Dimensions: diameter 15 μ , diameter of polar capsules 4.5 μ , length of polar filament 27 to 35 μ .

SINUOLINEA CAPSULARIS Davis

[Figs. 214 to 216]

1917 *Sinuolinea capsularis*

Davis

1917 : 233

Habitat: Urinary bladder of *Paralichthys albiguttus*, *P. dentatus*, *Spheroides maculatus*; Beaufort (July, August).

Vegetative form: Rounded to irregular shape. Body colorless or light yellow. Progressive movements slow. Pseudopodia large branched or arborescent, formed entirely of ectoplasm. Ectoplasm transparent and usually granular, merging gradually with the endoplasm. Endoplasm contains numerous fat globules. In large trophozoites, gemmules are observed. The gemmules are more finely granular and more transparent than the surrounding protoplasm and are practically identical with the small, free trophozoites. Trophozoites containing several gemmules are usually rounded and motionless and appear to be more or less degenerate. Disintegration of such trophozoites were actually observed. Sporulating trophozoites were rare and were never seen to contain gemmules. Size up to 40μ in diameter. Disporous and polysporous(?).

Spore: Spherical, sometimes slightly elongated. Sutural plane much twisted on its axis. Sutural ridge very distinct. Polar capsules and capsulogenous cells large occupying more than one-half of the cavity of spore. Coiled polar filament distinct. Sporoplasm granular contains numerous fat globules. Dimensions: diameter 12 to 14μ , diameter of polar capsules 4.5μ , length of polar filament 50μ .

SINUOLINEA ARBORESCENS Davis

[Figs. 217 and 218]

1917 *Sinuolinea arborescens*

Davis

1917 : 233

Habitat: Urinary bladder of *Siphostoma floridae*; Beaufort

Vegetative form: Rounded or irregular. Body colorless or light yellow. Actively ameboid, forming large arborescent pseudopodia of ectoplasm. Ectoplasm well developed, hyaline and homogeneous. Endoplasm coarsely granular, sometimes containing a few fat globules. Larger trophozoites are less active and the ectoplasm less distinct. In sporulating trophozoites the ectoplasm may entirely disappear, the entire trophozoite consisting of a coarsely granular mass. Diameter of rounded sporulating trophozoites 75μ . Polysporous.

Spore: Rounded, in front view, slightly elongated in the anterior end view. Polar capsules large. Sutural ridge prominent, makes a characteristic S-shaped turn on the anterior end. Coiled polar filaments distinct. Dimensions: length 15μ , breadth 12μ , diameter of polar capsules 5μ .

SINUOLINEA OPACITA Davis

[Fig. 219]

1917 *Sinuolinea opacita* Davis 1917 : 234

Habitat: Urinary bladder of *Paralichthys albiguttus*; Beaufort (August).

Vegetative form: Rounded or slightly irregular. Body colorless and opaque. Movements slow. Pseudopodia short lobose. Ectoplasm not distinct, except around ends of pseudopodia, where it forms a thin hyaline layer. Endoplasm opaque, finely granular, with numerous greenish-yellow fat globules varying greatly in size. Diameter of rounded sporulating trophozoites 22μ , exceptionally large trophozoites 100μ . Disporous.

Spore: Nearly spherical, with flattened, lateral appendages extending from the posterior side. Sutural plane slightly twisted on its axis. Sutural ridge distinct. Polar capsules large. Coiled polar filament distinct. Sporoplasm finely granular, containing several comparatively large fat globules. Dimensions: diameter 12 to 13μ , diameter of polar capsules 4μ .

SINUOLINEA BRACHIOPHORA Davis

[Fig. 220]

1917 *Sinuolinea brachiophora* Davis 1917 : 234

Habitat: Urinary bladder of *Paralichthys albiguttus*; Beaufort (August only in one fish).

Vegetative form: Rounded to somewhat irregular. Body colorless. Ectoplasm hyaline. Endoplasm granular, with numerous large fat globules. Disporous.

Spore: Nearly spherical, with a long lateral appendage from each valve. These appendages are empty except at extreme distal end, which contains a granular mass, probably the remains of the parietal cell. Sutural plane slightly oblique to longitudinal axis. Sutural ridge distinct. Polar capsules and capsulogenous cells large, occupying more than half of cavity of spore. Sporoplasm finely granular. Dimensions: length exclusive of appendages 9 to 11μ , length of appendages 18 to 22μ , breadth of spore 9μ , diameter of polar capsules 3.5μ .

Remarks: Davis mentions that in many respects this species is very similar to *S. opacita*, which occurs in the same host.

Suborder PLATYSPOREA nom. nov.

The definition of the suborder is recorded on page 57.

Family MYXIDIIDAE Thélohan

1892	<i>Myxididées</i>	Thélohan	1892 : 173, 175
1893	<i>Myxidiidae</i>	Gurley	1893 : 412

The characters of the family are described on page 57.

Genus MYXIDIUM Bütschli

1882 *Myxidium* Bütschli 1882 : Pl. 38

The characters of the genus are described on page 58.

Type species: *Myxidium lieberkühni* Bütschli.

MYXIDIUM LIEBERKÜHNI Bütschli

[Figs. 221 to 240]

1854		Lieberkühn	1854 : 5-6, 349
1879		Leuckart	1879 : 246
1881		Bütschli	1881 : 638-648
1882	<i>Myxidium lieberkühni</i>	Bütschli	1882 : 593-595
1883		Balbani	1883 : 201-202, 274-275
1891	<i>Myxidium lieberkühni</i>	Pfeiffer	1891 : 20, 91, 105, 127
1894	<i>Myxidium lieberkühni</i>	Gurley	1894 : 283-289
1895	<i>Myxidium lieberkühni</i>	Thélohan	1895 : 340
1895	<i>Myxidium lieberkühni</i>	Cohn	1895 : 5-36
1898	<i>Myxidium lieberkühni</i>	Doflein	1898 : 229, 341
1902	<i>Myxidium lieberkühni</i>	Prenant	1902a : 200-217
1902	<i>Myxidium lieberkühni</i>	Laveran and Mesnil	1902 : 469-472
1906	<i>Myxidium lieberkühni</i>	Léger and Hesse	1906 : 720
1909	<i>Myxidium lieberkühni</i>	Auerbach	1909a : 71
1912	<i>Myxidium lieberkühni</i>	Schröder	1912 : 326-327
1912	<i>Myxidium lieberkühni</i>	Parisi	1912 : 286
1916	<i>Myxidium lieberkühni</i>	Mavor	1916a : 66-68
1916	<i>Myxidium lieberkühni</i>	Mavor	1916b : 373-378

Habitat: Urinary bladder of *Esox lucius* L., *Lota lota* L. (*L. vulgaris*); France, Canada (Georgian Bay), U. S. A., (Wisconsin, Lake Mendota), Italy (Lago Maggiore, Lago di Como, Milano), Germany.

Vegetative form: Form variable with lobose or immovable filiform pseudopodia. Clear differentiation of protoplasm. Cohn described third layer of protoplasm (mesoplasm). Endoplasm yellowish in older trophozoites, contains yellow globules, fat globules and hematoidin crystals. Size varying with age up to a maximum length of 300 μ by a breadth of 136 μ (Bütschli). Plasmatomous multiplication active. Cohn described budding of larger forms, while Laveran et Mesnil observed only the division of smaller forms. Each pansporoblast develops into two spores. Polysporous.

Spore: Elongated fusiform. Shell with longitudinal striations. Polar capsule at each end of the spore. The longer axis of polar capsules coincides with that of spore. Dimensions: length 18 to 20 μ , width 5 to 6 μ . Mavor's measurement: polar capsules 5 μ by 2.5 to 3 μ , length of polar filament 40 to 45 μ .

MYXIDIUM INCURVATUM Thélohan

[Figs. 241 to 251]

1892	<i>Myxidium ? incurvatum</i>	Thélohan	1892a : 1093-1094
1895	<i>Myxidium incurvatum</i>	Thélohan	1895 : 341
1912	<i>Myxidium incurvatum</i>	Parisi	1912 : 286-287
1912	<i>Myxidium incurvatum</i>	Auerbach	1912 : 4, 39
1916	<i>Myxidium incurvatum</i>	Georgévitch	1916 : 90-91
1917	<i>Myxidium incurvatum</i>	Davis	1917 : 234-235

Habitat: Gall-bladder of *Nerophis aequoreus* L., *N. annulatus*, *N. lumbriciformis*, *Blennius pholis* L., *Callionymus lyra* L., *Fundulus majalis*, *Gambusia affinis*, *Hippocampus brevisrostris*, *Mugil cephalus*, *Scorpaena scrofa* L., *Syngnathus acus* L., *S. typhle*; Roscoff, Concarneau, Marseille, Banyuls, Napoli, Bergen, Monaco, Beaufort (July).

Vegetative form: Thélohan describes as follows: Trophozoites usually small, sometimes reaching a considerable size. Pseudopodia lobose. Protoplasm pale and finely granular with refractive globules. Disporous. According to Parisi and Davis rarely monosporous. Georgévitch observed apparently the polysporous form.

Parisi's form: ectoplasm hyaline, endoplasm granular. Monosporous form 25μ long.

Davis's form: lobose pseudopodia, occasionally being drawn out into a long process. Many trophozoites often cling together closely. Diameter of rounded disporous forms about 13 to 15μ , that of monosporous forms about 10 to 11μ .

Spore: Thélohan's description is as follows: Irregular fusiform. Longest axis curved into S-form, both ends sharply pointed and directed toward opposite directions. Polar capsule opening on opposite side of the spore, in some spores the axis of the polar capsules being parallel to each other. Dimensions: length 8 to 9μ , breadth 4 to 5μ , length of polar filament 10 to 15μ .

Parisi gave the following dimensions: length 10 to 12μ , breadth 5 to 6μ , length of polar capsule 3μ , length of polar filament 28μ .

According to Georgévitch, young spores are not curved (Fig. 245).

Davis's form; Polar filaments when extruded in HCl remained tightly coiled. Dimensions: length 8 to 9μ , width 5 to 6μ , diameter of polar capsule about 3μ .

Remarks: As are shown in figures, Davis's form seems to be somewhat different from the European forms.

MYXIDIUM SPHAERICUM Thélohan

[Fig. 252]

1895	<i>Myxidium sphericum</i> (corr. <i>sphaericum</i>)	Thélohan	1895 : 341-342
------	---------------------------------------------------------	----------	----------------

Habitat: Gall-bladder of *Belone acus* (*Belone belone* L.); Banyuls, Le Vivier-sur-Mer.

Vegetative form: Trophozoites spherical or subspherical, not exceeding 20 to 22 μ in diameter with lobose pseudopodia formed from the entire surface. Endoplasm granular, contains small refractive granules. Disporous.

Spore: Form similar to *M. incurvatum*, but much greater. Coiled polar filament distinctly visible in fresh spore. Dimensions: length 15 to 20 μ , width 7 to 8 μ , length of polar filament 60 μ (KOH).

MYXIDIUM HISTOPHILUM Thélohan

[Fig. 253]

1895	<i>Myxidium histophilum</i>	Thélohan	1895 : 341
------	-----------------------------	----------	------------

Habitat: Connective tissue of kidney and ovary of *Leuciscus phoxinus* L. (*Phoxinus laevis* Ag.); France.

Vegetative form: Small mass.

Spore: Fusiform, being compressed at the middle part. Shell with longitudinal striations. Length of the spore 15 μ .

MYXIDIUM sp. Gurley

[Fig. 254]

1851		Leydig	1851 : 226, 234
1852		Leuckart	1852 : 436
1894	<i>Myxidium</i> ? sp. incert.	Gurley	1894 : 290
1899	<i>Myxidium</i> sp.	Labbé	1899 : 92

Habitat: Gall-duct of *Raja batís* L.

Vegetative form: No description.

Spore: Not described. One figure.

MYXIDIUM DANILEWSKYI Laveran

[Figs. 255 to 257]

1887		Danilewsky	1887 : 35
1897	<i>Myxidium danilewskyi</i>	Laveran	1897 : 725-726
1898	<i>Myxidium danilewskyi</i>	Laveran	1898 : 27-30

Habitat: Kidney of *Emys orbicularis* L.; France.

Vegetative form: Form elongated, circular in cross-section, tapering toward the ends. Body of greenish color, occupying the lumen of the renal tubules of the kidney. Body bent along the cavity of the tubule. Endo-

plasm granular, ectoplasm covering the entire surface of the body as a thin layer. Each pansporoblast develops two spores. Polysporous.

Spore: Elongated fusiform, similar to *M. lieberkühni*, but much smaller. Polar capsule at each end, extrudes filament under the action of nitric acid. Sporoplasm granular with one nucleus. Dimensions: length 12μ , breadth 3 to 4μ .

MYXIDIUM GIGANTEUM Doflein

[Fig. 258]

1898 *Myxidium giganteum* Doflein 1898 : 285-286

Habitat: Gall-bladder of *Raja asterias*; Napoli.

Vegetative form: Rounded trophozoites. Lobose pseudopodia with slow movement, show remarkable dimensions. Posterior portion forms "Stemm-pseudopodien." Small form club-shaped. Endoplasm is of yellowish color. Diameter of large form 500μ , of medium sized 200μ , small individuals $70-90\mu$, quite young ones, polymorphous 8 to 40μ . Larger individual up to 700μ by 180μ . Many trophozoites form a cyst-like motionless stage, in which many individuals seem to be covered with a common gelatinous envelope. Each pansporoblast forms two spores. Polysporous.

Spore: Elongated. Fusiform in front view; in side view, one valve arch-form, the other being flat. Transparent. Two polar capsules, one at each end. Coiled polar filament is clearly seen in larger polar capsules. Dimensions: length 28μ , breadth 8μ , polar capsules 8μ by 4μ .

MYXIDIUM BARBATULAE Cépède

1906 *Myxidium barbatulae* Cépède 1906 : 67
1906 *Myxidium barbatulae* Cépède 1906a : 15-16

Habitat: Kidney of *Cobitis barbatula* L.; Isère.

Vegetative form: Trophozoites form cysts. Form and size vary greatly. Average size: 400 to 500μ in length and 200μ in breadth.

Spore: Irregular fusiform. Polar capsule at each end of the spore. Shell longitudinally striated, number being variable. Dimensions: length 12 to 15μ , breadth about 6μ , polar capsules 5μ by 2.5 to 3μ .

MYXIDIUM GIARDI Cépède

[Figs. 259 to 261]

1906 *Myxidium giardi* Cépède 1906a : 16; 1906b : 170-173
1908 *Myxidium giardi* Cépède 1908 : 93-95
1908 *Myxidium giardi* Cépède 1908a : 8

Habitat: Kidney of *Anguilla vulgaris* Flem.; Wimereux (August).

Vegetative form: Subspherical white cysts, 800 to 900μ in diameter, surrounded by a thick (up to 30μ) membrane, composed of the connective tissue of the host.

Spore: Irregular fusiform, greatly enlarged at the middle portion. Plane of symmetry of the spore coincides with the sutural plane. Shell thick with 9 to 11 longitudinal striations on each valve, which are more clearly seen on spores stained with iron hematoxylin. Polar capsule at each end. Coiled polar filament distinct. Sporoplasm finely granular with two nuclei and refringent globules. Dimensions *in vivo*: length 9 to 10 μ , width 5 to 5.6 μ , thickness 4.75 to 5 μ , polar capsules 3.5 μ by 2 μ .

MYXIDIUM PFEIFFERI Auerbach

[Figs. 262 to 265]

1908	<i>Myxidium pfeifferi</i>	Auerbach	1908 : 459-464
1910	<i>Myxidium pfeifferi</i>	Auerbach	1910c : 171-172

Habitat: Gall-bladder of *Tinca vulgaris* Cuv.; Karlsruhe.

Vegetative form: Observations in sections. More or less flattened, disc-form, often enrolled. The ectoplasm finely granular, without large pseudopodia. It is not usually distinguishable from the endoplasm which is highly alveolar and contains numerous nuclei, but no enclosures.

Spore: Form varies to some extent. Similar to *Myxidium lieberkühni*; slightly curved. Shell with fine longitudinal striations. Polar capsules two, one at each end. Polar filament is extruded by adding one drop of water to the smear of the spore, which had been desiccated for 24 hours. Sporoplasm with one or two nuclei, in one case with four small nuclei, which is thought to be an abnormal. Dimensions: length 13 to 18 μ , breadth 5.2 to 5.8 μ , length of polar capsule 5.2 to 6 μ , length of polar filament 45 to 54 μ .

MYXIDIUM INFLATUM Auerbach

[Fig. 266]

1909	<i>Myxidium inflatum</i>	Auerbach	1909 : 72-74
1909	<i>Myxidium inflatum</i>	Auerbach	1909a : 31
1910	<i>Myxidium inflatum</i>	Auerbach	1910c : 172
1912	<i>Myxidium inflatum</i>	Auerbach	1912 : 39

Habitat: Gall-bladder of *Cyclopterus lumpus* L.; Bergen (September).

Vegetative form: Extremely polymorphous. Rounded, spherical, or much elongated. Ameboid movements very active. Differentiation of protoplasm is sharp and clear, which is best observed in individuals in motion; highly hyaline ectoplasm forms very long lobose pseudopodia, into which granular endoplasm flows in slowly. Size variable. Rounded large form 44 to 45 μ in diameter. Fully grown spores are set free from the mother trophozoite in comparatively short time. Spore formation similar to that of *Myxidium bergense*. Disporous and polysporous (5 spores in maximum).

Spore: Very broad compared with the length. The longitudinal axis is curved in S shape. Polar capsule situated in opposite way at each end of the spore. Dimensions: length 20.8 to 23.4 μ , breadth 13 to 15.6 μ , polar capsules 7.8 μ , length of polar filament 90 to 100 μ (KOH).

MYXIDIUM BERGENSE Auerbach

[Fig. 267]

1908		Keysselitz	1908 : 289
1909	? <i>Myxidium sphaericum</i>	Auerbach	1909 : 75-76
1910	<i>Myxidium bergense</i>	Auerbach	1910 : 61
1910	<i>Myxidium bergense</i>	Auerbach	1910c : 172
1912	<i>Myxidium bergense</i>	Auerbach	1912 : 18-39
1915	<i>Myxidium bergense</i>	Mavor	1915 : 30-31

Habitat: Gall-bladder of *Gadus virens* L., *G. aeglefinis*, *G. merlangus*, *Pleuronectes platessa* and *Sebastes viviparus*, *Melanogrammus aeglefinis*; Norway (Bergen), Canada (St. Andrew, July to September).

Vegetative form: Rounded or elongated, as the result of formation of various pseudopodia. Trophozoites partly free, partly attached to the epithelium of the bladder. Size up to 54 μ in diameter. Pseudopodia of two kinds: lobose and long filose, sometimes slightly branched. Mavor observed a cyst-like stage under certain conditions, which, he thinks, may be due to some exceptional conditions of the parasite. Plasmogamy. Monosporous, disporous and polysporous.

Spore: Fusiform. Main axis curved into S shape. Form, roughly speaking, very much similar to that of *M. sphaericum* Thél. Dimensions: length 16.2 to 19 μ , breadth 7 to 9 μ , length of polar capsules 5.4 μ , length of polar filament about three times as that of spore.

MYXIDIUM PROCERUM Auerbach

[Fig. 268]

1910	<i>Myxidium procerum</i>	Auerbach	1910 : 61-62
1910	<i>Myxidium procerum</i>	Auerbach	1910c : 172-173
1912	<i>Myxidium procerum</i>	Auerbach	1912 : 4, 39

Habitat: Gall-bladder of *Argentina silus* As.; Bergen.

Vegetative form: Not observed.

Spore: Greatly elongated and narrow. Sporoplasm with one or two nuclei. Dimensions: length 21.6 to 25.2 μ , breadth 3.6 to 4 μ , length of polar capsule 7.2 μ .

MYXIDIUM MACKIEI Bosanquet

[Figs. 269 to 271]

1910	<i>Myxidium mackiei</i>	Bosanquet	1910 : 436-438
------	-------------------------	-----------	----------------

Habitat: Renal tubules of kidney of *Trionyx gangeticus*; Bombay. Observations on three slides.

Vegetative form: The largest trophozoite 160μ by 27μ . No distinction between ectoplasm and endoplasm could be drawn, except in a few individuals in which there was a cyst-wall. Spores are formed in pairs. Protoplasm with two kinds of nuclei, some vesicular, others smaller and compact. Polysporous.

Spore: Fusiform with rather pointed ends. Shell finely striated. Two comparatively small polar capsules, one at each end. Sporoplasm with one or two nuclei, contains, often, two large vacuoles. Dimensions: length 16μ (a few 17μ), breadth 5μ (many broader than this).

Remarks: The discoverer, J. P. Mackie mentioned that the parasites did not appear to excite any reaction in the tissue of the host, the animal's health being unaffected.

MYXIDIUM MACROCAPSULARE Auerbach

[Figs. 272 and 273]

1910 *Myxidium macrocapsulare* Auerbach 1910 : 440-441

Habitat: Gall-bladder of *Scardinus erythrophthalmus* L.; Karlsruhe.

Vegetative form: Not observed.

Spore: Elongated elliptical when viewed at right angles to sutural plane. Shell somewhat thick with longitudinal striations parallel to the sutural line. In side-view, both ends pointed in diagonally opposite directions. Polar capsules are comparatively large, one at each end, opening at the sharply pointed end. Dimensions: length 10 to 12μ , breadth 6μ , polar capsules 3 to 4μ .

Remarks: No pathological change. Bile was clear.

MYXIDIUM sp. Awerinzew

[Figs. 274 to 276]

1908	<i>Myxidium</i> sp.	Awerinzew	1908 : 33, 43, 45, 55
1909	<i>Myxidium</i> sp.	Awerinzew	1909 : 76, 78, 80, 81
1911	<i>Myxidium</i> sp.	Awerinzew	1911 : 199-204

Habitat: Gall-bladder of *Cottus scorpius*; Aleksandrowsk, North Sea.

Vegetative form: Trophozoites are small. The protoplasm is differentiated into ectoplasm and endoplasm in some specimens. Very active formation of filiform pseudopodia of various length. Degenerating trophozoites, with one or two empty spaces are often noticed. Each spore is formed independently from each other. Monosporous, disporous and polysporous (with three spores).

Spore: Form similar to *Myxidium incurvatum*. Young spores not curved. Dimensions: length 20 to 35μ , breadth 10 to 15μ .

MYXIDIUM DEPRESSUM Parisi

[Figs. 277 and 278]

1912 *Myxidium depressum* Parisi 1912 : 287

Habitat: Gall-bladder of *Citharus linguatula* Gthr.; Napoli (August).
Vegetative form: Not observed.

Spore: Fusiform with greatly attenuated extremities in front view; flattened and curved in S-form in profile. The axis of polar capsules parallel to each other. Coiled polar filament visible *in vivo*. Sporoplasm with two nuclei, occupies the extracapsular cavity of the spore. Dimensions: length 12 to 14 μ , breadth 5.5 to 6 μ , thickness 2.5 to 3 μ , polar capsules 5.5 to 6 μ by 2.3 μ , length of polar filament 30 μ .

MYXIDIUM OVIFORME Parisi

[Figs. 279 and 280]

1912 *Myxidium oviforme* Parisi 1912 : 287-288
1912 *Myxidium oviforme* Auerbach 1912 : 39

Habitat: Gall-bladder of *Apogon rex mullorum* Cuv., *Coris julis* Gthr., *Gadus callarias* L., *Trutta salar* L.; Napoli (August), Norwegian coast.

Vegetative form: Unobserved by Parisi. Auerbach's observation is as follows:

Trophozoites, small ameboid, usually spherical. Size 10 to 12 μ in diameter. Monosporous (probably).

Spore: Oval with rounded extremities, slightly pointed at the foramina of polar capsules. Shell with numerous fine striations running longitudinally. Polar capsules being often invisible, opening a little above and below of the hypothetical horizontal plane. Sporoplasm fills the extracapsular cavity of the spore, leaving little space at the extremities of the polar capsules. Dimensions: length 11 μ , breadth 8 to 8.5 μ , polar capsules 4.5 μ by 3 μ , length of polar filament 30 to 35 μ . Auerbach's measurements: length 12 to 13 μ , breadth 8 to 9 μ , polar capsules about 4 μ long.

MYXIDIUM ANGUILLAE Ishii

[Figs. 281 to 284]

1915 *Myxidium anguillae* Ishii 1915 : 372-382

Habitat: Integument of the side of the body of *Anguilla japonica* Temm. et Sch.; Schizuoka, Nippon (October). Number of the cysts visible to unaided eye, 10 and 9 on the left and the right side respectively.

Vegetative form: Trophozoites form white and sharply contoured cysts. Cysts, spherical or oval, surrounded by a membranous connective tissue (about 2 μ thick) of the host. Protoplasm is clearly differentiated into ectoplasm and endoplasm. Diffuse infiltration also occurs. Size measured along the skin, 1.2 to 2mm. in diameter; in sections 1.174mm. by 0.658mm.

Spore: Form similar to *Myxidium pfeifferi* Auerbach, but rather straight fusiform, rarely slightly bent. In many spores the shell tapers to a sharp point at each end. Shell striated longitudinally, 22 in all (2 sutural ridges?). Two polar capsules, one at each end. Sporoplasm usually with two nuclei. Dimensions: length 9.1μ , breadth 2.8μ , length of polar capsule 3.5μ .

MYXIDIUM sp. Mavor

[Figs. 285 and 286]

1915 *Myxidium* sp. Mavor 1915 : 32

Habitat: Gall-bladder of *Pseudopleuronectes americanus*; New Brunswick (Canada), of rare occurrence.

Vegetative form: Observations in smears are as follows: Spheroidal, with numerous long pseudopodia on one side, which suggests the attachment of the trophozoite to the bladder. Trophozoites without any spore. Pansporoblasts spherical, 15 to 16μ in diameter.

Spore: Spindle shaped. The long axis being slightly bent in S-form. Two pear shaped polar capsules, one at each end of spore. Coiled polar filament visible in fresh state. Dimensions: length 14 to 15μ , breadth 6 to 7.5μ , polar capsules 4μ by 2.5μ , length of polar filament 90 to 95μ (ammonia water).

MYXIDIUM GADI Georgévitch

[Figs. 287 to 290]

1916 *Myxidium gadi* Georgévitch 1916 : 88-89
 1917 *Myxidium gadi* Georgévitch 1917c : 797-799
 1919 *Myxidium gadi* Georgévitch 1919 : 251-289

Habitat: Gall-bladder of *Gadus pollachius*, *Solea vulgaris* Quens; Roscoff (September).

Vegetative form: Highly polymorphous. Spherical or oval. Large forms fill up the bladder. Ectoplasm hyaline and transparent, forming one long or many short lobose pseudopodia. Endoplasm colorless and finely granular, contains more or less large numbers of nucleus. Monosporous, disporous and polysporous.

Spore: Fusiform with attenuated ends. Young spores more attenuated than the fully grown forms. The main axis of the spore coincides with the longitudinal axis of the polar capsules, with slight deviation. Two nuclei of the sporoplasm, are always smaller than those of the shell-valves or of polar capsules. Dimensions: length 6 (?) to 14μ , breadth 4 to 6μ .

MYXIDIUM GLUTINOSUM Davis

[Fig. 291]

1917 *Myxidium glutinosum* Davis 1917 : 235

Habitat: Gall-bladder of *Cynoscion regalis*; Beaufort.

Vegetative form: Elongated or irregular. Slowly ameboid, moving by means of a broad, lobose pseudopodium of hyaline ectoplasm. Body colorless. Ectoplasm only distinct in pseudopodium. Endoplasm finely granular. The mature spores while still within the mother trophozoites, are surrounded by a clear, refractive gelatinous envelope. Diameter of rounded sporulating trophozoites 20μ . Disporous.

Spore: Cylindrical, ends of valves rounded except at one side, where the polar capsules open at the apex of a small, conical elevation. Spore characterized by the presence of a transparent, homogeneous, gelatinous envelope. Polar capsules pyriform, opening on each side nearly at right angles to the longitudinal axis. Dimensions: length 10 to 11μ , breadth 6μ , length of polar capsules 3μ .

MYXIDIUM PHYLLIUM Davis

[Figs. 292 and 293]

1917 *Myxidium phyllium* Davis 1917 : 235

Habitat: Gall-bladder of *Gambusia affinis*; Beaufort.

Vegetative form: Exceptionally large; flattened, leaflike, usually folded on itself; motionless. Pseudopodia were not observed. Ectoplasm forming a distinct transparent layer around entire body. After being on slide for some time ectoplasm usually becomes covered with very numerous, short, hairlike processes. Endoplasm finely granular, contains numerous fat globules. Diameter up to 1.35mm . Polysporous.

Spore: Fusiform, slightly truncated at each end where polar capsules open. Shell with numerous longitudinal striations. Sporoplasm finely granular, with several small fat globules. Dimensions: length 11μ , breadth 8μ , diameter of polar capsules 3μ .

MYXIDIUM STRIATUM Cunha et Fonseca

1917 *Myxidium striatus* Cunha et Fonseca 1917 : 321

Habitat: Gall-bladder of *Menticirrhus americanus* L., *Bairdiella ronchus* Cuv. et Val.; Brazil.

Vegetative form: More or less spherical. Body small and colorless. Endoplasm granular. Ectoplasm visible when pseudopodia are formed. Pseudopodia filiform, being projected radially. Size variable, 16μ in diameter in average.

Spore: Elliptical. Shell with fine longitudinal striations which run parallel to sutural line. Sutural plane oblique to the longitudinal axis of the spore which is thickened at the extremities. Two ovoidal polar capsules, one at each end. Dimensions: length 10 to 14μ , breadth 6 to 8μ , length of polar capsules 4μ , length of polar filament 30μ .

MYXIDIUM KAGAYAMAI nov. spec.

[Figs. 294 and 295]

1916 *Myxidium* sp.

Kudo

1916 : 6

Habitat: Gall-bladder of *Misgurnus anguillicaudatus* Cant.; Tokio (September), 2% of the fish examined infected.

Vegetative form: Not observed.

Spore: Fusiform; one valve being more convex than the other. Suture line straight. Shell with fine longitudinal striations. Dimensions in fixed preparations: length 15 to 18 μ , breadth 6 to 7 μ , length of polar capsules 7 to 8 μ , length of polar filament 60 to 70 μ .

Remarks: Tho the vegetative form is still unobserved, the author is compelled to consider the present form as a new species by careful reexamination of the material and proposes the name in honor of Dr. T. Kagayama, Tokio, Nippon.

MYXIDIUM AMERICANUM nov. spec.

[Figs. 622 to 627]

Habitat: In the lumen of urinary tubules of the kidney of *Trionyx spinifera*; Crystal Lake, Urbana, Ill. (July). A single host specimen showed a light infection in the above mentioned organ. No intracellular stage was detected.

Vegetative form: The young trophozoite in the lumen of the tubule of the kidney is multinucleate, and more or less irregular in shape which suggests the ameboid movements of the animal (Figs. 622, 623). The older form with mature spores is rather spherical in form with a distinct outline. The protoplasm is fairly well differentiated into ectoplasm and endoplasm (Fig. 624). The size of the trophozoites varies from 12 to 25 μ in diameter. A pansporoblast produces two spores. Polysporous.

Spore: Spindle-form; with the two pointed extremities stretched in opposite directions. Circular in cross-section. The shell is rather thin; sutural line is straight. Fine longitudinal striations on the shell, eight to ten in number on each valve. The polar capsules are nearly spherical, coiled polar filament being visible in fresh material (three turns). The polar filament is easily extruded from the fresh spores under the influence of potassium hydrate solution. The direction of the extruded polar filament forms an angle of about 45° with the main axis of the spore and the two filaments are parallel to each other. Preserved spores do not show any filament extrusion under the influence of the said chemical. The sporoplasm is finely granular, and shows, upon staining, two small nuclei of ring-shape, as their peripheral layer takes stain more deeply than the central portion. Average dimensions of fresh spores: length 15 to 16 μ , breadth 5.5 to 6 μ , polar capsule 4 μ by 3.5 μ , length of polar filament 25 to 32 μ .

Remarks: Two species of the genus Myxidium were reported to occur in chelonian hosts; i.e., *M. danilewskyi* (page 109) and *M. mackiei* (page 112). The former differs from the present form in having an elongated vegetative form which is greenish in color, and in having spores of different shape, dimensions and structure, not to speak of the difference of the host. The latter resembles closely to the species under consideration in dimensions of the spores, but differences in the trophozoite and in the structure of the spore do not allow one to consider two forms as identical. The species is therefore treated as new.

Genus SPHAEROMYXA Thélohan

1892 *Sphaeromyxa* Thélohan 1892a : 1091-1093

The characters of the genus are described on page 58.

Type species: *Sphaeromyxa balbianii* Thélohan.

SPHAEROMYXA BALBIANII Thélohan

[Figs. 296 to 307]

1892	<i>Sphaeromyxa balbianii</i>	Thélohan	1892a : 1091-1093
1895	<i>Sphaeromyxa balbianii</i>	Thélohan	1895 : 342
1912	<i>Sphaeromyxa balbianii</i>	Parisi	1912 : 288
1916	<i>Sphaeromyxa balbianii</i>	Georgévitch	1916 : 92-93
1917	<i>Sphaeromyxa balbianii</i>	Davis	1917 : 235-236

Habitat: Gall-bladder of *Motella tricirrata* Bl., *M. maculata* Risso., *Cepola rubescens* L., *Clupea pilchardus*, *Siphostoma floridae*, *S. louisianae*; Roscoff (September), Concarneau, Marseille, Banyuls, Napoli (September), Beaufort (June to August).

Vegetative form: Flattened leaf-like or disc-form, reaching 3 to 4mm. in diameter. Often forms spherical with opaque appearance. The protoplasm is distinctly differentiated into endoplasm and ectoplasm. Ectoplasm forms rounded lobes which exhibit slow movements and show a clear radially striated structure in sections. Endoplasm reticular, contains nuclei, young and mature spores and fat globules. Each pansporoblast develops two spores. Polysporous.

Davis mentions that the largest form he observed was 900 μ in diameter.

Georgévitch recognized a large number of small trophozoites which were formed by repeated plasmotomous multiplication.

Spore: Fusiform, with truncate ends. Shell longitudinally striated. One polar capsule at each end. Polar filament is wound around an imaginary axis perpendicular to the longitudinal axis of the spore. When extruded, the polar filament is seen as a short, conical and hollow thread-like structure. Sporoplasm finely granular with two nuclei. Dimensions: length 15 μ , width 5 μ , length of polar filament 15 μ .

Parisi gave the following dimensions: length 15 to 20 μ , width 5 to 6 μ , polar capsule 7 μ by 4.7 μ , length of polar filament 25 to 30 μ .

Davis' measurements are as follows: length 17 to 20 μ , breadth 5 to 6 μ , length of polar filament 20 μ .

Georgévitch observed young spores with both ends tapering into a point. Later they assume the typical form with truncated ends. He did not recognize the striations on the shell. He also mentions the occurrence of abnormal spores, such as elliptical, spherical forms, etc.; or with only one polar capsule.

SPHAEROMYXA IMMERSA (Lutz) Thélohan

[Figs. 308 to 311]

1889	<i>Cystodiscus immersus</i>	Lutz	1889 : 84-88
1895	<i>Sphaeromyxa immersa</i>	Thélohan	1895 : 343
1899	<i>Cystodiscus immersus</i>	Lühe	1899 : 291-293

Habitat: Gall-bladder of *Bufo marinus* L. and *Leptodactylus ocellatus* L.; Brazil.

Vegetative form: Leaf-like or disc form, visible thru the bladder wall. Upper and lower sides slightly convex. Size up to 1.5 or 2mm. in diameter; thickness being 1/20 to 1/10 of the diameter. Protoplasm is well differentiated. Ectoplasm transparent and membranous, often contains a large number of micrococcus-like bodies. No ameboid movements nor change of form. Endoplasm highly vacuolar, contains fat globules. Plasmotomic multiplication probably occurs. Spores always arranged in pairs. Polysporous.

Spore: Oval with rounded extremities. Shell more or less thick, with fine transverse striations. Spherical polar capsule at each end. Sutural plane is oblique to the longitudinal axis of the spore. Sporoplasm transparent. Dimensions: length 12 to 14 μ , breadth 9 to 10 μ , length of polar filament 50 to 70 μ (4 to 5 times that of the spore) (KOH).

SPHAEROMYXA INCURVATA Doflein

[Figs. 312 to 314]

1898	<i>Sphaeromyxa incurvata</i>	Doflein	1898 : 286-287
------	------------------------------	---------	----------------

Habitat: Gall-bladder of *Blennius ocellatus*; Napoli.

Vegetative form: Trophozoites are found in large masses (Plasmodia ?), in which they form a thin, hollow ball, 5-7mm. in diameter.

As the surface is greater than the inner surface of the bladder, some parts of the body are folded up. Body bluish white and transparent. Protoplasm highly vacuolar, contains numerous fat globules, nuclei and spores. Polysporous.

Spore: Curved to one side along sutural plane and also in a plane at right angles to it. Polar capsule at each end. Sporoplasm with two nuclei.

Polar filament is wound along the longer diameter of the capsule, and relatively thick, but thinner than that of *S. balbianii* Thél. Dimensions: length (along the inner side of the arch) 30 to 35 μ , breadth 8 μ , distance between two polar capsules 12 to 15 μ , polar capsules 12 to 15 μ by 4 to 5 μ .

SPHAEROMYXA SABRAZESI Laveran et Mesnil

[Figs. 315 and 322]

1900	<i>Sphaeromyxa sabrazesi</i>	Laveran et Mesnil	1900 : 380-382
1906	<i>Sphaeromyxa sabrazesi</i>	Schröder	1906 : 455-466
1907	<i>Sphaeromyxa labrazesi*</i>	Schröder	1907 : 359-381
1910	<i>Sphaeromyxa sabrazesi</i>	Schröder	1910 : 1-5
1912	<i>Sphaeromyxa sabrazesi</i>	Parisi	1912 : 288
1913	<i>Sphaeromyxa sabrazesi</i>	Jameson	1913 : 2
1916	<i>Sphaeromyxa sabrazesi</i>	Georgévitch	1916 : 91-92
1916	<i>Sphaeromyxa sabrazesi</i>	Georgévitch	1916a : 3

Habitat: Gall-bladder of *Hippocampus brevisrostris* Cuv., *H. guttulatus* Cuv.; *Syngnathus acus*, *Motella tricirrata*, *Nerophis annulatus*, *Siphonostoma rondeletii*; Arcachon, Rovigno, Napoli, Roscoff (September), Monaco, Villefrance, (March to June).

Vegetative form: Disc form. Diameter up to 2mm. Thickness variable. Body whitish in color. Ectoplasm thin, transparent and homogeneous. Young trophozoites may probably have lobose pseudopodia. Endoplasm highly vacuolated, contains nuclei of various sizes, pansporoblasts, spores and more or less refringent granules. Polysporous.

Schröder observed larger forms up to 5mm. Ectoplasm also was found to project numerous fine short (1 μ) hair-like processes from the surface. Each pansporoblast develops into two spores.

Spore: Cylindrical, bent in arch form; with truncated ends. Large cylindrical polar capsule at each end. Sporoplasm granular, contains one nucleus. Polar filament short and conical, is extruded under the action of nitric acid. Dimensions: length 28 μ , width 4.3 μ , polar capsule 9 to 10 μ by 3 μ , distance between the polar capsules 8 μ , length of polar filament 8 μ .

Schröder noticed the stained sporoplasm contained one or two nuclei. He observed indistinctly marked longitudinal striations on the shell. Dimensions: length 22 to 25 μ , breadth 3 to 4 μ , polar capsule 8 μ by 2 to 3 μ , length of polar filament about 12 μ .

Georgévitch described the presence of a hyaline substance, containing pale granules, in the spore cavity. Young spores were found to take the form of Myxidium type. In mature spores, he always found two nuclei by staining.

*Misprinted in Schröder's paper.

SPHAEROMYXA HELLANDI Auerbach

[Figs. 323 and 324]

1909	<i>Sphaeromyxa hellandi</i>	Auerbach	1909 : 78-79
1910	<i>Sphaeromyxa hellandi</i>	Auerbach	1910b : 772-774
1910	<i>Sphaeromyxa hellandi</i>	Auerbach	1910c : 174-175
1912	<i>Sphaeromyxa hellandi</i>	Auerbach	1912 : 4, 40

Habitat: Gall-bladder of *Molva vulgaris* Flem., *Centronotus gunellus*, *Brosmius brosmæ* Asc.; Bergen, Torghatten.

Vegetative form: Large and rounded disc form. Protoplasm is distinctly differentiated into ectoplasm and endoplasm. Thickness up to 160μ , folded in the bladder. Ectoplasm finely granular; in unstained specimens, it is recognizable as 10 to 12μ thick layer, in which about 2μ thick radially striated outer layer and 8 to 10μ thick inner finely granular region can be distinguished. In stained sections, the outer layer remains unstained. Endoplasm highly alveolar, contains refractive granules of different size which are not stained with Sudan III. Each pansporoblast develops into two spores. Polysporous.

Spore: Arch form in front view. The degree of curvature varies greatly. Sutural line curved in S-shape and well marked. Both ends more or less truncated. Polar capsule at each end. Polar filament being wound along the longest axis of the polar capsule and is extruded with KOH. Sporoplasm rounded with one or two nuclei. Dimensions: length 20.8 to 26μ , breadth and thickness 5.4μ , length of polar capsule 10 to 10.8μ .

SPHAEROMYXA EXNERI Awerinzew

[Figs. 325 and 326]

1913	<i>Sphaeromyxa exneri</i>	Awerinzew	1913a : 155
------	---------------------------	-----------	-------------

Habitat: Gall-bladder of *Thysanophris japonicus*; Lorenço Marques (Africa).

Vegetative form: Not observed.

Spore: Somewhat resembles that of *S. hellandi* Auer. in being bent to one side with sutural line of S-form but differs in dimensions. Both ends slightly tapering. Polar capsules two, one at each blunt end, in which the polar filament is wound parallel to its longer axis. Sporoplasm comparatively small and sharp-contoured, contains only one nucleus. Dimensions: length 75 to 80μ , breadth 18 to 20μ , length of polar capsule 30 to 35μ .

SPHAEROMYXA GASTEROSTEI Georgévitch

[Fig. 327]

1916	<i>Sphaeromyxa gasterostei</i>	Georgévitch	1916 : 88
------	--------------------------------	-------------	-----------

Habitat: Gall-bladder of *Gasterosteus spinachia*; Roscoff (September).

Vegetative form: Trophozoites form large plasmodia.

Spore: Large, elongated fusiform; ends less truncated than those of the spore of *Sphaeromyxa balbianii*. As the spore becomes mature, the ends assume more pointed shapes. Polar capsules two, one at each end. Sporoplasm with two nuclei, fills the extracapsular cavity. Dimensions: twice or three times larger than those of *Sphaeromyxa balbianii* Thélohan.

Genus ZSCHOKKELLA Auerbach

1910	<i>Zschokkella</i>	Auerbach	1910 : 62-63
			1910a : 240-256
			1910c : 175

The characters of the genus are described on page 58.

Type species: *Zschokkella hildae* Auerbach.

ZSCHOKKELLA HILDAE Auerbach

[Figs. 328 to 331]

1910	<i>Zschokkella hildae</i>	Auerbach	1910 : 62-63
1910	<i>Zschokkella hildae</i>	Auerbach	1910a : 240-254
1912	<i>Zschokkella hildae</i>	Auerbach	1912 : 40-41

Habitat: Urinary bladder of *Phycis blennioides* Br., *Gadus aeglefinis*, *G. callarias* L., *G. virens* L.; Norway.

Vegetative form: Trophozoites float in the bile or attach themselves to the epithelial layer of the bladder. Youngest ameboid form about 4.5 to 6 μ . In floating form, pseudopodia more or less long, lobose, are formed; while in the attached form those similar to the pseudopodia of *Myxidium bergense* Auer. are developed. Plasmogamy occurs. Size varies greatly according to the number of spores which are formed in each individual. Monosporous (with or without the remnant of protoplasm), disporous and polysporous (up to 4 spores).

Spore: Semicircular in front view, with slightly and equally attenuated ends. At each end, large spherical polar capsule is situated which opens not at the extremity, but on the flat surface. Shell bivalve and thick. Sutural line S-form. Sporoplasm with two nuclei. Dimensions: length 16 to 28.8 μ , breadth 13 to 18 μ , polar capsules 5.6 to 7.2 μ in diameter, length of polar filament 72 μ (KOH).

ZSCHOKKELLA NOVA Klokačewa

[Figs. 332 and 333]

1914	<i>Zschokkella nova</i>	Klokačewa	1914 : 184-186
------	-------------------------	-----------	----------------

Habitat: Gall-bladder of *Carassius vulgaris*; Russia?

Vegetative form: Not observed.

Spore: Outline irregular. Observations on fixed materials alone. Two large round polar capsules open at the side near ends. Sporoplasm with two nuclei. On some spores, striations that run parallel to the sutural line were observed. Dimensions: length 9.5 to 11.5 μ , breadth 6.5 to 7 μ , diameter of polar capsule 3 to 3.5 μ .

ZSCHOKKELLA ACHEILOGNATHI Kudo

[Figs. 334 to 338]

1916 *Zschokkella acheilognathi* Kudo 1916 : 3-5

Habitat: Gall-bladder and gall-duct of *Acheilognathus lanceolatum* Temm. et Schl.; Tokio (May). Over 80% of the fish examined were found to be infected.

Vegetative form: Disc-shape. In bile duct, large trophozoites are folded up. Body colorless and transparent. Protoplasm is well defined into two regions. Ectoplasm finely granular *in vivo*. In stained sections, it shows two layers; thin outer layer (2μ thick) presents very fine striations, while inner layer (6 to 8μ thick) is finely vacuolated without any enclosure. Endoplasm is highly vacuolated. Lobose pseudopodia formed only in younger individuals (15 to 30μ in diameter), in which ameboid movements are not slow. Size: up to 720μ by 550μ , thickness 5 to 30μ . Polysporous.

Spore: Form resembles *Zschokkella hildae*, but slightly elongated. Form varies to some extent. Some spores are of Myxidium type. Sutural line curved. Longitudinally striated. Spherical polar capsule at each end opening near the extremity. Dimensions: length 10 to 14μ , breadth 6 to 7μ , diameter of polar capsule 3 to 5μ , length of polar filament 65 to 70μ (KOH).

ZSCHOKKELLA GLOBULOSA Davis

[Figs. 339 and 340]

1917 *Zschokkella globulosa* Davis 1917 : 236

Habitat: Urinary bladder of *Spheroides maculatus*; Beaufort (August).

Vegetative form: Rounded; slowly ameboid, forming short, lobose pseudopodia. Body colorless and transparent. Ectoplasm not distinct. Protoplasm granular, characterized by the presence of several large fat globules. Sporulating trophozoites about 15 to 16μ in diameter. Monosporous and disporous.

Spore: Semicircular. Sutural line twisted on its axis and oblique to longitudinal axis; sutural ridge distinct. Polar capsules opening on flat surface. Sporoplasm finely granular and very transparent. Dimensions: length 11μ , breadth 7μ , diameter of polar capsules 3μ .

Family MYXOSOMATIDAE Poche

1913 *Myxosomatidae* Poche 1913 : 230

The characters of the family are described on page 58.

Genus MYXOSOMA Thélohan

1892 *Myxosoma* Thélohan 1892 : 175

The characters of the genus are described on page 58.

Type species: *Myxosoma dujardini* Thélohan.

MYXOSOMA DUJARDINI Thélohan

[Figs. 341 to 343]

1841		Müller	1841 : 486-487
1845		Dujardin	1845 : 644
1892	<i>Myxosoma dujardini</i>	Thélohan	1892 : 175
1895	<i>Myxosoma dujardini</i>	Thélohan	1895 : 343-344
1905	<i>Myxosoma dujardini</i>	Nufer	1905 : 77, 79, 186
1910	<i>Myxosoma dujardini</i>	Wegener	1910 : 72-73
1916	? <i>Myxosoma dujardini</i>	Kudo	1916 : 3

Habitat: Branchial lamellae of *Scardinius erythrophthalmus* L., *Perca fluviatilis*, *Leuciscus rutilus* L. and *Cyprinus carpio* L.; France, Frisches Haff, Kurisches Haff (February, April, May), Tokio (May), Switzerland.

Vegetative form: White cysts being branched, rounded, spherical or irregular; 1 to 1.5mm. in diameter.

Wegener's form 1 to 1.7mm. long.

Spore: Ovoidal, flattened, with attenuated anterior end which is slightly bent laterally. Two pyriform polar capsules at the anterior end. Sporoplasm without any iodophilous vacuole. Dimensions: length 12 to 13 μ , breadth 7 to 8 μ .

Wegener's form: polar capsules 6 μ by 3 μ .

Kudo's form: polar capsules 6 to 7 μ by 2 μ , length of polar filament 70 μ .

MYXOSOMA (?) LOBATUM Nemeček

[Fig. 348]

1911	<i>Myxosoma (?) lobatum</i>	Nemeček	1911 : 160-162
------	-----------------------------	---------	----------------

Habitat: Branchiae of *Leuciscus leuciscus* L. and *Aspius rapax* Ag.; Austria.

Vegetative form: Cysts spherical, oval or elongated; of white color. Size from 0.5 to 3mm. by 0.5 to 1mm. Those in *Aspius rapax*, oval to spindle-shape, 1 to 3mm. long and 1 to 1.5mm. wide.

Spore: Ovoidal; anterior end narrowly pointed and straight; posterior end rounded, with lobose appendix (about 6 μ long). A transverse fold on the shell behind the polar capsules in fresh as well as preserved spores. No iodophilous vacuole. Dimensions: length 12.6 μ , breadth 8.2 μ , length of polar capsule 4.2 μ , length of polar filament 80 to 90 μ . Spores found in *Aspius rapax*, had slight difference in dimensions, the structure, however, being similar to the above.

Remarks: Nemeček doubts if this form is really *Myxosoma* because of the following: 1) different shape of the cysts compared with that of the type species as described by Thélohan; 2) spores observed might develop later into other forms like *Henneguya*.

MYXOSOMA FUNDULI Kudo

[Figs. 344 to 347]

1918 *Myxosoma funduli* Kudo 1918 : 12-14

Habitat: Branchiae of *Fundulus majalis* Wal. and *F. heteroclitus* L.; Woods Hole (August, September).

Vegetative form: Cysts. Spherical and small; 150μ in average diameter. Largest form observed 360μ by 264μ . Spores, young and mature, were found in the cysts. Polysporous.

Spore: Pyriform. Shell uniformly thick with 7 to 10 folds on sutural edge at the posterior portion. Sutural ridge, fairly well marked. Two polar capsules pyriform and of equal size at the anterior end. Sporoplasm finely granular with two nuclei but without any iodophilous vacuole. Dimensions: length 14μ , breadth 8μ , thickness 6μ , polar capsule 8μ by 2μ , length of polar filament 38 to 42μ (perhydrol, KOH).

Remarks: The writer could not find any evidence of an iodophilous vacuole by treatment with various iodine mixtures, which is the most important characteristic of the genus. Hahn's form (*Myxobolus funduli*, p. 151) should be distinguished from the present form.

Genus LENTOSPORA Plehn

1905 *Lentospora* Plehn 1905 : 150

The characters of the genus are described on page 58.

Type species: *Lentospora cerebralis* (Hofer) Plehn.

LENTOSPORA CEREBRALIS (Hofer) Plehn

[Figs. 349 to 354]

1903	<i>Myxobolus cerebralis</i>	Hofer	1903 : 8
1904	<i>Myxobolus chondrophagus</i>	Hofer	1904 : 53
1905	<i>Lentospora cerebralis</i>	Plehn	1905 : 145-166
1909	<i>Lentospora cerebralis</i>	Plehn	1909 : 38
1910	<i>Lentospora cerebralis</i>	Auerbach	1910c : 176

Habitat: Cartilage and perichondrium of *Trutta iridea* Gibb., *Salmo fontinalis* Mitch., *Trutta salar* L.; Germany (Karlsruhe and other localities).

Vegetative form: Ameboid form. Size varies greatly. Small ameboid form probably grows up into large individual which has often fifty or more ringform nuclei and breaks up into numerous small forms by division. No sporous character is observed except a figure of a disporous form.

Spore: Circular in front view; lenticular in side view, with more or less extensive variation in length and breadth. Shell smooth. Sutural ridge distinctly thickened. Two polar capsules pyriform and convergent, are usually of same size. Extruded polar filaments cross each other. Sporoplasm with two ring-form nuclei but without any iodophilous vacuole.

Dimensions: diameter 6 to 10μ , length of polar capsule $2/5$ that of the spore, length of polar filament 40 to 50μ (limewater, 1% KOH).

Remarks: Plehn noticed that the present form causes the chronic form of "Drehkrankheit" among young fish in German waters. She was unable to extrude the polar filament with mineral (?) acids. Auerbach, however, could extrude the filament by means of acids.

LENTOSPORA MULTIPLICATA Reuss

[Fig. 355]

1906 *Lentospora multiplicata* Reuss 1906 : 203

Habitat: Muscle of *Idus melanotus* Heck.; Volga?, Russia.

Vegetative form: Not described.

Spore: Oval. Sutural edge broad with many folds. No iodophilous vacuole. Dimensions: length 12μ , breadth 9.5μ , thickness 6μ , polar capsules 4μ by 2.25μ .

LENTOSPORA ENCEPHALINA Mulsow

1911 *Lentospora encephalina* Mulsow 1911 : 483-485

Habitat: In the blood vessel of the brain, especially of the mid-brain of *Cyprinus carpio* L.; Munich (spring). Blood vessels are the only seat of infection. In most cases many individuals lie parallel to one another. The infection occurs frequently and heavily. The effect, however, is undetermined.

Vegetative form: Trophozoite elongated, worm-like and circular in cross section. The body is covered with a pellicula. The protoplasm is distinctly differentiated into homogeneous ectoplasm layer and inner endoplasm. In the latter are found numerous granules, small nuclei and spores.

Spore: Almost circular in front view; profile? No iodophilous vacuole is found. The polar filament is easily extruded by means of a highly diluted KOH solution. Diameter: 5 to 5.5μ .

LENTOSPORA ASYMMETRICA Parisi

[Figs. 356 to 359]

1912 *Lentospora asymmetrica* Parisi 1912 : 292-293

Habitat: Connective tissue of kidney of *Crenilabrus pavo* C. et V.; Napoli (September).

Vegetative form: One trophozoite found; a small, rounded form with thin and hyaline ectoplasm which could be distinguished from the endoplasm with coarse yellowish globules, containing two spores. Disporous?

Spore: Oval from the front; flattened and fusiform in profile. Sutural edge with many triangular folds, which are more clearly seen in material

preserved in formalin than in fresh condition. Two polar capsules of same size, are situated asymmetrically, opening at the side near the anterior end. Sporoplasm granular and with two nuclei, but without any iodophilous vacuole. The polar filament not being extruded by ordinary reagents, probably because the spores were not full-grown. Dimensions: length 10 to 11 μ , breadth 6.5 to 7 μ , length of polar capsules 5 μ .

LENTOSPORA ACUTA (Fujita) Kudo

[Figs. 360 to 362]

1912 *Sphaerospora acuta* Fujita 1912 : 260-261

Habitat: Epithelium of branchial lamellae of *Carassius auratus* L.; Sapporo, Nippon.

Vegetative form: Fujita's description is simply as follows: Sporoblast contains about two spores.

Spore: Spherical in front view, with slightly pointed anterior end; spindle shaped in side-views. Shell thin and smooth. Two convergent polar capsules are of different sizes, occupying about 5/8 in space of the spore. No vacuole could be made out in sporoplasm. Dimensions: length 8 to 10 μ , breadth 7 to 8 μ , thickness 5 to 6 μ , polar capsules 5 μ by 4 μ .

Remarks: This species, recorded incompletely by Fujita as *Sphaerospora*, shows characters of the genus *Lentospora* in spore form so that it is provisionally placed here.

LENTOSPORA DERMATOBIA Ishii

[Figs. 594 to 596]

1916 *Lentospora dermatobia* Ishii 1916 : 472-474

Habitat: In the integument of *Anguilla japonica* Temm. et Schl.; Shizuoka, Nippon. From the same specimen which harboured *Myxidium anguillae*, see page 114. The number of cysts reaches probably "several hundreds."

Vegetative form: Cysts, beneath the epidermis, usually subcircular, more or less irregularly triangular or quadrilateral under the magnifier, with the largest diameter of from 142 to 267 μ . The epidermis is slightly lifted up by the cyst. No chromatophore on the surface of the cyst. The cysts separated from each other, are found mostly in the central region of the body, head and fins being free from cysts. In cross-section, cysts exhibit oval or lenticular shape with the longest diameter, which is twice as long as the depth, placed parallel to the surface of the skin. No particular pathological change was noticed.

Spore: Circular in front view; broad fusiform or lenticular in side view. Sutural ridge fairly distinct. Sutural edge comparatively broad, especially

at the posterior margin, where a few folds (three are figured) are seen. Two oval polar capsules convergent and of equal size. Sporoplasm is sharply contoured, no iodophilous vacuole being recognized. Dimensions in preserved material (?): diameter 6.3 to 7 μ , thickness 4.2 to 4.9 μ , length of polar capsule 2.8 to 3.5 μ .

Family MYXOBOLIDAE Thélohan

1892	<i>Myxobolidées</i>	Thélohan	1892 : 173, 176
1893	<i>Myxobolidae</i>	Gurley	1893 : 412, 413
1895	<i>Myxobolidées</i>	Thélohan	1895 : 347

The characters of the family are described on page 58.

Genus MYXOBOLUS Bütschli

1882	<i>Myxobolus</i>	Bütschli	1882 : Pl. 38 : 6-10
------	------------------	----------	----------------------

The characters of the genus are described on page 58.

Type species: *Myxobolus mülleri* Bütschli.

MYXOBOLUS MÜLLERI Bütschli

[Figs. 397 to 403]

1881		Bütschli	1881 : 630
1882	<i>Myxobolus mülleri</i>	Bütschli	1882 : 595
1895	<i>Myxobolus mülleri</i>	Thélohan	1895 : 349
1905	<i>Myxobolus mülleri</i>	Nufer	1905' : 77, 79, 186
1906	<i>Myxobolus mülleri</i>	Cépède	1906 : 64-65
1906	<i>Myxobolus mülleri</i>	Schröder	1906 : 195
1908	<i>Myxobolus mülleri</i>	Auerbach	1908 : 456
1909	<i>Myxobolus mülleri</i>	Auerbach	1909a : 54, 71
1910	<i>Myxobolus mülleri</i>	Wegener	1910 : 81
1911	<i>Myxobolus mülleri</i>	Nemeczek	1911 : 160
1912	<i>Myxobolus mülleri</i>	Parisi	1912 : 293

Habitat: Air bladder and branchiae of *Leuciscus cephalus* L.; kidney and ovary of *L. phoxinus* L.; eye of *Crenilabrus melops* L. and *Alburnus lucidus*; branchiae of *Aspro asper* L., *Barbus vulgaris* Flem., *Leuciscus rutilus* L., *Squalius cephalus* L., *S. agassizii* Heckel, *Lota vulgaris* L., *Phoxinus loevis* Ag.; pseudobranchiae of *Cottus gobio* L.; intestine of *Mugil auratus* Risso.; France, Germany [Karlsruhe, Alle (October), Pregel, Frisches Haff], Switzerland (Neuchatel Lake), Italy (Napoli, September).

Vegetative form: White cysts in the connective tissue. Form elongated oval, 2 to 3mm. in diameter. No clear differentiation of protoplasm is observed even in young forms. In sections, some cysts show radiate striations in the thick granule-free ectoplasm. Endoplasm filled with nuclei.

Cépède writes as follows: Cysts in branchiae, subspherical or elliptical, 1.5mm. by 0.5mm.

Wegener's form: Cysts small and rounded, 0.2 to 0.3mm. in diameter.

Spore: Ordinarily spherical or subspherical. Two polar capsules

with a small triangular intercapsular appendix. Polar capsules pyriform and of same size. Sutural edge exhibits folds (7 to 9).

Thélohan's dimensions: length 10 to 12 μ , breadth 9 to 11 μ , length of polar capsule 5 μ .

Cépède gave the following dimensions in vivo: length 10 μ , breadth 9 μ , thickness 6 μ , length of polar capsule 5 μ .

Wegener's form. Usually oval, often almost spherical. Length 10 to 11 μ , breadth 8 to 9 μ , diameter of spherical form 9 μ , polar capsule 4 to 5 μ by 2 to 3 μ .

MYXOBOLUS PIRIFORMIS Thélohan

[Figs. 363 to 364]

1852		Remak	1852 : 144
1883		Balbiani	1883 : 197-198
1884		Balbiani	1884 : 125
1891		Pfeiffer	1891 : 132
1892	<i>Myxobolus piriformis</i>	Thélohan	1892 : 177
1893	<i>Myxobolus piriformis</i>	Gurley	1893 : 414
1894	<i>Myxobolus piriformis</i>	Gurley	1894 : 211
1895	<i>Myxobolus piriformis</i>	Thélohan	1895 : 348
1905	<i>Myxobolus piriformis</i>	Nufer	1905 : 77, 186
1910	<i>Myxobolus piriformis</i>	Plehn	1910 : 22-27
1910	<i>Myxobolus piriformis</i>	Wegener	1910 : 73

Habitat: Branchiae, spleen, kidney of *Tinca tinca* L., *Cobitis fossilis* L. and subcutaneous connective tissue, spleen, liver, connective tissue of the intestine of *Leuciscus* sp.; France, Germany (Pregel), Switzerland.

Vegetative form: Small, long thread-like cysts. Color white. Polysporous.

Wegener's form: average size, length 1mm., breadth 0.09 to 0.1mm

Spore: Elongated oval; flattened. Anterior end highly attenuated and slightly bent to one side. One pyriform polar capsule at this end. Dimensions: length 16 to 18 μ , breadth 7 to 8 μ , length of polar filament 30 μ .

Wegener gives the following dimensions: length 18 μ , breadth 7.5 μ , polar capsule 7.5 μ by 3.5 μ .

MYXOBOLUS UNICAPSULATUS Gurley

[Figs. 365 to 366]

1841		Müller	1841 : 487
1893	<i>Myxobolus unicapsulatus</i>	Gurley	1893 : 414
1894	<i>Myxobolus unicapsulatus</i>	Gurley	1894 : 210-211

Habitat: In the skin of *Labeo niloticus* For.; Nile.

Vegetative form: Cysts very small pustules in the skin of the head.

Spore: Form similar to *Myxosoma dujardini*. A single polar capsule at the anterior end, obliquely directed. Dimensions: length 0.0051''', breadth 0.0034'''.

MYXOBOLUS FUHRMANNI Auerbach

[Fig. 367]

1909	<i>Myxobolus fuhrmanni</i>	Auerbach	1909 : 65-68
1910	<i>Myxobolus fuhrmanni</i>	Auerbach	1910c : 178-179

Habitat: Connective tissue under the mucous membrane of the mouth of *Leuciscus rutilus* L.; Neuchatel Lake.

Vegetative form: Cysts, of pea-size, surrounded by several membranous layers of connective tissue with a few nuclei. Finely granular ectoplasm forms outer layer. Endoplasm is dense and contains faintly stained nuclei. Pansporoblasts and spores are found in the central portion of the cyst. Polysporous.

Spore: Elongated pyriform, with attenuated anterior and rounded posterior ends. Majority with a single polar capsule; spores with two polar capsules were also observed. Shell thick, at the posterior end. 4 to 6 notch-like markings on the posterior part of the shell. Sutural ridge thickened and fairly well marked. Coiled polar filament visible in preserved material. The opening of the polar capsule is either at the anterior end or near it. Sporoplasm with two nuclei of unequal size and a comparatively large iodophilous vacuole, stained brown with iodine alcohol. Dimensions: length 18 to 20 μ , breadth about 8 μ , thickness 6 μ , length of polar capsule 9 to 10 μ .

MYXOBOLUS OCULI-LEUCISCI Trojan

[Fig. 368]

1909	<i>Myxobolus oculi-leucisci</i>	Trojan	1909 : 679-682
------	---------------------------------	--------	----------------

Habitat: Vitreous body of the eye of *Leuciscus rutilus* L.; Prague (May?).

Vegetative form: Two cysts, spherical and subspherical, 100 to 180 μ in diameter. Ectoplasm finely granular. Outer portion of endoplasm with small nuclei, then larger nuclei each surrounded by protoplasm, while the central portion contains spores. Polysporous.

Spore: Elongated oval, flattened dorso-ventrally. Posterior margin rounded. At the anterior end, a single polar capsule with distinctly visible coiled polar filament. Shell smooth without any markings. Sporoplasm with one nucleus, usually elongated oval (2.8 μ in diameter) and one vacuole, occupies more than half of the space of the spore. Dimensions: length 9 to 10 μ , breadth 4.5 to 5.5 μ , thickness 3 μ , polar capsule 5 μ by 2 μ .

MYXOBOLUS TOYAMAI Kudo

[Figs. 369 to 370]

1915	<i>Myxobolus toyamai</i>	Kudo	1915 : 517-523
1917	<i>Myxobolus toyamai</i>	Kudo	1917 : 163-170

Habitat:—Connective tissue of branchial lamellae of *Cyprinus carpio* L.; Tokio (July).

Vegetative form: Cysts, ovoidal or in shape of calabash. Small form 67 by 50 μ , shows clear differentiation of protoplasm. Ectoplasm radially striated, often, differentiates fine processes (2 to 3 μ long). Endoplasm coarsely granular, contains nuclei from 1 to 4 μ in diameter. Size up to 190 μ in greatest diameter in sections. Two spores are formed in each pansporoblast. Polysporous.

Spore: Pyriform, with attenuated anterior and rounded posterior ends. No bilateral symmetry. Lateral sides are curved. Calabash shaped spores often occur. Shell without any marking, thickened at the anterior end. Sutural ridge shows sometimes a short (1.5 μ long) tail-like process at the posterior tip. A single pyriform polar capsule at the anterior end; in stained preparations, a small, oblong mass of protoplasm is seen between the polar capsule and the shell. Coiled polar filament distinct. Sporoplasm with two nuclei of usually same size and a relatively large iodophilous vacuole, 3 μ in diameter. Dimensions: length 15 μ , breadth 7 to 8 μ , thickness 5 to 6 μ , polar capsule 7 to 8 μ by 3 to 4 μ , length of polar filament 40 to 45 μ (pressure, perhydrol, KOH).

MYXOBOLUS NOTATUS Mavor

[Figs. 371 to 372]

1916	<i>Myxobolus notatus</i>	Mavor	1916a : 70-71
------	--------------------------	-------	---------------

Habitat: Connective tissue of the voluntary muscles on the sides or tail of *Pimephales notatus* Raf.; Georgian Bay, Canada (Summer).

Vegetative form: Cysts as large as 3mm. in diameter, are surrounded by a layer of columnar epithelial cells (origin and significance?) and a dense layer of connective tissue. Protoplasm is not clearly differentiated, tho the cyst is surrounded by an area devoid of nuclei. In the outer region of endoplasm, numerous nuclei each with a caryosome, are recognized. In the course of spore formation, two nuclei for polar capsules appear at first, one of which degenerates later. Polysporous.

Spore: Pyriform, with a posterior extension forming a process, 5 μ in length and as broad as the spore. A single polar capsule at the anterior end. An iodophilous vacuole in the sporoplasm. Dimensions: length 17 to 18 μ , breadth 7.5 to 8 μ , polar capsule 7 μ by 4 μ , length of polar filament 95 μ .

MYXOBOLUS sp. Kudo

1918 *Myxobolus* sp. Kudo 1918 : 15

Habitat: Spleen of *Perca flavescens*; West Falmouth, Mass. (August). Isolated spores were noticed in one fish, in smears and section preparations.

Vegetative form: Not observed.

Spore: Ovoidal, attenuated at the anterior end. Shell uniformly thick. A single polar capsule opens at the anterior tip. Sporoplasm contains an iodophilous vacuole and two nuclei of equal size (2μ). Dimensions: length 18 to 20μ , breadth 8μ , polar capsule 7 to 9μ by 3 to 6μ .

MYXOBOLUS ROHITAE Southwell et Prashad

[Figs. 373 to 374]

1918 *Myxobolus rohita* Southwell et Prashad 1918 : 344-347

Habitat: Branchiae of *Labeo rohita*; Turag river, Mirpur, Dacca district, Bengal (June). Type specimens of Indian Museum P48/1. Infection was heavy. In one case 53 cysts were found on one surface of a single gill.

Vegetative form: Cysts in the gill-filaments. The cysts preserved in alcohol are of a creamy-yellow color, oval to cylindrical in form, lying with the long axis parallel to the gill-filaments. Cysts attached to the gill-filaments with the flattened surface. Size: length 3.1 to 3.8mm.; breadth 0.8 to 1.2mm. Cyst-wall striated vertically, covered with an epithelium, two to three layers thick. In the central portion and at the periphery, mature spores and pansporoblasts as well as immature spores were found respectively. Polysporous.

Spore: Elongated pyriform, rounded at the posterior end and acutely pointed anteriorly. Sutural ridge slightly raised. One polar capsule present, being of conspicuous size. Coiled polar filament is distinctly observed in the polar capsule. An iodophilous vacuole, 3.6μ in diameter, in the sporoplasm. "Lying just posterior to it is the nucleus of the spore. A few granules of chromatin were also seen lying scattered in the protoplasm." Dimensions: length 30 to 32μ , breadth 7 to 8μ , length of the polar capsule 22 to 23μ , that of polar filament 92 to 97μ .

MYXOBOLUS SENI Southwell et Prashad

[Figs. 375 to 376]

1918 *Myxobolus seni* Southwell et Prashad 1918 : 347

Habitat: On the median and caudal fins of *Labeo rohita*; Mirpur, Dacca (January). Type specimens in Indian Museum numbered P 53/1.

Vegetative form: Trophozoites form cysts which are elongated ellipsoidal. Size from 4.7mm. to 5.4mm. in length, 2.9mm. to 3.7mm. in

breadth. Color of the cyst whitish with black scattered granules on the surface.

Spore: Oval, much wider behind than in front and pointed at the anterior end. Sutural ridge is slightly thickened. A single polar capsule, showing much coiled polar filament. Iodinophilous vacuole 2.3μ in diameter. Dimensions: length 13.2 to 13.6μ , breadth 10.1 to 10.3μ , length of polar capsule 4μ , length of polar filament 43μ (in one case).

MYXOBOLUS MISGURNI nov. spec.

[Figs. 377 to 378]

1916 *Myxobolus fuhrmanni* Kudo 1916 : 5

Habitat: Gall-bladder of *Misgurnus anguillicaudatus*; Tokio (September). About 50% of the fish examined showed a few isolated spores floating in the bile.

Vegetative form: Unobserved.

Spore: Form elongated pyriform, with attenuated anterior and rounded posterior ends. Shell uniformly thick. Over sutural edge, shell exhibits many (up to 12) triangular markings. Sutural ridge distinct. A single pyriform polar capsule at the anterior end. Sporoplasm contains an iodophilous vacuole and two nuclei. Coiled polar filament distinct in vivo. Dimensions of fresh spores: length 14 to 15.5μ , breadth 6 to 7.3μ , thickness 5 to 6μ , polar capsule 6.3μ by 2 to 3μ , length of polar filament up to 100μ .

Remarks: The writer reported this species as identical with *Myxobolus fuhrmanni* Auerbach. By repeated reexamination and comparison with Auerbach's description, however, he came to the conclusion that the present form should be treated as a new species, on account of the difference of the host and the characters of the spore.

MYXOBOLUS PFEIFFERI Thélohan

[Figs. 379 to 385]

1890	<i>Myxosporidian</i>	Pfeiffer	1890 : 30-37
1891	<i>Myxosporidian</i>	Pfeiffer	1891 : 100, 105-110, 130
1893	<i>Myxosporidian</i>	Pfeiffer	1893 : 118-130
1895	<i>Myxobolus pfeifferi</i>	Thélohan	1895 : 350
1898	<i>Myxobolus pfeifferi</i>	Doflein	1898 : 306, 320, etc.
1906	<i>Myxobolus pfeifferi</i>	Cépède	1906 : 59
1906	<i>Myxobolus pfeifferi</i>	Stazzi	1906 : 14-19
1906	<i>Myxobolus pfeifferi</i>	Mercier	1906 : 427-428; 1906a : 763-764
1908	<i>Myxobolus pfeifferi</i>	Keysselitz	1908 : 253-273, 286-306
1909	<i>Myxobolus pfeifferi</i>	Mercier	1909 : 5-30

Habitat: Muscle and connective tissue of kidney, spleen, intestine, ovary, etc., of *Barbus barbus* L., and branchiae of *B. fluviatilis* Ag. and *B. plebejus* Val.; Drac (June), Neckar, Prag, Milano. The cause of well known "Boil disease" (Beulenkrankheit) or Myxoboliasis tuberosa (Hofer) of the barbels in European waters. Among many observers Keysseltiz made a thoro study of the parasite. His observations are as follows: The disease occurs among the fish at any stage of growth. About 8% of the fish, 7 to 15cm. long, caught in May and June between Conz and Trier were infected with the parasites. The heaviest infection, however, occurs among fish up to 40cm. in length; fish 50cm. long or larger show the tumors caused by the parasites, rather rarely. Most of the fish die as the result of the infection between the early part of April and the end of October. The highest mortality is reached in the hottest months, i.e., July and August. The temperature greatly affects the growth of the parasites. Fish kept in the aquarium at a temperature of 25° C. or higher demonstrate the growth of the boil in size daily. The boils are not noticed during the winter and spring, they are formed from the early part of April to the middle of October.

Vegetative form: The parasites develop tumors of conspicuous size.

Keysseltiz's observations are as follows: The tumor varies in size from millet-grains to hen's eggs. Form spherical, oval or elongated. The number of cysts on a single fish, is usually 3 to 4; often one, in some fish, however, 23 were recognized on one fish. Usually tumors separated from each other, rarely many forming one tumor. In one fish, 27cm. in length, a tumor of 7cm. long, 4cm. broad and 3cm. thick, was observed in July. The seat of infection is: the muscle of the body, muscle of pectoral and anal fins, often in peritoneum and rarely in intestine. As the result of breaking up of the cyst membrane, spores are also found in the testis, liver and kidney.

The tumor is composed of many vegetative forms, rounded, oval, elongated, variously branched or flattened. Size reaches to 1.5mm. in diameter. Protoplasm is usually differentiated into ectoplasm and endoplasm. The surface is not often smooth, but shows irregular outline. Ectoplasm is seen often as a very thin, uniformly hyaline, indistinctly granular or radially striated layer, giving the network-like appearance to the surface of the body. Endoplasm, stained more deeply around the peripheral part than other portion, shows a coarsely alveolar structure in the central region. It contains vegetative nuclei, developmental stages of propagative nuclei, granules, fat-like, often leucocytes and red blood corpuscles. The leucocytes, uninuclear or multinuclear, were seen at the periphery, apparently in the course of degeneration. Red blood corpuscles were found, in section, inside of the apparently intact parasite. Each pansporoblast develops into two spores. Polysporous.

Cépède observed one cyst, about 2mm. in diameter, in the connective tissue of the third gill arch.

Spore: Thélohan described as follows: Ovoidal. Sutural edge shows folds. A small triangular intercapsular appendix. Dimensions: length 12 μ , breadth 10 μ . Cépède's form showed exactly the same dimensions.

Keysselitz gave the characters of the spore as follows:

Flattened oval. Shell smooth. A small intercapsular appendix. Sutural edge having a number of small flat enlargement, size and number being variable. Two convergent narrow canals (foramina) penetrate the shell at the anterior end. Two polar capsules, pyriform and of equal or nearly equal size, are located at the anterior half. Coiled polar filament distinct, coiled 7 to 8 times. No distinct connection between polar capsule and the filament. Sporoplasm fills the posterior half of the spore, extending into intercapsular cavity. It is finely reticular, exhibits one or two rounded or oval vesicular nuclei and an iodophilous vacuole. Fat-like substance is often seen around the polar capsules. Spores kept in water for four months remain intact in large numbers. Dimensions: length 12 to 12.5 μ , breadth 10 to 10.5 μ , length of polar capsule, 5.5 to 6 μ , length of polar filament 28 to 34 μ .

MYXOBOLUS INAEQUALIS Gurley

[Fig. 411]

1841		Müller	1841 : 487-488
1893	<i>Myxobolus inaequalis</i>	Gurley	1893 : 414
1894	<i>Myxobolus inaequalis</i>	Gurley	1894 : 212

Habitat: In the skin of the head of *Piramutana blochi* Cuv. et Vil. and *Synodontis schall* Bl. Schn.; Guiana, Surinam.

Vegetative form: Very small pustules in the skin of the head.

Spore: Ovoidal. Two polar capsules of unequal size at the anterior end. Dimensions: length 0.0052''', breadth 0.0033'''.

MYXOBOLUS DISPAR Thélohan

[Fig. 386]

1895	<i>Myxobolus dispar</i>	Thélohan	1895 : 348
1904	<i>Myxobolus dispar</i>	Hofer	1904 : 50
1910	<i>Myxobolus dispar</i>	Wegener	1910 : 73-74
1911	<i>Myxobolus dispar</i>	Nemeczek	1911 : 145

Habitat: Branchiae of *Carassius carassius* L., branchiae and epithelium of intestine of *Cyprinus carpio* L., also muscle and spleen of *Scardinius erythrophthalmus* L. and in the skin and the connective tissue of *Alburnus lucidus* Heck.; France, Austria, Königsburg (March, July, September).

Vegetative form: Not described by Thélohan.

Wegener's description is as follows: Cysts: white in color; spindle shape with pointed ends. Cysts in *Carassius carassius* L. smaller and oval.

Size 3.5mm. by 0.8mm. Cysts are surrounded by thick layers (7 to 8 μ) of the connective tissue of the host. Ectoplasm seems to be undifferentiated. Endoplasm granular, contains a larger number of spores. Polysporous.

Spore: Thélohan's diagnosis is as follows:

Ellipsoidal or slightly oval. Shell with 3 to 5 folds along sutural edge. Polar capsules of unequal size, with a small intercapsular body. The vacuole is difficult to stain with iodine. Dimensions: length 10 to 12 μ , breadth 8 μ , polar capsule 7 μ by 5 μ .

Wegener's form is as follows: length 11 to 12 μ , breadth 7.5 to 8 μ , larger polar capsule 6 to 7 μ by 3.5 μ , smaller one 4 μ by 2.5 to 3 μ . The sporoplasm is shifted toward the smaller polar capsule.

MYXOBOLUS ELLIPSOIDES Thélohan

[Figs. 387 to 389]

1852		Remak	1852 : 144-146
1892	<i>Myxobolus ellipsoides</i>	Thélohan	1892 : 177
1895	<i>Myxobolus ellipsoides</i>	Thélohan	1895 : 350-351
1898	<i>Myxobolus ellipsoides</i>	Doflein	1898 : 324, etc.
1905	<i>Myxobolus ellipsoides</i>	Nufer	1905 : 77, 79, 186.
1910	<i>Myxobolus ellipsoides</i>	Wegener	1910 : 74-75
1912	<i>Myxobolus ellipsoides</i>	Lo Giudice	1912 : 1-79

Habitat: Connective tissue of air bladder, branchiae, kidney, spleen, liver and cornea of *Tinca tinca* L., branchiae of *Abramis brama* L., *Alburnus lucidus* Heck., *Leuciscus rutilus* L., *Squalius cephalus* L., *Abramis vimpa* Cuv., *Blicca björkna* L., *Idus melanotus*; France, Vierwaldstätter See, Prague, Masurische See, Italy.

Vegetative form: Thélohan does not describe.

According to Wegener, white cysts, elongated oval; 2mm. by 0.5mm. in size. Polysporous.

Spore: Thélohan described as follows: Flattened elliptical, rather elongated. Sutural edge broad without any folds. Shell with no marking. Form of the spore somewhat variable. Two polar capsules of equal size, capsulogeneous nuclei present even when fully grown. Abnormal spores are of frequent occurrence. Dimensions: length 12 to 14 μ , breadth 9 to 11 μ , length of polar capsule 4 μ .

Wegener's form: length 14 to 15 μ , breadth 10 to 11 μ , polar capsule 4 to 5 μ by 3 μ . Shell comparatively thick. One spore had a tail 5 μ long.

MYXOBOLUS EXIGUUS Thélohan

[Figs. 390 to 395]

1891	<i>Myxosporidium mugilis</i> ?	Perugia	1891 : 23
1895	<i>Myxobolus exiguus</i>	Thélohan	1895 : 349-350
1906	<i>Myxobolus exiguus</i>	Schröder	1906 : 195
1910	<i>Myxobolus exiguus</i>	Wegener	1910 : 75
1912	<i>Myxobolus exiguus</i>	Parisi	1912 : 294-295

Habitat: Branchiae of *Abramis brama* L. and *Chondrostoma nasus* L., wall of stomach, pyloric coecum and intestine, branchiae, spleen, kidney of *Mugil chelo* Cuv., *M. capito* Cuv. and *M. auratus* Riss.; Le Vivier-sur-mer, Banyuls, Marseille, Heidelberg, Pregel, Frisches Haff, Kurisches Haff, Geneva, Napoli.

Vegetative form: No description by Thélohan.

Wegener writes as follows:

Cysts of variable size. Color white. Usually small and narrow, 0.5 to 0.7mm. long and 0.2mm. wide. Frequently large round cysts of 1.2 to 1.5mm. in diameter, filling the lamella. Cysts are surrounded by 10 to 11 μ thick membrane composed of the connective tissue of the host. Ectoplasm, 5 μ thick, is faintly stained by hematoxylin. Outer region of endoplasm, alveolar and densely loaded with nuclei, while in the central portion with mature spores in granular ground-mass.

Parisi's observations are as follows:

Cysts in the intestinal wall of *Mugil auratus*, large; reaching a length of 3mm.

Spore: Thélohan's description is as follows:

Flattened ovoidal, with more or less attenuated anterior end. Sutural edge shows fairly noticeable folds. A small triangular intercapsular appendix. Vacuole in the sporoplasm is usually hard to stain with iodine. Dimensions: length 8 to 9 μ , breadth 6 to 7 μ , length of polar filament 15 μ (KOH).

Wegener observed as follows: Rounded with slightly pointed anterior end. Length 8 to 9.5 μ , breadth 6 to 7.5 μ , polar capsule 4.5 μ by 2 to 3 μ . Shell exhibits small folds around the sporoplasm. An intercapsular triangular body indistinctly visible.

Parisi gave the following dimensions: length 8 to 8.5 μ , breadth 6 to 7 μ , thickness 5.5 μ , polar capsule 3 to 4 μ by 1.5 to 2 μ , length of polar filament 30 μ (alkaline). Folds usually 6 in number. Coiled polar filament visible in vivo.

MYXOBOLUS OVIFORMIS Thélohan

[Fig. 396]

1854		Lieberkühn	1854 : 21-22
1892	<i>Myxobolus oviformis</i>	Thélohan	1892 : 177
1895	<i>Myxobolus oviformis</i>	Thélohan	1895 : 351
1905	<i>Myxobolus oviformis</i>	Nufer	1905 : 77, 186
1906	<i>Myxobolus oviformis</i>	Cépède	1906 : 60
1910	<i>Myxobolus oviformis</i>	Wegener	1910 : 76-78

Habitat: Fin (subcutaneous tissue), spleen, kidney and liver of *Gobio gobio* L.; branchiae of *Alburnus lucidus* Heck., *Cyprinus carpio* L., *Blicca björkna* L., *Abramis brama* L. and *A. vimba* L.; France (Isère), Frisches Haff (especially spring months), Switzerland.

Vegetative form: Thélohan gave no description.

Wegener's observations are as follows:

Cysts, white, 0.75 to 1.7mm. by 0.4 to 0.7mm. In sections, cysts are shown to be surrounded by a thick (10 to 20 μ , average 16 μ) layer of connective tissue. Ectoplasm a thin (6 to 8 μ thick) layer, exhibits a transverse striation. The striation is often absent at places in ripe cysts. Endoplasm finely granular. In young cysts, it is, however, reticulated, with nuclei of 1.5 μ in diameter.

Spore: Thélohan described as follows: Flattened ovoidal with pointed anterior end. Shell smooth. No folds. Polar capsule comparatively large. Dimensions: length 10 to 12 μ , breadth 9 μ , polar capsule 6 μ .

Cépède observed numerous spores in the liver and kidney of *Gobio gobio*. Dimensions in vivo: length 10 to 12 μ , breadth 9 μ , length of polar capsule 6 μ . Polar capsules of equal size. Coiled polar filament distinct.

Wegener's form: length 10.5 to 11 μ , breadth 7.5 to 8 μ , polar capsule 5 to 6 μ by 3 μ .

Remarks: Wegener recognized another form, which seems to be of very rare occurrence and which can not be distinguished distinctly from the above described form. Cysts at the end of the branchial lamellae. Size 1.7 to 2mm. in largest length. Spore resembles more closely the figure given by Thélohan for *Myxobolus oviformis* than the above mentioned form which he observed. A small intercapsular appendix (rounded) indistinct. Sporoplasm comparatively small. Length 12.5 to 13.5 μ , breadth 9 μ , polar capsule 7.5 μ by 3 μ .

MYXOBOLUS LINTONI Gurley

[Figs. 404 to 408]

1891		Linton	1891 : 99-102
1893	<i>Myxobolus lintoni</i>	Gurley	1893 : 414
1894	<i>Myxobolus lintoni</i>	Gurley	1894 : 238

Habitat: Superficial musculature and subcutaneous tissue of *Cyprinodon variegatus*; Woods Hole (August).

Vegetative form: Cysts, not closed, but fungoid masses of an irregular shape, varying in size from 4mm. by 2.5mm. to 10mm. by 4mm., projecting as much as 3mm. above general surface of skin. The skin of the host overlying these tumors, is more or less cracked and broken, the scales being scattered.

Spore: Elliptical in the front view; lenticular in side view. Shell thick. Sutural ridge marked. Two polar capsules, convergent, at the anterior end. Spores kept in alcohol, extruded polar filaments under the action of iodine water and sulphuric acid. Sporoplasm with a large iodophilous vacuole. Dimensions: length 13.9 μ , breadth 11 μ , thickness 8 μ .

MYXOBOLUS GLOBOSUS Gurley

[Figs. 409 and 410]

1893	<i>Myxobolus globosus</i>	Gurley	1893 : 415
1894	<i>Myxobolus globosus</i>	Gurley	1894 : 241

Habitat: Branchial lamellae of *Erimyzon sucetta oblongus* Lac. (*Catostomus tuberculatus* Le Sueur); Kinston (N.C.), Columbia, (S.C., March), tributaries of Fox River.

Vegetative form: Cysts, whitish, elongated elliptical or rod-shaped, surrounded by very thin membrane? Size up to 0.5mm. in max. length. Polysporous.

Spore: Globose, subcircular in outline. Shell thin and very transparent. Sutural ridge very wide, being one third of the thickness of the spore. Polar capsules two, of equal size, divergent. Vacuole present, but not clearly contoured. Dimensions: length 7 to 8 μ , breadth 6 to 7 μ , thickness 5 μ .

MYXOBOLUS OBLONGUS Gurley

[Figs. 412 to 416]

1841		Müller	1841 : 487-490
1893	<i>Myxobolus oblongus</i>	Gurley	1893 : 414
1894	<i>Myxobolus oblongus</i>	Gurley	1894 : 234-238

Habitat: Beneath the skin, chiefly of the head of *Erimyzon sucetta oblongus* Lac. (*Catostomus tuberculatus* Le Sueur); Kinston, tributaries of Fox River.

Vegetative form: Cysts, round or elliptic, not over 1mm. in diameter, covered by resistant membrane. Color whitish. Polysporous.

Spore: Spatular, approaching roundish-oblong. Shell thin and transparent. Sutural ridge wide. Two polar capsules, pyriform, of equal size. Sporoplasm extending forward along the upper surface. Vacuole could not be detected. Dimensions: length 14 to 17 μ , breadth 8.5 μ , thickness 5 to 6 μ .

MYXOBOLUS TRANSOVALIS Gurley

[Figs. 417 and 418]

1893	<i>Myxobolus transovalis</i>	Gurley	1893 : 415
1894	<i>Myxobolus transovalis</i>	Gurley	1894 : 242

Habitat: Under scales on external surface of *Phoxinus (Clinostomus) funduloides* Girard; 4 Mile Run, Carlisle, Va., tributary of Potomac River (June). No fish of the same species caught from the same locality on August 29 of the same year was found infected.

Vegetative form: It is not certain whether cysts exist or not. Spores in mass, appear to be held together by a small gelatinous or mucoid mass

which has no attachment to the subjacent connective tissue. It forms a thin discoidal mass situated in the center of the concave surface of the scale. The color of the mass slightly more yellowish than the surrounding tissue, when coagulated. It is exceedingly difficult to detect its presence in the fresh state.

Spore: Elliptical, with the largest diameter passing thru two polar capsules. Shell thin. Sutural edge narrow. Two polar capsules of equal size convergent. Polar filament is extruded under the action of glycerine and sulphuric acid. The vacuole in the sporoplasm is difficult to detect. Sporoplasm also contains two nuclei, rarely one, 1 to 1.5μ in diameter. Dimensions: length 6 to 7μ , breadth 8μ .

MYXOBOLUS OBESUS Gurley

[Figs. 419 and 420]

1883		Balbiani	1883 : 203
1893	<i>Myxobolus obesus</i>	Gurley	1893 : 415
1894	<i>Myxobolus ? obesus</i>	Gurley	1894 : 239
1899	<i>Myxobolus obesus</i>	Labbé	1899 : 100
1906	<i>Myxobolus obesus</i>	Cépède	1906 : 60-61

Habitat: On *Alburnus alburnus* L.; branchiae and kidney of *A. lucidus* Heck. (*A. mirandella* Bl.); Lac du Bourget.

Vegetative form: Balbiani gave no observation.

Cépède observed as follows: Cysts, ovoidal, more or less elongated or variable in form, not exceeding 800μ in length. In kidney, numerous cysts were of subspherical, ovoidal or rarely irregularly elongated form. Sub-spherical cysts 500 to 600μ in average diameter. Polysporous.

Spore: Cépède describes as follows: Subcircular or ovoidal in front view; lenticular in side view. Sutural edge exhibits variable numbers (4 to 5) of fold-like markings on the shell. Polar capsules pyriform and of equal size. Coiled polar filament distinct. A small triangular intercapsular appendix. Sporoplasm with a subspherical and clearly outlined vacuole and two nuclei. Dimensions in vivo: length 11.5 to 12μ , breadth 7.5 to 8μ , thickness 5μ . Those of fixed and stained spores: length 11.25 to 11.50μ , breadth 7.25 to 7.50μ , length of polar capsule 5μ .

Remarks: Cépède mentions that *Alburnus alburnus* L. mentioned by Gurley is "without doubt" identical with *A. lucidus* Heckel.

MYXOBOLUS CYCLOIDES Gurley

[Fig. 421]

1841		Müller	1841 : 481, 486
1893	<i>Myxobolus cycloides</i>	Gurley	1893 : 415
1894	<i>Myxobolus cycloides</i>	Gurley	1894 : 239
1906	<i>Myxobolus cycloides</i>	Cépède	1906 : 61-63
1910	<i>Myxobolus cycloides</i>	Wegener	1910 : 79-80

Habitat: Opercle, pseudobranchiae and kidney of *Leuciscus rutilus*; branchiae of *Scardinius erythrophthalmus*, *Blicca björkna* L., *Gobio gobio* L., *Abramis vimba* L., *A. brama* L., *Rhodeus amarus* Bl., *Alburnus alburnus* L., *Lota lota* L.; France (Isère), Germany (Pregel, Frisches and Kurisches Haff, Masurische See, January, May).

Vegetative form: Wegener observed cysts as follows. A type: 1 to 2mm. by 0.4 to 0.7mm. Form exactly like that of *Myxobolus oviformis*. B type: small and round, present in groups. C type: small 0.5mm. by 0.2mm.

Spore: Gurley gave the following short diagnosis from the observations of J. Müller: subcircular-ovate or broadly rounded elliptic, length 12μ .

Cépède distinguishes three different types of spores as follows: Lenticular in side view; subcircular (13.5μ by 13μ), oval (14.7μ by 11.4μ) and ovoidal (16μ by 11μ) in front view. Two polar capsules of equal size (6μ by 4μ), closely set or separated (3μ apart) from each other. Coiled polar filament distinct. A small triangular intercapsular appendix. Sporoplasm refractive and finely granular. Sutural edge exhibits folds of variable number at the posterior portion. Dimensions of fixed and mounted spores: length 10.5 to 12μ , breadth 7.5 to 8μ .

Wegener, without noticing Cépède's paper, also mentions three different types chiefly distinguished by the spore as follows:

A type (common form), in the branchiae of *Lota lota*, *Abramis brama*, *A. vimba*, *Blicca björkna*, *Leuciscus rutilus*, *Alburnus alburnus* and *scardinius erythrophthalmus*. Cysts mentioned above.

Spore. Rounded or oval; flattened. A tail, 15μ long, was noticed twice. A triangular intercapsular appendix. Sutural edge usually having folds. Polar capsules often differ in form and size in different cysts, tho they are constant in one and the same cyst, causing the variability in size of sporoplasm. Dimensions: length 11 to 12.5μ , breadth 8 to 9μ , polar capsule 4.5 to 6μ by 3 to 3.7μ , in many cysts 7.5μ by 4μ .

B type. In the fifth gillarch of *Gobio gobio* L. Cysts mentioned above.

Spore. Elongated oval. A triangular intercapsular appendix. Indistinct folds on sutural edge. Dimensions: length 12.5 to 13.5μ , breadth 8 to 10μ , polar capsule 5 to 6μ by 3 to 4μ .

C type. In the branchiae of *Rhodeus amarus* Bl. and *Alburnus alburnus* L. (April and May). Cysts mentioned above.

Spore. Rounded. Distinct intercapsular appendix. Folds distinct on sutural edge. Dimensions: length 12 to 15μ , breadth 9 to 10μ , polar capsule 5 to 7μ by 3 to 4μ .

MYXOBOLUS SPHAERALIS Gurley

1874		Claparède	1874 : 113-114
1893	<i>Myxobolus sphaeralis</i>	Gurley	1893 : 415
1894	<i>Myxobolus sphaeralis</i>	Gurley	1894 : 240

Habitat: Mucosa of branchiae of *Coregonus lavaretus* L. (*C. fera*); Lake Geneva.

Vegetative form: Cysts, 0.25 to 0.33mm. in diameter. Polysporous.

Spore: Spherical, 9μ in diameter.

MYXOBOLUS ANURUS Cohn

[Figs. 422 and 423]

1895	<i>Myxobolus anurus</i>	Cohn	1895 : 42-43
1896	<i>Myxobolus anurus</i>	Cohn	1896 : 266
1899	<i>Henneguya psorospermica</i> <i>anura</i>	Labbé	1899 : 102
1910	<i>Myxobolus anurus</i>	Wegener	1910 : 76
1911	<i>Henneguya psorospermica</i> <i>anura</i>	Nemeczek	1911 : 146

Habitat: Branchiae of *Esox lucius* L.; Königsberg (March, December), Frisches Haff, Pregel, Masurische See, Lotzen, (September, October).

Vegetative form: Cysts small rounded and of white color. Cohn measures length 0.6mm., breadth 0.34mm. Wegener's form: length 0.3 to 0.5mm. and breadth 0.2 to 0.3mm.

Spore: Cohn's descriptions are as follows: More or less oval. Dimensions: length 12 to 15μ , breadth 4 to 6.8μ , polar capsule 5.5 to 7μ by 2.1 to 2.5μ , length of polar filament 32 to 38μ .

Wegener's form: Elongated and narrow, often with a tail. Dimensions: length 15μ (maximum up to 18μ), breadth 6 to 7μ , polar capsule 8μ by 3μ .

Remarks: Tho Labbé classified this as a subspecies of *Henneguya psorospermica* Thélohan, Wegener's observation gives stronger basis for placing this form in the genus *Myxobolus*.

MYXOBOLUS sp. Gurley

[Fig. 424]

1882		Bütschli	1882 : 590
1894	<i>Myxobolus</i> sp. incert.	Gurley	1894 : 214
1899	<i>Myxobolus</i> sp.	Labbé	1899 : 100

Habitat: *Nais lacustris* L. (*N. proboscidea*); Locality?

Vegetative form: Cysts, 8mm. by 4.25mm. Polysporous.

Spore: Oval or circular; tailed or untailed. These spores of different form occur, often, without order in the same cyst.

MYXOBOLUS sp. Gurley

[Fig. 425]

1894	<i>Myxobolus</i> sp. incert.	Gurley	1894 : 239
------	------------------------------	--------	------------

Habitat: Body cavity of *Carassius carassius* L.; Leipsic.

Vegetative form: Not observed.

Spore: Broadly elliptic; shell bivalve; valves equally convex. Sutural ridge. Two equal polar capsules. Sporoplasm with a vacuole. Dimensions: length 14μ , breadth 10μ , thickness 5μ .

Remarks: This species seems to be very similar to *M. carassii* Klokačewa (page 150).

MYXOBOLUS sp. Gurley

[Figs. 426 to 429]

1841		Müller	1841 : 480
1894	<i>Myxobolus</i> sp.	Gurley	1894 : 240-241
1899	<i>Myxobolus</i> sp.	Labbé	1899 : 100

Habitat: Skin of opercle, in the branchiae, on the head or on the fin of *Lucioperca lucioperca* L.; Germany, Don.

Vegetative form: Cysts 1.09 to 2.18mm. in diameter. Color whitish. Polysporous.

Spore: Rounded. Thickness equal to half the breadth. Sutural ridge. Two polar capsules, of equal size, converging.

MYXOBOLUS CYPRINI Doflein

[Figs. 430 to 432]

1896		Hofer	1896 : 2, 38-39
1898	<i>Myxobolus cyprini</i>	Doflein	1898 : 288, 320, 325
1904	<i>Myxobolus cyprini</i>	Hofer	1904 : 66-67
1909	<i>Myxobolus cyprini</i>	Doflein	1909 : 780-783
1916	<i>Myxobolus cyprini</i>	Doflein	1916 : 1026-1027

Habitat: Suppurative connective tissue and epithelium of kidney, liver and spleen of *Cyprinus carpio* L., rarely *Tinca vulgaris* Cuv. and *Abramis brama* L.; Germany, Austria. According to Hofer the parasites cause so-called "small pox of carp" among carp in German waters.

Vegetative form: Small ameboid. Form irregular. The youngest form with a single or many nuclei, is found in the epithelium of the kidney. Multiplication by multiple division, the nuclei undergoing amitotic division. Endoplasm contains homogeneous, yellow and refractive bodies. Also found in the state of diffuse infiltration. Spores are found in the parenchym of the kidney.

Spore: Oval. Shell thickened (1.5μ wide) along the sutural edge. Two converging polar capsules cross each other, in front view, at the anterior tip. Sporoplasm with an iodophilous vacuole. Dimensions: length 21μ , breadth 15μ , length of polar capsule 6μ . Doflein (1916:1027) gives the following dimensions: length 10 to 16μ , breadth 8 to 9μ .

Hofer gives the following dimensions: length 10 to 12μ (up to 16μ), breadth 8 to 11μ , polar capsule 5 to 6μ by 3μ , sutural edge 1.5μ .

MYXOBOLUS NEUROBIUS Schuberg et Schröder

[Figs. 433 to 436]

1905 *Myxobolus neurobius* Schuberg and Schröder 1905 : 49-56Habitat: Nervous tissue of *Trutta fario* L.; Gutach (May?).

Vegetative form: Cysts, usually elongated, often spherical. Elongated form 0.9mm. by 0.02mm. The seat of the cysts is between the medullary sheath and sheath of Schwann. Neither medullary sheath nor axis-cylinder was infected. Cyst-membrane could not be made out. Cysts contained only full-grown spores without any younger stage. Polysporous.

Spore: Broad oval in front view; spindle shaped in side view. Anterior end attenuated, posterior end rounded. Shell somewhat thick. Sutural ridge is not particularly marked. Edge without any fold. No intercapsular appendix. Sporoplasm, with a large and spherical iodophilous vacuole and a single nucleus, occupies less than one half of the inner space of the spore. Two polar capsules, pyriform, fuse into one at the anterior end. Coiled (8 to 10 times) polar filament distinct. Dimensions: length 10 to 12 μ , breadth 8 μ , thickness 6 μ , polar capsule 6 to 7 μ by 2 μ .

MYXOBOLUS AEGLEFINI Auerbach

[Figs. 437 to 441]

1906	<i>Myxobolus aeglefini</i>	Auerbach	1906 : 568-570
1906	<i>Myxobolus aeglefini</i>	Auerbach	1906a : 115-119
1907	<i>Myxobolus esmarkii</i>	Johnstone and Woodcock	1907 : 204-208
1909	<i>Myxobolus aeglefini</i>	Auerbach	1909 : 76-78
1910	<i>Myxobolus aeglefini</i>	Auerbach	1910c : 181-182
1911	<i>Myxobolus aeglefini</i>	Nemeczek	1911 : 162

Habitat: Cartilage and bone of cranium and eye of *Gadus aeglefinis*, *G. callarias*, *G. merlangus* L., *G. morrhua* L., *G. esmarkii* and *Molva vulgaris* Flem.; Norway, Morecambe (March).

Vegetative form: Cysts in cartilage and bone of cranium and in cartilaginous layer of the sclerotic of the eye. Protoplasm is distinctly differentiated. Ectoplasm somewhat vacuolated; endoplasm granular with numerous small nuclei. Polysporous.

Johnstone's observations are as follows: Round the peripheral part of the cornea, and covered loosely by conjunctiva are a number of milk-white rounded or oval bodies, from about 1 to 3mm. in diameter. Several of these fused to form elongated mass which lie along the curvature of the periphery of the eye. These cysts also invade the lateral and posterior parts of the bulbus oculi. In sections, the cysts lie within the thickness of cartilaginous layer of the sclerotic. This latter is enlarged into thick layer (2mm.) by the presence of the cysts.

Nemeczek mentions irregular cysts of 1.5mm. in diameter.

Spore: Elliptical in front view. Two polar capsules convergent. No intercapsular appendix. Sutural edge rather thick with a number of folds on the posterior margin. Sporoplasm with two nuclei and an iodophilous vacuole. Dimensions: length 10.8 to 11.7 μ , breadth 9.9 to 10.4 μ , thickness 7.2 to 9 μ , length of polar capsule 4.5 to 5 μ .

Woodcock's form has a spore with the following characters:

Slightly ovoid. Sporoplasm always contains a large and well defined vacuole and two nuclei. Dimensions: length 10 μ , breadth 8 μ , length of polar capsule 3.25 to 3.5 μ .

MYXOBOLUS GIGAS Auerbach

[Figs. 442 to 445]

1906	<i>Myxobolus gigas</i>	Auerbach	1906 : 386-391
1910	<i>Myxobolus gigas</i>	Auerbach	1910c : 182
1912	<i>Myxobolus gigas</i>	Parisi	1912 : 293-294

Habitat: Subcutaneous connective tissue of the operculum of *Abramis brama* L.; Karlsruhe, Pavia. Parisi observed cysts on the side, on the caudal fin (5 cysts on rays), on other fins, branchiae and in the internal organs of the fish.

Vegetative form: Cysts, spherical or ovoidal. No cyst membrane composed of the connective tissue of the host. Protoplasm is indistinctly differentiated. Ectoplasm thin and radially striated, which gradually turns into endoplasm. Endoplasm finely granular, contains numerous nuclei (2.5 to 2.7 μ in diameter). Size of greatest form 360 μ by 290 to 300 μ . According to Parisi size up to 1.5mm.

Spore: Elliptical when viewed from the front. Sutural edge somewhat narrow, having a number of folds at the posterior portion. Sporoplasm with an iodophilous vacuole and two nuclei. Dimensions: length 16.9 to 21.6 μ , breadth 13 to 16.2 μ thickness 9 μ , length of polar capsules 7.8 μ , length of polar filament 90 μ (sulphuric acid).

Parisi gives 150 μ for the length of polar filament.

MYXOBOLUS VOLGENSIS Reuss

[Figs. 446 to 448]

1906	<i>Myxobolus volgensis</i>	Reuss	1906 : 200-201
------	----------------------------	-------	----------------

Habitat: Branchiae, cornea and dorsal fin of *Lucioperca volgensis* Pall; Volga.

Vegetative form: Cysts, spherical, 0.3 to 1mm. in diameter. Polysporous.

Spore: Broad elliptic or rounded. Sutural edge has at least 3 folds. Sporoplasm with an iodophilous vacuole. Dimensions: length 8.25 to 9.5 μ , breadth 7.25 to 8.25 μ , thickness 4.5 to 5.5 μ , polar capsule 3 μ by 2 μ .

MYXOBOLUS SCARDINII Reuss

[Fig. 449]

1906 *Myxobolus scardinii* Reuss 1906 : 201Habitat: Branchiae of *Scardinius erythrophthalmus* L.; Volga.

Vegetative form: Cysts, elongated oval. Smaller cysts rounded oval, 0.8mm. by 0.5mm., the larger forms elongated, 1.2mm. by 0.5mm. Polysporous.

Spore: Broad elliptical. Sutural edge narrow, having folds. A larger triangular intercapsular process. An iodophilous vacuole in sporoplasm. Dimensions: length 11 to 12 μ , breadth 9 to 9.5 μ , thickness 4.5 to 5 μ , polar capsules 5 μ by 2.5 μ .

MYXOBOLUS PHYSOPHILUS Reuss

[Figs. 450 and 451]

1906 *Myxobolus physophilus* Reuss 1906 : 201-202

Habitat: Surface of air bladder of *Scardinius erythrophthalmus* L.; Volga.

Vegetative form: Cysts, rounded, 1.5mm. in diameter. Polysporous.

Spore: Oval, with attenuated anterior end. Sutural edge narrow and smooth. Polar capsules rather large. An iodophilous vacuole in sporoplasm. Dimensions: length 12 to 13 μ , breadth 8.25 to 9 μ , thickness 6.5 to 7 μ , polar capsules 6 μ by 2.5 μ .

MYXOBOLUS MACROCAPSULARIS Reuss

[Fig. 452]

1906 *Myxobolus macrocapsularis* Reuss 1906 : 202Habitat: Branchiae of *Blicca björkna* L.; Volga.

Vegetative form: Cysts, Elongated oval. Size: 1mm. by 0.5mm. Polysporous.

Spore: Oval with greatly attenuated anterior portion. Sutural edge broad and without any fold. Polar capsules rather large. An iodophilous vacuole in sporoplasm. Dimensions: length 11 to 13 μ , breadth 8.25 to 9.25 μ , thickness 5.5 μ , polar capsules 6 μ by 2.5 to 3 μ .

MYXOBOLUS SANDRAE Reuss

[Fig. 453]

1906 *Myxobolus sandrae* Reuss 1906 : 202-203Habitat: Muscle of *Lucioperca sandra* Cuv.; Volga.

Vegetative form: Cysts. Rounded, 0.5mm. in diameter. Polysporous.

Spore: Oval. Sutural edge broad with many distinct folds. An iodophilous vacuole in sporoplasm. Dimensions: length 9.25 to 10 μ , breadth 7.25 to 8.25 μ , thickness 4 to 5 μ , polar capsules 3.5 μ by 2 μ .

MYXOBOLUS BRAMAE Reuss

[Fig. 454]

1906 *Myxobolus bramae* Reuss 1906 : 203-204Habitat: Branchiae of *Abramis brama* L.; Volga.

Vegetative form: Cysts. Oval, 0.5mm. long, 0.25mm. broad. Polysporous.

Spore: Oval to nearly spherical. Sutural edge narrow and with indistinct folds. Two polar capsules, with a small triangular intercapsular process. An iodophilous vacuole. Dimensions: length 11 to 12 μ , breadth 9.25 to 10 μ , thickness 4.5 to 5.5 μ , polar capsules 4 to 5 μ by 2.25 μ .

MYXOBOLUS CYPRINICOLA Reuss

[Fig. 456]

1906 *Myxobolus cyprinicola* Reuss 1906 : 204Habitat: Branchiae of *Cyprinus carpio* L.; Volga.

Vegetative form: Cysts, oval, 0.5mm. by 0.3mm. Polysporous.

Spore: Elongated oval. Sutural edge narrow with many indistinct folds. An iodophilous vacuole. Dimensions: length 9.25 to 10 μ , breadth 7 to 7.25 μ , thickness 5 to 5.5 μ , polar capsules 4.5 μ by 2.5 to 3 μ .

MYXOBOLUS BALLERI Reuss

[Fig. 455]

1906 *Myxobolus balleri* Reuss 1906 : 204-205Habitat: Branchiae of *Abramis ballerus* L.; Volga.

Vegetative form: Cysts. Elongated, 1.5mm. by 0.5mm. Polysporous.

Spore: Oval, slightly pointed at the anterior end. A triangular intercapsular appendix. Sutural edge smooth. An iodophilous vacuole. Dimensions: length 11 to 12 μ , breadth 9.25 to 10 μ , thickness 5.5 to 6.5 μ , polar capsules 5.5 μ by 2.75 μ .

MYXOBOLUS SQUAMAE Keysselitz

[Figs. 457 to 459]

1908 *Myxobolus squamae* Keysselitz 1908 : 273-274Habitat: Inner surface of the scales of *Barbus fluviatilis* Agass.; Mosel and Neckar.Vegetative form: Form variable; rounded, oval, elongated or rarely branched. The outline of the body is not smooth but irregular with numerous small tooth-like projections with which the body comes in contact with the surrounding substance. The parasites seem to be able to dissolve the substance composing the scale. Length 50 to 800 μ . In one scale, one or many, up to 8, individuals were found. All showed only advanced stages of spore formation. The parasites are surrounded by a variously developed envelope of connective tissue. Polysporous.

Spore: Elongated oval. Two polar capsules, with 7 to 8 times coiled polar filament. A triangular intercapsular projection. Sporoplasm with an iodophilous vacuole. Dimensions: length 10 to 10.5 μ , breadth 8 to 8.5 μ , length of polar capsule 4.5 μ .

MYXOBOLUS CORDIS Keysselitz

[Figs. 460 and 461]

1908 *Myxobolus cordis* Keysselitz 1908 : 279-282

Habitat: Muscle of ventricle, rarely that of bulbus arteriosus of *Barbus fluviatilis* Ag., spores found in kidney, liver and spleen in the condition of somewhat scattered infiltration; Germany (Mosel and Neckar).

Vegetative form: Elongated, oval, sausage or club form. The body whitish, later yellowish. Size from 0.25 up to 4mm., usually 1 to 1.5mm. in length. Propagative stage and cysts observed. One end of the body is held more or less deeply in the muscle and is covered by cellular envelope as in *Myxobolus musculi*, while remaining larger portion of the body is suspended freely inside of the ventricle, covered with a thin layer probably of endocardiac cells. Fish 30 to 45cm. long harboured 40 to 60 parasites. No movements. Ultimately the cysts are formed with differentiated protoplasm. Polysporous.

Spore: Oval. Shell very thin at the anterior end. At the posterior end, cell-like appendage, 2 to 3 μ wide which is probably formed by both valves, is present. Two pyriform polar capsules at the anterior end, which show the polar filament coiled 7 to 8 times. Sporoplasm with a comparatively large and oval iodophilous vacuole and two nuclei, rarely one (syncaryon). Dimensions: length 12 μ , breadth 10 μ , length of polar capsule 4.5 μ .

MYXOBOLUS MUSCULI Keysselitz

[Figs. 462 to 464]

1908 *Myxobolus musculi* Keysselitz 1908 : 282-286

Habitat: Muscle of the main body, rarely that of fins and operculum, and kidney of *Barbus fluviatilis* Agass. of various size (youngest fish found infected, 2 months old), spores in liver, spleen, kidney and ovary (not the ovum) in diffuse infiltration; Mosel and Neckar.

Vegetative form: Elongated. Body whitish opaque, with differentiated protoplasm. Smallest individual observed, 24 μ . Large form 2mm. in length. Many trophozoites are found closely situated, forming a large mass of parasites that reached dimensions of 4mm. by 2mm. The surrounding envelope, varying in thickness, composed of cells with elongated nuclei as those of perimysium. Young cysts surrounded by thin layer of ectoplasm. Polysporous.

Spore: Oval. Two polar capsules usually unequal. Shell as in *M. cordis* with a small peg closer to the anterior end, polar filament coiled 4 to 5 times, visible in the capsule. Sporoplasm with rarely one (syncaryon), but usually two nuclei and an iodophilous vacuole. A posterior process as is seen in the spores of *M. cordis*, but much smaller, was occasionally observed. Dimensions: length 11μ , breadth 8μ , polar capsules 6μ and 4μ long.

MYXOBOLUS sp. Miyairi

1909 *Myxobolus* sp. Miyairi 1909 : 126

Habitat: Branchiae of loach (*Misgurnus anguillicaudatus* Cant.?); Fukuoka? (Nippon).

Vegetative form: Cysts were not observed.

Spore: No description.

MYXOBOLUS sp. Wegener

[Fig. 465]

1910 *Myxobolus* sp. Wegener 1910 : 78

Habitat: Branchiae (gill-arch) of *Perca fluviatilis* L.; Germany (Frisches Haff, March). Only one case.

Vegetative form: Cysts on a gill-arch, white and round, with a diameter of 1.1mm. Polysporous.

Spore: Form and size very variable. Rounded or elliptical, pointed at the anterior end. Sutural edge showing folds at the posterior portion. Dimensions: length 8 to 10μ (in round form) and 11μ (in elliptical form), breadth 8 to 9μ , polar capsules 4 to 5μ by 2 to 3μ , length of polar filament 40μ .

MYXOBOLUS PERMAGNUS Wegener

[Fig. 466]

1910 *Myxobolus permagnus* Wegener 1910 : 78-79

Habitat: Branchiae and operculum of *Perca fluviatilis* L., air bladder of *Scardinius erythrophthalmus* L.; Königsberg (May), Pregel (March).

Vegetative form: Cysts rounded in form and white in color, resemble to those of *M. gigas*. No clear ectoplasm layer, nor typical protoplasmic structure. Polysporous.

Spore: Oval, sharply pointed at the anterior end. Sutural edge with 5 to 6 distinct folds at the posterior portion. Polar filament visible in the polar capsules. Dimensions: length 17 to 18μ , breadth 10 to 13μ , polar capsules 7 to 8μ by 3.5 to 4μ .

MYXOBOLUS ROTUNDUS Nemeček

[Fig. 467]

1911 *Myxobolus rotundus* Nemeček 1911 : 156-157

Habitat: Branchiae of *Abramis brama* L.; Austria.

Vegetative form: Cysts, ovoidal or spindle form, 1 to 3mm. long and 1 to 1.5mm. wide. Body white. An extraordinary large number of spores were found in the cysts. Polysporous.

Spore: Round or slightly oval, when viewed from the front. Greatly flattened in side view. Polar capsules convergent, with no intercapsular body. Shell smooth. Sutural edge narrow, without folds. Dimensions: length 10μ , breadth 9.8μ , thickness 3μ , polar capsules 3.8 to 5μ long, length of polar filament 40μ .

MYXOBOLUS MINUTUS Nemeček

[Fig. 468]

1911 *Myxobolus minutus* Nemeček 1911 : 160

Habitat: Branchiae of *Leuciscus* sp.; Austria.

Vegetative form: Cysts spherical, oval or elongated with white color. Size: 0.5 to 3mm. by 0.5 to 1mm. Polysporous.

Spore: Rounded oval, similar to that of *Myxobolus rotundus*. Shell smooth. Sutural edge narrow without folds. Sporoplasm with an iodophilous vacuole. No intercapsular appendix. Dimensions: length 6μ , breadth 4.2 to 5μ , polar capsule 3μ by 2μ , length of polar filament 50 to 60, often 70μ .

MYXOBOLUS sp. Lebzelter

1912 *Myxobolus* sp. Lebzelter 1912 : 296-297

Habitat: Gall-bladder of *Thymallus thymallus* L.

Vegetative form: Not observed.

Spore: Sutural ridge distinct. Dimensions, length 5μ , breadth 3μ .

MYXOBOLUS MAGNUS Awerinzew

[Figs. 469 and 470]

1913 *Myxobolus magnus* Awerinzew 1913 : 75-76

Habitat: Eye of *Acerina cernua* L.; Petrograd.

Vegetative form: Trophozoites form white spots in the tissue of iris, with many spores (300 to 400). Each pansporoblast forms in most cases two, sometimes 3 or 5 spores! Polysporous.

Spores: Large, elongated roundish, slightly flattened. Sutural edge somewhat thick, forming a wide ridge, with 4 to 5 folds at the posterior portion. Polar capsules do not cross each other. Sporoplasm with an iodophilous vacuole and two nuclei. Dimensions: length 38 to 45μ , breadth 32 to 38μ , thickness 28 to 35μ , length of polar capsules 15 to 17μ , diameter of the vacuole 12 to 16μ .

MYXOBOLUS CARASSII Klokačewa

[Figs. 471 to 473]

1914 *Myxobolus carassii* Klokacewa 1914 : 182-184

Habitat: Body cavity, liver and intestine of *Carassius vulgaris* L.; Petrograd?

Vegetative form: Cysts spherical. Those in liver and intestine yellowish, surrounded by an envelope composed of fibrous connective tissue. Secondary cysts are formed. Polysporous.

Spore: Oval, in front view. Two ovoidal polar capsules convergent at the slightly attenuated anterior end. Coiled polar filament visible. Sporoplasm with an iodophilous vacuole and two nuclei. Sutural edge shows folds in some cases. Dimensions: length 13 to 17 μ , breadth 8 to 10 μ , thickness 5 to 7 μ , polar capsules 6 to 7 μ long.

Remarks: Compare with *Myxobolus* sp. Gurley on page 142.

MYXOBOLUS sp. Southwell

1915	<i>Myxobolus</i> sp.	Southwell	1915 : 312-313
------	----------------------	-----------	----------------

Habitat: Subcutaneous intermuscular tissue of *Rasbora (Cyprinus) daniconius* Day; from a stream near Katwan, Mirzapore (U.P.), India.

Vegetative form: 6 cysts found on four fish. The seat is immediately below the scales, in the epidermis. Color milky white. Soft, flattened and roughly oval in shape. Greatest length found, 1.1mm. No pigment was present on the cyst.

Spore: Two equal capsules, with a very short tail-like process. Sporoplasm with vacuole; iodine treatment could not be carried out. Dimensions: length 13 μ , breadth 13 μ , polar capsule 4 μ by 4 μ (?).

Remarks: Dimensions, especially that of polar capsule seem to be misprinted. Southwell gave one figure of a fish with a cyst near the dorsal fin. He thinks that "it is quite possible that our parasites may belong to *Myxobolus cyprini*." The incomplete observation without any figure, leads the writer to leave the form also as *Myxobolus* sp. Southwell.

MYXOBOLUS FUNDULI Kudo

[Figs. 474 to 476]

1915	<i>Myxobolus musculi</i>	Hahn	1915 : 201-205
1917	<i>Myxobolus musculi</i>	Hahn	1917 : 91-104

Habitat: Branchiae and muscle of *Fundulus heteroclitus*, *F. majalis*; Woods Hole. Hahn claims that he succeeded in causing experimental infection in *F. diaphanus* and *Cyprinodon variegatus* by inoculation.

Vegetative form: Hahn uses quite a number of different terms from those that are ordinarily used in describing Myxosporidia, without giving any definitions. Naturally it is hard to put what he wrote in several pages in the following lines. Granular vegetative forms produce a great many pansporoblasts, each with a single spore. "Trophoplasm" is difficult to stain. Size: 74 by 33 μ , 24 by 19 μ . Cysts within and between the muscle fibers, containing several hundred spores.

Spore: Hahn's descriptions may be summarized as follows: Dimensions: length 14.3 μ , breadth 6.7 μ , thickness 6.7 μ to 2/3 of width, polar

capsule, 6.5μ by 2μ , polar filament 3 to 4 times the length of the spore (42.9 to 57.2μ). Polar filament coiled 10 to 14 times. Shell thin, almost invisible. The spores found in the gill: length 12 to 13.4μ , breadth 6μ to 10.4μ . A vacuole is present in the sporoplasm.

Remarks: Examination of Hahn's first paper suggested that he was dealing with the present form as a new species tho he did not mention at all Keysselitz who gave the name *Myxobolus musculi* Keysselitz to the parasite of *Barbus fluviatilis* from German rivers. I was informed by Hahn that he gave the name, *Myxobolus musculi*, without knowing the fact that it was preoccupied by Keysselitz (1908) (see page 148) and that tho he became aware of it later, he can not determine differences by which the two forms can be distinguished. A comparison of the descriptions of Keysselitz and Hahn, however, shows that these two forms differ in several respects. Hence the latter form is recorded here as a distinct species under the new name.

It is interesting to note that very similar forms, one without an iodophilous vacuole at any stage of spore-formation (*Myxosoma funduli* Kudo, see page 125) and the other with a vacuole, occur in the same hosts in the same locality. As mentioned above, the reader is requested to refer to Hahn's original paper for further data.

MYXOBOLUS PLEURONECTIDAE Hahn

[Fig. 477]

1917 *Myxobolus pleuronectidae* Hahn 1917 : 160-161

Habitat: Subcutaneous muscular tissue of *Pseudopleuronectes americanus*; Woods Hole.

Vegetative form: Similar to that of *Myxobolus funduli*.

Spore: Hahn writes as follows: Shape and appearance resembles *Myxobolus pfeifferi*. Dimensions: length 14.5μ , breadth 11.9μ , polar capsules 6μ by 3.7μ .

MYXOBOLUS CAPSULATUS Davis

[Fig. 478]

1917 *Myxobolus capsulatus* Davis 1917 : 237

Habitat: Visceral connective tissues of *Cyprinodon variegatus*; Beaufort.

Vegetative form: Irregular form. In the state of diffuse infiltration. Polysporous.

Spore: Pyriform, flattened. Polar capsules large and pyriform, filling almost entire cavity of the spore. Sporoplasm relatively small. Iodophilous vacuole visible in living spore. Dimensions: length 16μ , breadth 10 to 11μ , polar capsules 11μ by 4μ , length of polar filament 84μ .

MYXOBOLUS NODULARIS Southwell et Prashad

[Figs. 479 and 480]

1918 *Myxobolus nodularis* Southwell and Prashad 1918 : 347

Habitat: In the muscles of *Rasbora daniconius* occurring in two fish on the sides, and in another as a globular cyst near the anus; Mirpur, Dacca (June). Type specimens, numbered P 52/1.

Vegetative form: Cysts rounded or slightly elongated, varying in length 3.5 to 3.8mm. and 2.3 to 2.8mm. in width. Creamy yellow in color, in one case appearing blackish owing to the large number of black granules scattered in its surface.

Spore: Ovoidal. Sutural ridge very wide (about 1/5 thickness of the spore). Two polar capsules of equal size, which show coiled polar filaments clearly. Dimensions: length 9μ , breadth 7.2μ , length of polar capsule 3.4μ , that of polar filament 18.3μ .

MYXOBOLUS HYLAE Johnston et Bancroft

[Figs. 591 to 593]

1888		Fletcher	1888 : 337
1890		Haswell	1890 : 661
1909	<i>Myxobolus</i> sp.	Johnston	1909 : 29
1910	<i>Myxobolus</i> sp.	Cleland and Johnston	1910 : 25
1918	<i>Myxobolus hylae</i>	Johnston and Bancroft	1918 : 171-175

Habitat: In the testes, vasa efferentia and oviducts of *Hyla aurea*; Sidney, Australia (April, other months not mentioned). Fletcher observed the parasites also in the urinary bladder of both sexes, which fact was not confirmed by Johnston and Bancroft on account of the scarcity of the material. The latter authors could not infect *Hyla caerulea* by feeding infected testes or the cysts, giving the conclusion that the parasite is specific to *H. aurea*. The male is more often attacked by the parasite than the female. The infected animal appeared sickly and emaciated. As to the infection in kidneys, they write as follows: In one male specimen both testes and both kidneys were affected, and the upper parts of the ureters adjacent to the kidneys were swollen and milky in appearance. In another, in addition to the testes, the adjacent kidney and mesentery were attacked. No spores have yet been detected by them in sections of the kidney tubules.

Vegetative form: Johnston and Bancroft describe as follows:

Cysts: in male, either imbedded in the tissue or may project freely into the coelom of the testes; in female, lying between the layers of the wall, being projected into the lumen of the oviduct. Size from those of microscopic dimensions up to 2 to 3mm. in diameter. In sections, the protoplasm

is differentiated into two regions. The outer layer (ectoplasm) surrounds the body as a thin, light-staining region, while the endoplasm being denser and of more or less granular structure filled with spores especially in the central portion.

Spore: Johnston and Bancroft describe as follows:

Form somewhat variable, caused by the reduction in length. Oval, egg-shaped or nearly circular in front view. Sutural ridge slightly thickened. Two pyriform polar capsules are located at the anterior end. Sporoplasm with an iodophilous vacuole (2μ in diameter), shows usually two distinct nuclei, rarely but one. Dimensions: length variable, diameter of circular form 7 to 8μ , breadth 8 to 10μ , thickness about 6μ , thickness of shell 1μ , polar capsules 4 to 5μ by 2μ , length of polar filament 90 to 98μ (acids or alkalies).

MYXOBOLUS AUREATUS Ward

[Figs. 643 to 649]

1919

Myxobolus aureatus

Ward

1919 : 49

Habitat: Between the ectodermal layers of the fin membrane of *Notropis anogenus*; Put-in-Bay, Lake Erie (August). Out of thirty fish, two to three cm. in length, seven were found to be infected. The infected fish were not inferior in size or vigor to others of the same species. The most heavily infected one was the most vigorous of all. The number of cysts in the individual fish, varied from one to forty, being confined in the fin. The cysts are always separated from each other, tho in a few instances they were apparently connected.

Vegetative form: The parasite forms cysts between the ectodermal layers of the fin membrane. The cyst is a smooth margined ellipsoid, measuring from 1 to 1.6 mm. in layer diameter and from 0.8 to 1.2 mm. along its transverse axis. The opaque cyst is of a clear orange yellowish color. This gilt color is contained in the cyst wall, fading away in alcohol and formol. The chromatophores of the skin of the host are distinctly more abundant on the cyst than in other parts of the skin, and the older the cyst the more abundant the chromatophores. The wall of the cyst is noticeably tough and thick. In section, the protoplasm shows a poor differentiation into ectoplasm and endoplasm. The former granular and reticular, covers the entire surface as a thin layer, while the latter is highly vacuolated, containing only mature spores. Polysporous.

Spore: Ovoid; slightly pointed anterior and rounded posterior ends in front view; slightly compressed in lateral view. Sutural ridge distinct. The shell is of moderate thickness, and bears a flange at the posterior half in some spores. Two pyriform polar capsules, frequently of slightly different dimensions, are at the anterior part of the spore. No intercapsular appendix is present. When the spore is allowed to stand for 24 hours

or more in water, the polar filaments are extruded. The binucleated finely granular sporoplasm shows an iodophilous vacuole. Dimensions: length 12.4 to 13.5 μ , breadth 6.5 to 7.5 μ , thickness 5 μ , length of polar capsule 6 to 7 μ (rarely 7.5 μ), length of polar filament about 20 to 26 μ , diameter of iodophilous vacuole about 2 μ .

MYXOBOLUS MIYAIRII nov. spec.

[Fig. 481]

1909 *Myxobolus* sp.

Miyairi

1909 : 130, 131-132

Habitat: Intestinal wall of *Parasilurus asotus* L.; Fukuoka ? (Nippon)

Vegetative form: Cysts. Size rather small up to 0.5mm. Full-grown spores as well as those in developmental stages fill the central portion of cysts, while numerous nuclei are chiefly found along the periphery of endoplasm.

Spore: Elongated elliptic. Two polar capsules of nearly same size. Sporoplasm with a comparatively large iodophilous vacuole. Dimensions: length 13 to 14.5 μ , breadth 6 to 7 μ , length of polar capsules 4.5 μ , length of polar filament 30 to 35 μ .

Remarks. As the descriptions show the form and structure are distinguishable from other species, the writer establishes the present species.

MYXOBOLUS KOI nov. spec.

[Figs. 482 to 485]

Habitat: In the connective tissue of the gill filament of *Cyprinus carpio* L.; Tokio (April). One fish was found infected in a slight degree.

Vegetative form: Cysts small and spherical; white in color. Size up to 230 μ in largest diameter. The seat similar to *Myxobolus toyamai*. The structure of the cysts, observed in section preparations, is also similar to the above mentioned unicapsular *Myxobolus*.

Spore: Oval with attenuated anterior and rounded posterior ends in front view; elongated pyriform in side view. Shell comparatively thin. No marking on shell. Sutural ridge fairly well marked. No intercapsular appendix. Two polar capsules are pyriform, large, and of usually equal form and size. Coiled polar filament distinct in vivo. Sporoplasm rather small, finely granular, shows two nuclei in almost all spores. An iodophilous vacuole is deeply stained by Lugol's solution. Dimensions: 14 to 16 μ , breadth 8 to 9 μ , thickness 5 to 6 μ , polar capsule 8 to 9 μ by 2.5 to 3 μ , length of polar filament 72 μ in average (KOH).

MYXOBOLUS ORBICULATUS nov. spec.

[Figs. 566 to 576]

Habitat: Muscle of myotomes of *Notropis gilberti* J. et M.; Stony Creek, Ill. (November). The fish was kept alive in an aquarium from November

11, 1918, until March 10, 1919, when it was killed, being then nearly dead. The material was examined on March 15. A few isolated spores occurred in the muscle of *Notropis blennioides* (Homer Park, Ill., November).

Vegetative form: In and between the muscle bundles of the myotomes. Size variable. Color opaque white under the dissecting microscope. Smallest rounded ameboid forms with a single or numerous nuclei, in the muscle bundle, have the size of from 10μ to 30μ in greatest diameter (Figs. 573 to 575). The largest form observed was 400μ by 120μ . Young forms without any differentiated protoplasm, shows indistinct granular and reticular structure with deeply staining spherical or ring-form chromatinic granules. The number of the nuclei increases with the growth of the body. Larger form (Fig. 576), spindle shape, circular in cross-section, lies with its long axis parallel to the muscle fibres. The protoplasm vacuolated, contained mostly mature spores. Spores were also found in the state of diffuse infiltration. Polysporous.

Spore: Form somewhat variable. Typical form almost circular, slightly pointed at the anterior end (Fig. 566) in front view; spindle shaped in profile (Figs. 569 and 570). Sutural ridge marked. Shell uniformly thick, usually exhibiting four triangular folds on the surface along the posterior margin (Figs. 566, 568 and 571). No intercapsular appendix. Two pyriform polar capsules are, as a rule, of the same size and form. Frequent occurrence of the inequality of the polar capsules together with abnormalities in the form of the spore, were noticed especially among comparatively young spores. The granular sporoplasm, shows two spherical nuclei when stained. The iodophilous vacuole, spherical and 2μ in average diameter, is deeply stained with Lugol's solution. Dimensions of unstained preserved spores: length and breadth 9 to 10μ , thickness 6.5 to 7μ , polar capsule 6 to 7.5μ by 2.5 to 3μ .

MYXOBOLUS DISCREPANS nov. spec.

[Figs. 597 to 601]

Habitat: Branchial lamellae of *Carpiodes difformis*; Salt Fork, Urbana, U.S.A. (May). One fish caught, died (soon after the capture) two hours before being fixed. Length 8.5cm.

Vegetative form: The parasites formed numerous cysts on the branchial lamellae. Cysts slightly yellowish white and mostly rounded or elongated along the lamella, occur in groups, often occupying the entire lamella. Infection was fairly heavy. Every gill arch harbored ten to twenty cysts mostly on the outer surface. Size of the cyst varies, small rounded one 500μ in diameter up to elongated forms 2mm. by 0.5mm., the majority being from 0.5 to 1mm. in diameter. The cyst is surrounded by a thin connective tissue layer of the host. The protoplasm shows little differen-

tiation. The ectoplasm is a rather narrow zone around the entire body and the endoplasm is filled with various nuclei, several stages of developing pansporoblasts, and mature spores. Each pansporoblast produces two spores. Polysporous.

Spore: Approximately circular with broad anterior and more or less narrower posterior end in front view; broadly fusiform in profile. Shell uniformly thin with 5 to 6 markings on the posterior margin. Two polar capsules broadly oval and convergent, fill the anterior half of the spore. A small triangular intercapsular appendix presents. Coiled polar filament is fairly visible in vivo. The spores from the cysts which were fixed with alcohol-acetic and preserved in 95 per cent alcohol, showed the extrusion of the polar filament under the influence of potassium hydrate solution (35 per cent) even after a considerable length of time as is shown in the following:

Material fixed on May 29.

June 2; Extrusion took place in almost all spores.

June 10; Extrusion took place in almost all spores.

June 26; Extrusion took place in almost all spores.

July 28; Extrusion took place in almost all spores.

August 29; Extrusion took place in numerous spores.

September 29; Extrusion took place in about 70 per cent of the spores, some filaments being rather short, and not fully extended.

October 20; Extrusion took place in about 50 per cent of the spores, most filaments being short, and not fully extended.

Sporoplasm coarsely granular shows clearly two ring-form nuclei in fresh preparations. Dimensions of preserved spores: length 11.4 to 13.5 μ , breadth 9.5 to 11 μ , thickness 8.5 to 9.5 μ , polar capsule 5.5 to 6 μ by 3.5 to 4 μ , length of polar filament 50 to 55 μ .

Remarks: The present species differs from the hitherto known species. *Myxobolus lintoni* (page 138) and *Myxobolus orbiculatus* (page 155) which are the nearest to the present form, differ from *Myxobolus discrepans* in the host, organ of infection, vegetative form and form and structure of the spore.

MYXOBOLUS MESENTERICUS nov. spec.

[Figs. 628 to 631]

Habitat: In the mesentery, liver, spleen and wall of stomach, pyloric coecum, intestine, and gall-bladder of *Lepomis cyanellus*; Crystal Lake, Urbana, Ill. (June and July). Out of thirty-six host fish, 10 cm. in average length, seven were found to be infected. In every case, except one, the mesentery was the main seat of infection, harboring conspicuous cysts. The number of cysts found in the host body varied from three to seven. The infected fish did not exhibit any recognizable pathological changes. Other species of fish caught at the same time, were free from the infection.

Vegetative form: The cysts are mostly spherical in form, and are covered by a tough resistant envelope composed of the connective tissue of the host. They are uniformly white in color, and have the variable dimensions of from 0.5 to 1.5mm. in diameter. In section, the protoplasm shows a coarsely reticulated structure without distinct differentiation. In all cysts of various sizes fully mature spores were only observed. The spore formation could not be worked out. Polysporous.

Spore: Broadly oval with a slightly truncated anterior end in front view (Fig. 628), lenticular in side or end view (Fig. 629). No intercapsular appendix is seen. The shell is rather thick, and shows about eight folds on the sutural edge, two of which located laterally being more conspicuous than others. The sutural ridge is rather fine. Two convergent polar capsules equal in size occupy the anterior half of the spore. The coiled polar filament becomes more distinctly visible with the addition of Lugol's solution, altho it is faintly observable in fresh state. Fresh spores extruded their polar filaments under the action of potassium hydrate solution. In some spores, the extruded filaments cross each other near the foramina. The preserved spores showed no extrusion of the filament as in the last species. The sporoplasm is extremely finely granulated. The iodophilous vacuole is comparatively large. When stained, the spore shows two nuclei in the sporoplasm. Dimensions of fresh material: length 10 to 11.5 μ , breadth 8.5 to 9.5 μ , thickness 6.5 μ , polar capsule 4.75 μ by 1.5 to 2 μ , length of polar filament 32 to 40 μ . Average dimensions of unstained preserved spores: length 9.5 μ , breadth 8 μ , polar capsule 4.75 μ by 2 μ .

Remarks: The habitat and the structure of the spores, lead the writer to record the species as a new species.

Genus HENNEGUYA Thélohan

1892	<i>Henneguya</i>	Thélohan	1892 : 167, 176
1895	<i>Henneguya</i>	Thélohan	1895 : 352

The characters of the genus are described on page 59.

Type species: *Henneguya psorospermica* Thélohan.

HENNEGUYA PSOROSPERMICA Thélohan

[Figs. 486, 487 and 496]

1895	<i>Henneguya psorospermica</i>	Thélohan	1895 : 353
1896	<i>Myxobolus psorospermica</i> s. str. Cohn		1896 : 261
1899	<i>Henneguya psorospermica</i> <i>typica</i>	Labbé	1899 : 101
1905	<i>Henneguya psorospermica</i>	Nufer	1905 : 77, 185
1910	<i>Henneguya psorospermica</i>	Wegener	1910 : 81-82
1911	<i>Henneguya psorospermica</i> <i>typica</i>	Auerbach	1911 : 5, etc.

Habitat: Branchiae of *Esox lucius* L. and *Perca fluviatilis*; France,

Frisches and Kurisches Haff, Pregel, Masurische Seen (all the year round, but rarer in Winter) Switzerland.

Vegetative form: Thélohan's observations on the structure of the cyst, are as follows: The surface of the cyst is covered by a layer, homogeneous, refringent and deeply stained, with which the cyst comes in direct contact with the surrounding epithelial cells of the host. Inside of this layer, there is a "pseudoectoplasmic" zone, in which the protoplasm is dense at places, forming radiate irregular striations, enclosing numerous irregular masses which are composed of apparently the same substance that forms the external layer. Toward the central portion of the cyst, there are masses of spores (Fig. 496).

Cohn's descriptions are as follows: The purely white cyst is elliptical; length 1.15mm. and breadth 0.85mm. The seat is under the epidermis. It is surrounded by the host tissue with small, elongated and scattered nuclei. The outer layer of the cyst is a thin membranous protoplasm.

Wegener writes as follows: The white cysts are round or elliptical, usually on the upper end of the branchial lamella. Size of larger cysts, 1.5 to 2mm. long and 1.1 to 1.5mm. wide.

Spore: Elongated; anterior part fusiform and anterior end blunt. Polar capsules elongated and parallel to each other. Coiled polar filament visible in fresh conditions. Shell unstriated. Dimensions: total length 40μ in average, largest breadth 7μ , length of polar capsule 7 to 8μ .

Cohn's form is described by him as follows: Spore narrow with blunt anterior end. Sporoplasm with 6 horns (no figure to explain this expression!). When kept in water, sporoplasm takes round form and becomes highly refractive. Dimensions: length 29 to 38μ , length between the tip and the posterior margin of the cavity (15 to 20μ) 18μ , breadth 9 to 10μ , polar capsule (8 to 11μ) 9μ by 2μ , length of "starren Fäden" 14μ , length of tail 14 to 18μ .

Wegener's form is as follows: total length 35 to 38μ , breadth 7 to 8μ , length of the spore cavity 15μ , length of tail 15 to 20μ , polar capsule 8μ by 2 to 3μ .

HENNEGUYA TEXTA (Cohn) Labbé

1895	<i>Myxobolus textus</i>	Cohn	1895 : 38-39
1899	<i>Henneguya psorospermica texta</i>	Labbé	1899 : 101
1910	<i>Henneguya texta</i>	Wegener	1910 : 82-83

Habitat: Branchiae of *Perca fluviatilis* L.; Pregel, Frisches and Kurisches Haff (all the year round).

Vegetative form: Cohn observed as follows: Cyst distinctly elliptical. Length 0.75mm., breadth 0.375mm. The cysts surrounded by a thick layer of the host tissue. In the peripheral portion of the cyst, the protoplasm exhibits a network-like structure which forms a fibrous structure further inside, crossing the cyst at right angles to the long axis of the cyst.

Wegener writes as follows: The white cysts are elongated, 1.2 to 1.8mm. long and 0.5 to 0.7mm. wide.

Spore: Cohn mentions dimensions exactly the same as those of *Henneguya psorospermica* and can not distinguish the two species by the spore.

Wegener gives the following dimensions: length 30 to 40 μ , breadth 7 to 8 μ , length of the cavity of spore 15 to 18 μ , length of tail 15 to 25 μ , polar capsule 8 μ by 2 to 3 μ .

HENNEGUYA MINUTA (Cohn) Labbé

[Figs. 488 and 489]

1895	<i>Myxobolus minutus</i>	Cohn	1895 : 39-40
1899	<i>Henneguya psorospermica</i> <i>minuta</i>	Labbé	1899 : 102

Habitat: Branchiae of *Perca fluviatilis* L.; Frisches Haff, Lesina.

Vegetative form: Cohn's description is as follows: Cysts oval and small, difficult to distinguish them from those of *Henneguya psorospermica*. Size, 130 μ by 115 μ . The parasite was met only once. But the number of the cysts was far greater than that of *Henneguya psorospermica*, often 5 to 6 on one lamella, reaching up to 200 cysts on a single gillarch.

Spore: Cohn gives the following dimensions: total length (28 to 45 μ) about 36 μ , length from the tip to the end of cavity (20 to 28 μ) about 26 μ , breadth 10 to 11 μ , thickness 8 μ , polar capsule 11 to 14 μ by 2 to 3 μ , length of polar filament 42 to 45 μ , length of tail (8 to 17 μ) 12 μ . Cohn gives a figure (Fig. 489) of a spore with two vacuoles(?).

HENNEGUYA OVIPERDA (Cohn) Labbé

[Figs. 490 and 491]

1892		Weltner	1892 : 28-36
1895	<i>Myxobolus oviperdus</i>	Cohn	1895 : 40-41
1899	<i>Henneguya psorospermica</i> <i>oviperda</i>	Labbé	1899 : 102
1904	<i>Henneguya psorospermica</i> <i>oviperda</i>	Fuhrmann	1904 : 469-471
1911	<i>Henneguya psorospermica</i> <i>oviperda</i>	Auerbach	1911 : 5-22
1911	<i>Henneguya psorospermica</i> <i>oviperda</i>	Nemeczek	1911 : 146

Habitat: Ovary of *Esox lucius* L.; Switzerland, Berlin, Frisches Haff (all the year round), Upsala (May), Austria (December).

Vegetative form: Cohn writes as follows: No real cyst exists. The parasite occupies the ovum.

Auerbach, however, mentions the presence of cysts in the connective tissue and follicle epithelium of the ovary. Dimensions, 1mm. up to 5 or 6mm. in diameter.

Spore: Cohn states the form and dimensions are very much similar to those of *H. psorospermica*.

HENNEGUYA LOBOSA (Cohn) Labbé

[Figs. 492 and 493]

1895	<i>Myxobolus lobosus</i>	Cohn	1895 : 42
1899	<i>Henneguya psorospermica lobosa</i>	Labbé	1899 : 102
1910	<i>Henneguya(?) lobosa</i>	Wegener	1910 : 83
1911	<i>Henneguya(?) lobosa</i>	Auerbach	1911 : 22-25

Habitat: Branchiae of *Esox lucius* L.; Frisches Haff, Pregel, Karlsruhe.
Vegetative form: Cysts irregular in shape, size up to 2.5mm.

Wegener noticed that the cyst resembles that of *Myxosoma dujardini* with the dimensions of 2.2 to 2.8mm. by 1 to 1.1mm.

Spore: Cohn gives the dimensions as follows: total length 30 to 40 μ , length from the tip to the posterior margin of cavity 11.5 to 15 μ , breadth 5 to 6.5 μ , polar capsules 6.5 to 8 μ by 2 to 2.5 μ , length of tail 22 to 28 μ .

Wegener's form: oval; length 35 to 40 μ , breadth 5 μ , polar capsule 6 to 7 μ by 2.5 to 3 μ , length of the cavity of spore 13 to 15 μ , length of tail 20 to 25 μ , the iodophilous vacuole could not be detected.

Auerbach gave the following dimensions: total length 30 μ , breadth 4 to 6 μ , length of polar capsule 6 μ , length of polar filament 48 to 54 μ .

Remarks: Wegener and Auerbach did not observe the iodophilous vacuole.

HENNEGUYA PERI-INTESTINALIS Cépède

1906	<i>Henneguya psorospermica peri-intestinalis</i>	Cépède	1906 : 67
1907	<i>Henneguya psorospermica peri-intestinalis</i>	Cépède	1907 : 137
1912	<i>Henneguya psorospermica peri-intestinalis</i>	Parisi	1912 : 295

Habitat: Intestine of *Esox lucius* L.; Lac du Bourget, Pavia. (June).
Vegetative form: Cysts.

Spore: Cépède mentions that it resembles that of *Henneguya psorospermica*.

HENNEGUYA MEDIA Thélohan

[Figs. 494 and 495]

1890		Thélohan	1890 : 198-200
1892	<i>Henneguya media</i>	Thélohan	1892 : 177
1894	<i>Myxobolus medius</i>	Gurley	1894 : 248
1895	<i>Henneguya media</i>	Thélohan	1895 : 353
1898	<i>Henneguya media</i>	Doflein	1898 : 342

Habitat: Renal tubules of kidney and ovary of *Gasterosteus aculeatus* and *G. pungitius* L.; France. Mixed infection with *Sphaerospora elegans*.

Vegetative form: Rounded or elongated. In larger individuals, clear differentiation of protoplasm. Monosporous (?) and polysporous.

Spore: Fusiform. Shell striated. A vacuole in sporoplasm. Dimensions: length 20 to 24 μ , breadth 5 to 6 μ , polar capsules 4 to 5 μ . Tail short.

HENNEGUYA BREVIS Thélohan

1854		Lieberkühn	1854 : 357
1892	<i>Henneguya brevis</i>	Thélohan	1892 : 177
1895	<i>Henneguya brevis</i>	Thélohan	1895 : 354

Habitat: Similar to *H. media* Thélohan.

Vegetative form: Undescribed.

Spore: Fusiform with short tail. Dimensions: length 14 to 15 μ , breadth 5 to 6 μ , polar capsules 1.4 to 5 μ , tail 4 to 5 μ long.

HENNEGUYA SCHIZURA (Gurley) Labbé

[Figs. 497 to 499]

1841		Müller	1841 : 477-478
1893	<i>Myxobolus schizurus</i>	Gurley	1893 : 417
1894	<i>Myxobolus schizurus</i>	Gurley	1894 : 255
1899	<i>Henneguya schizura</i>	Labbé	1899 : 102-103

Habitat: In cellular tissue of the eye muscles, in that of the sclerotic, and in that between the sclerotic and choroid of *Esox lucius* L.; Germany, U. S. A.

Vegetative form: Cysts white; membrane delicate; 0.44 to 1.09mm. in diameter.

Spore: Oval. Dimensions: length 12 μ , breadth 6 μ , thickness one-half the breadth, tail 3 to 4 times length of the body.

HENNEGUYA CREPLINI (Gurley) Labbé

[Figs. 500 to 503]

1842		Creplin	1842 : 61-63
1894	<i>Myxobolus creplini</i>	Gurley	1894 : 248-249
1899	<i>Henneguya creplini</i>	Labbé	1899 : 103
1910	<i>Henneguya creplini</i>	Wegener	1910 : 84

Habitat: Branchiae of *Acerina cernua* L.; Pregel (March), Frisches and Kurisches Haff.

Vegetative form: Wegener describes as follows: Cysts, usually elongated oval and are located at the end of branchial lamella. Color white. Size 1 to 1.1mm. by 0.5mm. During winter, the cyst has only pansporoblasts, but no fully grown spores.

Spore: Creplin writes as follows: Elongated elliptical. Length 1/120''', breadth 1/360''', tail about as long as or a little longer than the body.

Wegener's form: elongated spindle shape; length 20 μ , breadth 8 to 9 μ , polar capsule 8 μ by 2 to 3 μ (parallel to each other).

Remarks: Wegener thinks that the present species and *Henneguya acerinae* Schröder, are one and the same species, and that the differences between the dimensions are due to the miscalculation of measurement in lines given by Creplin on the part of Gurley and Labbé.

HENNEGUYA LINEARIS (Gurley) Labbé

[Fig. 504]

1841		Müller	1841 : 489
1893	<i>Myxobolus linearis</i> (part)	Gurley	1893 : 417
1894	<i>Myxobolus linearis</i>	Gurley	1894 : 255
1899	<i>Henneguya linearis</i>	Labbé	1899 : 103

Habitat: Membrane lining branchial cavity of *Pimelodus sebae* Cuv. et Val., branchiae of *Platystoma fasciatum* L.; South American rivers.

Vegetative form: Not described.

Spore: Very narrow. Length 3 to 4 times breadth.

HENNEGUYA GURLEYI Kudo

[Fig. 505]

1893	<i>Myxobolus linearis</i> (part)	Gurley	1893 : 417
1894	<i>Myxobolus cf. linearis</i>	Gurley	1894 : 253-254
1899	<i>Henneguya linearis</i> var.	Labbé	1899 : 103

Habitat: Base of spines of the second dorsal fin of *Ameiurus melas* Raf.; Iowa (Storm Lake) (August).

Vegetative form: Spherical cysts, 1mm. in diameter.

Spore: Lanceolate. Dimensions: length of the body 19μ , width 5 to 6μ , thickness about 3μ .

Remarks: The species is most probably different from *Henneguya linearis* judging from the difference in the form and structure of spores, the seat of infection, and host species. Hence, it is recorded here as an independent species.

HENNEGUYA STRONGYLURA (Gurley) Labbé

[Fig. 506]

1841		Müller	1841 : 480
1894	<i>Myxobolus strongylurus</i>	Gurley	1894 : 249
1899	<i>Henneguya strongylura</i>	Labbé	1899 : 103

Habitat: Skin of cephalic region of *Synodontis schall* Bl. Schn.; Nile.

Vegetative form: Cysts over 2.18mm. in diameter.

Spore: Dimensions: length of the body 9μ , breadth 5.4μ . Tail always undivided. Two polar capsules of equal size.

HENNEGUYA MONURA (Gurley) Labbé

[Fig. 507]

1880		Ryder	1880 : 211-212
1893	<i>Myxobolus monurus</i>	Gurley	1893 : 416
1894	<i>Myxobolus monurus</i>	Gurley	1894 : 249-250
1899	<i>Henneguya monura</i>	Labbé	1899 : 103

Habitat: Subcutaneous intermuscular tissue of *Aphredoderus sayanus* Gill.; New Jersey (Woodbury).

Vegetative form: Cysts, lenticular, large, white, opaque and numerous (20). Membrane thin.

Spore: Lenticular or slightly obovate. Tail 2 to 3-times longer than the body.

HENNEGUYA KOLESNIKОВI (Gurley) Labbé

[Fig. 508]

1886		Kolesnikov	1886 : 242-248
1894	<i>Myxobolus kolesnikovi</i>	Gurley	1894 : 256-257
1898	<i>Myxobolus bicaudatus</i> (part)	Zschokke	1898 : 602-604, 646-655, 699-703
1899	<i>Henneguya kolesnikovi</i>	Labbé	1899 : 103-104

Habitat: Interstitial connective tissue of the thoracic and intercostal muscles of *Coregonus lavaretus* L.; Russia.

Vegetative form: Cysts numerous (80), spherical or oval; length 10 to 30mm., breadth 7 to 20mm.

Spore: Oval with a pointed anterior end. Tail three times longer than the body.

Remarks: Zschokke thinks the present species is identical with *Henneguya zschokkei*. But the evidence is not clear enough to bring one to agree with him due to the incomplete description of the present species.

HENNEGUYA MACRURA (Gurley) Thélohan

[Figs. 509 to 512]

1893		Evermann	1893 : 76
1894	<i>Myxobolus macrurus</i>	Gurley	1894 : 250-253
1895	<i>Henneguya macrura</i>	Thélohan	1895 : 354

Habitat: Subcutaneous connective tissue of head of *Hybognathus nuchalis* Ag.; Neches River, Texas (November, temperature of water 9°.4C.) Of frequent occurrence.

Vegetative form: Cysts, elongated 6mm. by 2mm. or less.

Spore: Rounded oblong. Dimensions: length 10 to 11 μ , breadth 6 to 8 μ , thickness 4 μ . Shell-valves unequally convex. Tail 30 to 40 μ .

HENNEGUYA ZSCHOKKEI (Gurley) Doflein

[Fig. 513]

1884		Zschokke	1884 : 234-235
1894	<i>Myxobolus</i> (?) <i>zschokkei</i>	Gurley	1894 : 244
1898	<i>Myxobolus bicaudatus</i> (part)	Zschokke	1898 : 602-607, 646-655, 699-703
1898	<i>Myxobolus bicaudatus</i>	Zschokke	1898a : 213-214
1901	<i>Henneguya zschokkei</i>	Doflein	1901 : 202
1904	<i>Henneguya zschokkei</i>	Hofer	1904 : 56
1905	<i>Henneguya zschokkei</i>	Nufer	1905 : 77, 185

Habitat: Subcutaneous and superficial intermuscular tissue of *Coregonus fera*, *C. schinzii* Fatio, *C. hiemalis* Jur. and muscular tissue and branchia of *C. wartmanni nobilis* and *C. exiguus albellus*; Neuchâtel-see, Zurich See, Genfer-see, Thuner-see, Vierwaldstätter-see.

Vegetative form: Zschokke writes as follows: Cysts rounded or oval surrounded by a compact membrane with many nuclei. The largest 32mm. by 16mm. Protoplasm granular. Polysporous.

Spore: Rounded oval in front view; broad elliptical in side view. Anterior end rounded; posterior end tapering, forming tail. Sutural ridge distinct. Tail is either bifurcated along the entire length or a single form, no intermediate form being observed. Dimensions: total length 55 μ , length of the body 10 μ , breadth 7 μ , length of tail 4 to 5 times the length of the spore-body, length of polar filament 6 to 10 times that of the body of the spore.

Remarks: Zschokke thinks that *Henneguya kolesnikovii*, *H. zschokkei* and *H. sp.* Gurley are one and the same species, for which he proposed the name *Myxobolus bicaudatus*.

HENNEGUYA sp. (Gurley) Labbé

[Fig. 514]

1886		Benecke	1886 : 211
1894	<i>Myxobolus sp. inc.</i>	Gurley	1894 : 244
1899	<i>Henneguya sp.</i>	Labbé	1899 : 104
1904	<i>Henneguya sp.</i>	Hofer	1904 : 51

Habitat: Integument (?) of *Leuciscus rutilus* L. The parasites formed boil-like enlargement in the skin.

Vegetative form: Not described.

Spore: Not described.

HENNEGUYA sp. (Gurley) Labbé

1874		Claparède	1874 : 114
1894	<i>Myxobolus sp. inc.</i>	Gurley	1894 : 253
1898	<i>Myxobolus bicaudatus</i> (part)	Zschokke	1898 : 602-607, 646-655, 699-703
1899	<i>Henneguya sp.</i>	Labbé	1899 : 104

Habitat: Branchial-arches of *Coregonus fera*; Genfer-see.

Vegetative form: One cyst, 1mm. in diameter.

Spore: Tail short. Zschokke quotes: length 8 to 10 μ .

Remarks: According to Zschokke, this species is identical with *H. zschokkei*.

HENNEGUYA TENUIS Vaney et Conte

[Fig. 515]

1901 *Henneguya tenuis* Vaney and Conte 1901 : 103-106

Habitat: Connective tissue of alimentary tract of *Acerina cernua* L.; Lyon (February).

Vegetative form: Numerous cysts particularly in the pyloric coecum. Usually spherical. Size: 30 to 150 μ in diameter.

Spore: Oval and small. Tail short. Two polar capsules at the anterior end. Sporoplasm with a nucleus, rod-shaped, with somewhat enlarged ends which is located at right angles to the longitudinal axis. Iodinophilous vacuole could not be traced. Dimensions: length 4 μ , breadth 2 μ .

HENNEGUYA NÜSSLINI Schuberg et Schröder

[Figs. 516 and 517]

1905 *Henneguya nüsslini* Schuberg and Schröder 1905 : 56-59

Habitat: Subcutaneous connective tissue at the base of dorsal fin of *Trutta fario* L.; Gutach.

Vegetative form: Trophozoites form cysts (2 cysts found). Cysts lenticular, 1.5 to 2mm., surrounded by many concentric layers of fibrous connective tissue. Cysts containing only mature spores.

Spore: Broad oval form, flattened. Anterior end rounded. Tail at the posterior end. Shell somewhat thick, often shows sutural ridge. Tail filaments two. A "dark part" which in side-view is of triangular form, runs into the tail. Sporoplasm, occupying the posterior half of the spore, projects a narrow portion between the polar capsules beyond the middle of the capsules. Sporoplasm, uniformly granular, contains an iodophilous vacuole and one, sometimes two nuclei connected by nuclear bridge. Polar capsules, pyriform, opening independently. Coiled polar filament observable, coiled 6 to 7 times. Dimensions: length excluding tail 12 μ , length with tail 32 μ , breadth 8 to 9 μ , polar capsules 5 μ by 3 μ , length of polar filament 4 to 5 times longer than that of spore excluding tail (48 to 60 μ).

HENNEGUYA LÉGERI Cépède

[Figs. 518 to 523]

1905	<i>Henneguya légeri</i>	Cépède	1905 : 905-913
1906	<i>Henneguya légeri</i>	Cépède	1906 : 66
1913	<i>Henneguya légeri</i>	Cépède	1913 : 302-305

Habitat: Urinary bladder of *Cobitis barbatula* L.; Isère (January).

Vegetative form: Young trophozoites subcircular, irregularly elliptical or elongated with distinct differentiation of protoplasm into ectoplasm and endoplasm. Plasmotomic multiplication takes place during winter months, when no spore is formed.

Spore: Oval with short tail, mostly bifurcated at the free end. The anterior end is more rounded, occasionally acuminated. Two polar capsules of equal size. Coiled polar filament distinct in vivo. Sporoplasm granular, contains two nuclei and a vacuole. The spore often shrinks in fresh conditions, probably owing to the poorly developed thin valves. Dimensions of spores mounted in balsam: length variable. Examples: Total length 22.5μ , tail 8.5μ ; total length 19.5μ , tail 8μ ; length of main part 8.5μ , breadth (comparatively constant) 6μ .

HENNEGUYA ACERINAE Schröder

[Figs. 525 and 526]

1906	<i>Henneguya acerinae</i>	Schröder	1906 : 186-196
1910	<i>Henneguya creplini</i>	Wegener	1910 : 84
1911	<i>Henneguya acerinae</i>	Nemeczek	1911 : 155

Habitat: Branchiae of *Acerina cernua* L., *Aspro zingel* Cuv., *Lucioperca lucioperca* L. and *L. sandra* Cuv. (?); Heidelberg (Necker), Apatin, Komitat Baco-Bodrog, Hungary (May).

Vegetative form: Schröder describes as follows: Rounded or spherical cysts in the connective tissue of branchial lamella. Full-grown cysts up to 300μ in diameter. Protoplasm is differentiated into ectoplasm and endoplasm. Ectoplasm shows fine radial striations. Endoplasm granular, contains many nuclei, especially lying in the middle portion. Well developed cyst, containing only spores, is surrounded by a membrane. On the surface of the ectoplasm, numerous edge-like elevations, branched and joining together, were recognized. Polysporous.

Nemeczek observed the largest cyst, spherical and 600μ in diameter.

Spore: Pyriform in front view; flattened. The anterior end is more or less blunt. Shell uniformly thin. Sutural edge slightly enlarged. Sporoplasm finely granular, contains an iodophilous vacuole and two nuclei. Polar capsules approximated closely, each having an independent opening. Dimensions: length 20 to 22μ , breadth 8 to 9μ , thickness 6 to 7μ , length of tail 50 to 60μ , polar capsules 10μ by 2 to 3μ , length of polar filament 80 to 90μ (water and nitric acid).

Nemeczek's form is as follows:

The tail is bifurcated along its entire length. In one case (May, 1909), however, all the spores had no bifurcated tail, while the polar capsules were of unequal size. Dimensions in fresh state: total length 37.6 to 41.8μ , length, excluding tail 12.6 to 16.8μ , breadth 4.5μ , length of polar capsule 6.3 to 8.4μ , length of polar filament 67μ , length of tail 25μ .

Nemeczek observed two more different (?) forms. One form found in *Lucioperca sandra*, tho the size differs from the dimensions given by Schröder, is thought to be identical with the present species. Another form in the branchiae of *Aspro zingel*, which is also to be one and the same species with the present species has the following dimensions: total length 35μ , length of spore excluding tail 15μ , breadth 5μ , length of polar capsule 6μ , length of tail 20μ .

HENNEGUYA GIGANTEA Nemeczek

[Figs. 527 to 535]

1911	<i>Henneguya gigantea</i>	Nemeczek	1911 : 146-154
1914	<i>Henneguya gigantea</i>	Georgévitch	1914 : 387-409

Habitat: Branchiae of *Lucioperca sandra* Cuv.; Apatin, Komitat Bacs-Bodrog, Hungary, Petrograd. Nemeczek mentions that the infection takes place only among young fish.

Vegetative form: Cysts numerous and of conspicuous size in the free end of branchial lamella. In average, each gill-arch has about 100 cysts which are of creamy color. Young cysts 400 to 450μ in diameter. They gradually begin to increase the size, from autumn until toward the end of spring, during which period, the contents remaining in the stages of pansporoblasts formation. Older cysts rounded spindle shape with the length of 4 to 7mm. and the breadth of 2 to 3mm. The connective tissue and epithelial cell layers form the cyst membrane. The connective tissue either simply surrounds the parasite or branches in the surface of the parasite, increasing in thickness and forming more or less enclosed chambers of the parasite. The membrane of the cyst which contains mature spores is usually very thin. Throughout the growth of the cyst, "chromatoid body" is seen in the endoplasm, which appears first as a filiform structure, stained deeply with nuclear stain. Later they gather together and form a compact body, situated excentrically. Fine branches from it become directed toward the surface of the body, anastomosing each other so that a network is formed on the surface of the cyst. The latter develops small ovoidal or columnal bodies (1.2μ long and about 1μ wide), which are arranged radially and densely. The number and quantity of these bodies increase in proportion to the number of propagative nuclei and they begin to disappear, first in the central portion, then in the periphery, so that in fully grown cysts (in summer months) these chromatoidal bodies are more rudimentary. Differentiated protoplasm is only recognized in young individuals, in which case ectoplasm is homogeneous and endoplasm reticular. Polysporous.

Spore: Nemeczek gives the following accounts.

Spindle shape, with truncate anterior end and very long thread like tail at the posterior end. The tail seems split into two at about the middle part of its length. Gentian violet stains the tail so intensively that its entire length could easily be made out. Dimensions: total length 87.5 to 110.5 μ , length of the body 10.5 μ , breadth 5 μ , length of tail 77 to 100 μ , length of polar capsule 5 μ , length of polar filament 70 μ (pressure or dessication followed by immersion in water).

Georgévitch's form: length excluding tail 15 μ , breadth 6 μ , length of tail 75 μ , length of polar capsule 6 μ , length of polar filament 75 μ , diameter of the iodophilous vacuole 4 μ .

Remarks: Nemeček mentions that from October on, cysts had no spores, only containing propagative cells. The velocity of the development of spores depends upon the temperature of water.

Georgévitch worked out the spore formation of the species and observed that the binucleated sporeplasm emerged from the posterior end of the spore.

HENNEGUYA (?) sp. Nemeček

[Figs. 536 to 539]

1911 *Henneguya* sp. Nemeček 1911 : 157-159

Habitat: Branchiae of *Abramis brama*; Komorn, Komitat Komorn, Hungary (March).

Vegetative form: Cysts in the branchiae.

Spore: Besides normal spores of *Myxobolus rotundus* (page 149), spores of *Henneguya* type in small number were found. The anterior part of these spores resembles that of the species mentioned above, while the breadth is much smaller (8 μ) than the latter. Majority of spores have a thread like tail, 10 to 15 μ long, which was often bifurcated. An iodophilous vacuole was fairly marked.

Remarks: It is placed here as a species of *Henneguya* by reason of the bifurcate tail.

HENNEGUYA GASTEROSTEI Parisi

[Figs. 540 to 543]

1912 *Henneguya gasterostei* Parisi 1912 : 296-297

Habitat: Kidney of *Gasterosteus aculeatus* L.; Lago di Garda (February).

Vegetative form: Rounded or oval, usually with two, but rarely with four spores. Ectoplasm thin and hyaline. Endoplasm contains numerous granules, most probably of fatty nature and decreasing in number as spores grow. Free full-grown spores were seen abundantly in the connective tissue of renal tubules, glomeruli, etc. Disporous and polysporous.

Spore: Oval with slightly attenuated anterior end; posterior end tapering into tails, which end in one point or bifurcated; asymmetrical in shape, one valve is more curved than the other. This asymmetry of the shell-valves in profile enables the present species to be distinguished from other species. Shell striated longitudinally. Two polar capsules pyriform and well developed, reaching to the middle of the spore. Sporoplasm with a round iodophilous vacuole. Dimensions: total length 38 to 48 μ , length of the cavity of the spore 15 μ , breadth 6 to 7.5 μ , polar capsules 7.5 to 9 μ by 3 to 3.5 μ , length of polar filament 50 μ .

HENNEGUYA NEAPOLITANA Parisi

[Figs. 544 and 545]

1912 *Henneguya neapolitana* Parisi 1912 : 297-298

Habitat: Connective tissue of the renal tubule of kidney of *Box salpa* C. et V.; Napoli (August).

Vegetative form: Small cyst (40 to 50 μ in diameter) surrounded by thin membrane, containing a number of spores, numerous pigment granules and coarse yellowish globules.

Spore: Oval, slightly flattened. Anterior end rounded when seen from the front, but attenuated in profile. Shell tapering into a long fine tail posteriorly. The fine distal portion of the tail wraps around the thicker part. Two polar capsules, pyriform, occupying the anterior half of the cavity of the spore, cross each other when seen from the front. Sporoplasm finely granular with two nuclei, the iodophilous vacuole being hardly visible. Dimensions: total length 50 to 60 μ , length of the cavity of spore 8.5 to 9.5 μ , breadth 8.5 to 9.5 μ , internal breadth 6.3 to 7 μ , thickness 8 μ , polar capsules 4.7 to 5.5 μ by 3 μ .

HENNEGUYA WISCONSINENSIS Mavor et Strasser

[Figs. 558 and 559]

1916 *Henneguya wisconsinensis* Mavor et Strasser 1916 : 676-682

Habitat: Urinary bladder of *Perca flavescens*; Lake Mendota, Wisconsin (April).

Vegetative form: Trophozoites are usually elongated and have the general form and shape of a limax ameba. It may reach a size of 300 μ by 70 μ . Clear differentiation of ectoplasm and endoplasm. Pseudopodia lobose. Two spores are formed in each pansporoblast. Polysporous.

Spore: Ovoid, bilaterally symmetrical, and have a bifurcated caudal process. Two polar capsules at anterior end. Coiled polar filament visible *in vivo* (5 windings). Dimensions: length excluding tail 11.5 μ , breadth 7 μ , tail 9.6 μ , polar capsules 3.5 μ by 2.5 μ , length of filament 33 μ .

HENNEGUYA BRACHYURA Ward

[Figs. 650 to 653]

1919 *Henneguya brachyura* Ward 1919 : 57

Habitat: In the cartilaginous fin ray of the caudal fin of *Notropis anogenus*; Put-in-Bay, Lake Erie (August). The species was found encysted in the same fish which was heavily infected by *Myxobolus aureatus*.

Vegetative form: Cysts rounded with slightly irregular contour imbedded in the fin ray. The size varies from 160μ in diameter up to 360μ by 240μ . No particular cyst membrane could be recognized. The differentiation of the protoplasm into ectoplasm and endoplasm is distinct. The ectoplasm covering the entire surface of the parasite as a layer 4 to 6μ thick, shows structure of a very finely granular nature. The endoplasm coarsely alveolar, is filled with mature spores in the central portion, while numerous nuclei and young spores in various developmental stages are present at the peripheral portion. Polysporous.

Spore: Rounded oval in front view; spindle shape with symmetrically built valves in profile. Shell rather thick. Sutural ridge fairly well marked; sutural edge exhibiting a variable number of folds (8 to 10). Two pyriform polar capsules are usually of the same size and form. The tail is a single process, usually more or less bent or irregularly curved, very rarely being straight. In general, it is sinuous with two or three shallow curves and is rather short, tapering gradually to a point. In young spores which are less deeply stained by any stain, various developmental stages of the tail are readily recognized. Giemsa solution stains the shell proper in clear blue, while the tail takes on a beautiful pink color, a distinct difference in affinity for dyes between the material in the tail and the shell. It seems probable that the tail of this type is entirely different in its development from that of the ordinary bifurcated type. Dimensions in section: length 10 to 11.5μ , breadth 8 to 8.75μ , thickness 4 to 5μ , polar capsules 3 to 4μ by 2μ , length of the tail up to 17μ .

HENNEGUYA SALMINICOLA Ward

[Figs. 654 to 656]

1914 ?*Henneguya zschokkei* Zschokke and Heitz 1914 : 200-201
 1919 *Henneguya salminicola* Ward 1919 : 59

Habitat: In the sub-dermal tissue of *Onchorhynchus keta* and *O. kisutch* (Zschokke and Heitz, Kamtschatka) and in the connective tissue in body muscles of *Oncorhynchus keta*, Stickeen River, Alaska (Ward, September). The last named author undertook a careful examination of a part of the infected tissue preserved in formol. The species forms conspicuous cysts in the muscle from the sub-peritoneal to the sub-dermal connective tissue, tho all are sub-peritoneal in position.

Vegetative form: Ward describes as follows: The whitish opaque cysts are pyriform, and fairly uniform in size (3 to 6mm. in diameter). The cyst is covered by numerous layers of connective tissue which form a tough membrane around the parasite. The cyst contains young spores in various stages of development, which showed that two spores are formed in one pansporoblast, and mature spores thickly massed together in the central area. Polysporous.

Spore: Oval with rounded anterior and more or less attenuated posterior ends; elliptical in profile with attenuated anterior end. Shell smooth. Sutural edge exhibits folds variable in number (usually 6 to 7). Tail double, composed of two fine and equal halves which are the prolongation of the shell valves. The processes usually run roughly parallel to each other. Two pyriform polar capsules are of slightly different dimensions. Coiled polar filament is indistinct in preserved unstained specimens. Sporoplasm finely granular, shows a large iodophilous vacuole. Dimensions of stained and mounted spores: total length 47μ (42.75 to 52.44μ), length of the main part 12μ (11.97 to 14.25μ), breadth 8μ (7.12 to 8.43μ), thickness 4.78μ , length of tail 35μ (30.78 to 38.19μ), polar capsule 3.70 to 4.55μ by 1.59 to 2.85μ .

Remarks: Zschokke and Heitz (1914) observed a species from Kamtschatka, which they thought to be identical with *Henneguya zschokkei* (page 165). The writer is inclined to think that the species is identical with the species just described from Alaska.

HENNEGUYA MIYAIRII nov. spec.

[Fig. 524]

1909 *Henneguya* sp.

Miyairi

1909 : 127-129

Habitat: Subcutaneous tissue of head of *Carassius auratus* L.; Fukuoka (Nippon).

Vegetative form: Trophozoites form cysts and are also found in the condition of diffuse infiltration around the cysts. Cyst-membrane fibrous and thin. Ectoplasm and endoplasm fairly well differentiated, though the border line is not sharply marked. At the periphery of endoplasm, pansporoblasts with 7 to 12 nuclei are present (Two spores are formed in each pansporoblast?). Polysporous.

Spore: Oval, with broadly rounded anterior and slightly elongated posterior ends, the latter ending in long and fine tails. Two polar capsules at the anterior portion, are pyriform, small and convergent. Sporoplasm with an iodophilous vacuole. Dimensions: length 12μ , breadth 8μ , length of the tails 10 to 30μ , length of polar filaments 23 to 40μ .

Remarks: As the description gives the details by which the species can be distinguished from other species, the writer establishes it on an independent basis.

HENNEGUYA MICTOSPORA nov. spec.

[Figs. 546 to 557]

Habitat: Urinary bladder of *Lepomis cyanellus* Raf., *L. humilis* Gir. and *Micropterus salmoides* Lac.; Stony Creek, Ill. (November).

In one out of three (6.5 to 8cm. long) of the first, in one out of two (7 and 9.5cm. long) of the second and in one of the third species, examined in the middle of November, was found the present form. None showed a heavy infection, a number of scattered trophozoites and spores being observed. The host did not show any pathological change.

Vegetative form: Polymorphous. Generally rounded or elongated oval. In small monosporous and disporous forms, the tail of the spores developed inside, is extruded from the body, so that these trophozoites show long processes (Figs. 546, 553, 555). Pseudopodia lobose, and extruded from the entire surface of the body (Fig. 547), tho sometimes they are well formed at one end of the body. Protoplasm is differentiated distinctly into ectoplasm and endoplasm. Ectoplasm is homogeneous and hyaline, forming the outer layer. Endoplasm is of reticular structure. The body is colorless, often yellowish, when the endoplasm is loaded with numerous yellowish coarse granules. The size varies from 6 or 7 μ up to 60 μ . In a rounded form of 38 μ in longest diameter, five pansporoblasts, each developing two spores and many nuclei were observed. In another oval form of 45 μ by 60 μ in size, numerous nuclei were stained, showing that no development of pansporoblast has yet taken place. Disporous, polysporous and monosporous, tho of rare occurrence.

Spore: Broad spindle shape with attenuated anterior end. Shell rather thin. Each valve has 6 to 8 longitudinal striations on the surface. A long tail composed of two halves, is developed at the posterior end. Two pyriform polar capsules with distinctly visible coiled polar filament opens at the anterior tip. Sporoplasm, finely granular, contains an iodophilous vacuole which is made distinctly visible by treating with Lugol's solution. When stained two typical nuclei are recognized in the sporoplasm. Dimensions of the fresh spores: length excluding tail 13.5 to 15 μ , breadth 8 to 9 μ , thickness 6 to 7.5 μ , length of tail 30 to 35 μ , often up to 40 μ , polar capsule 5 to 6 μ by 3 μ , length of polar filament 40 μ .

Genus HOFERELLUS Berg

1898	<i>Hoferellus</i>	Berg	1898 : 41
1898	<i>Hoferia</i>	Doflein	1898 : 288-289

The characters of the genus are described on page 59.

Type and only species: *Hoferellus cyprini* Doflein.

HOFERELLUS CYPRINI Doflein

[Figs. 577 to 581]

1898	<i>Hoferia cyprini</i>	Doflein	1898 : 289-290
1908	<i>Hoferellus cyprini</i>	Mercier	1908 : LIII-LIV
1910	<i>Hoferellus cyprini</i>	Plehn	1910 : 20-22

Habitat: In lumen and epithelial cells of renal tubules of kidney of *Cyprinus carpio* L.; France and Germany.

Vegetative form: Young trophozoites live in epithelium. Adults free in the urinary tubules. Form rounded or oval. No clear differentiation of protoplasm. Pseudopodium unobserved. Endoplasm contains numerous granules and many nuclei. Each pansporoblast forms two spores. Smaller individuals 20 to 30 μ in diameter. Polysporous.

Spore: Pyramidal with two short tail-like processes at the posterior end, which are formed from the shell-valves like those of *Henneguya*. Between these two processes, rarely small protoplasmic pointed processes occur. Each shell-valve has 9 to 10 longitudinal striations on it. Two polar capsules at the anterior part, show clearly the coiled polar filaments. Sporoplasm has two nuclei and an iodophilous vacuole. Dimensions: total length 10 to 12 μ , breadth 8 μ , tail-process 2 μ long, polar capsule 3 μ long.

MYXOSPORIDIA GENERA ET SPECIES INCERTAE

Gen. et spec. incert. Leydig

1851		Leydig	1851 : 222
1894	Gen. et spec. incert.	Gurley	1894 : 186

Habitat: Cysts in the root of tongue of *Chondrostoma nasus* L.; Germany.

Gen. et spec. incert. Leydig

1851		Leydig	1851 : 222
1894	Gen. et spec. incert.	Gurley	1894 : 186

Habitat: Heart (auriculo-ventricular valve) of *Leuciscus rutilus* L.

Gen. et spec. incert. Leydig

1851		Leydig	1851 : 223
1894	Gen. et spec. incert.	Gurley	1894 : 186

Gen. et spec. incert. Heckel et Kner

1851		Heckel and Kner	1851 : 12
1894	Gen. et spec. incert.	Gurley	1894 : 186-187

Habitat: Branchiae of *Lucioperca lucioperca* L.; Austria.

Gen. et spec. incert. Borne

1886		Borne	1886 : 211
1894	Gen. et spec. incert.	Gurley	1894 : 187

Habitat: *Scomber scombrus* L.

Genus incert. MERLUCII Perugia

[Figs. 582 and 583]

1891	<i>Myxosporidium merlucii</i>	Perugia	1891 : 22-24
1894	<i>Myxobolus ? merlucii</i>	Gurley	1894 : 242-243
1899	<i>Myxobolus merlucii</i>	Labbé	1899 : 100

Habitat: Gall-bladder of *Merluccius merluccius* L.; Italy.

Vegetative form: Various form. No differentiation of protoplasm. Disporous (?).

Spore: Oval, with two polar capsules.

Remarks: The species was placed in the genus *Myxobolus* by previous authors. The figures given by Perugia show that the spores are at least dimorphous. From the habitat and the disporous characters, one should place it rather in one of the genera of the Family Ceratomyxidae.

Genus incert. CONGRI Perugia

[Figs. 584 and 585]

1891	<i>Myxosporidium congri</i>	Perugia	1891 : 24-25
1894	Genus incert. <i>congri</i>	Gurley	1894 : 182
1912	<i>Myxobolus congri</i>	Parisi	1912 : 284

Habitat: Gall-bladder of *Conger conger* L.; Genova.

Vegetative form: Floating in the bladder. Form variable. Movements incessant, slow and ameboid.

Spore: Not described.

Gen. et spec. incert. Linton

[Fig. 590]

1891		Linton	1891 : 359-361
1894	Gen. et spec. incert.	Gurley	1894 : 182-183

Habitat: Subcutaneous tissue of *Notropis megalops* Raf.; Ohio (Black River; September, October).

Vegetative form: Cysts. Globular, discrete or aggregated into clusters, white, with minute patches of black pigment from host; size varying from 2.5mm. (single cyst) to 7mm. by 5mm. (clusters); cyst-membrane composed of connective tissue.

Spore: Top-shaped, somewhat flattened; with pointed anterior and broadly rounded posterior end. Shell thick, with elevated sutural ridge. Polar capsules could not be detected. Protoplasm finely granular. Dimensions: length 17μ , breadth 10μ , thickness 6μ .

Remarks: The cysts and figures of spores given by Linton suggest that it is most probably a unicapsular *Myxobolus*. As Linton could not detect (?) the polar capsule, tho his figures faintly show the said structure, it is placed in this group.

Gen. et spec. incert. Mingazzini

1892		Mingazzini	1892 : 398
1899		Labbé	1899 : 113

Habitat: Ovarian egg of *Lacerta* sp.

Vegetative form: Ameboid with hyaline pseudopodia and granular protoplasm.

Spore: Not observed.

Gen. et spec. incert. Nufer

1905	<i>Myxobolus</i> sp.	Nufer	1905 : 71, 77, 79, 85, 186
------	----------------------	-------	----------------------------

Habitat: In the connective tissue of branchia of *Chondrostoma nasus*; Lake of Lucerne. A single cyst in a single host fish.

Vegetative form: Cyst white, and of 1mm. in diameter.

Spore: With two polar capsules at one pole and the sporoplasm. Dimensions or any other characters are not given.

Remarks: Altho Nufer placed the form in the genus *Myxobolus*, this must be brought into the present group in view of the fact that the iodophilous vacuole was not detected, and that the observation is too incomplete to place it to any one of the genera.

Gen. et spec. incert. Mavor

[Figs. 586 and 587]

1915		Mavor	1915 : 27-28, 32-33
1916		Mavor	1916 : 553-554

Habitat: Gall-bladder of *Urophycis chuss*; St. Andrews (July to September).

Vegetative form: Mavor writes as follows:

Attached, usually in large numbers, to the epithelium of the bladder, occurs a spherical or ellipsoidal trophozoite which in stained preparations is found to contain numerous nuclei. Very often clusters of *Ceratomyxa acadiensis* are found adhering to the free surface of myxosporidium. In fresh preparations the appearance is that of budding from a parent organism. An examination of sections has shown a sharp division between the myxosporidium and *Ceratomyxa acadiensis*.

Spore: Not found.

Remarks: Mavor supposed that the form under discussion probably was some species of *Myxidium* or *Chloromyxum*.

Gen. et spec. incert. Mavor

[Figs. 588 and 589]

1916		Mavor	1916a : 68-69
------	--	-------	---------------

Habitat: Urinary bladder of *Stizostedion vitreum* Mitch.; Georgian Bay (Canada).

Vegetative form: Free forms vary greatly in shape, being rounded, elongated or branched. The largest individual 200 μ . Ectoplasm layer clearly visible, sometimes projecting many bristle-like short processes. Endoplasm contains greenish granules. Trophozoites also attached to the epithelium by means of deeply stainable portion of the body.

Spore: Not observed.

Remarks: Mavor mentions resemblance of the present form to *Myxidium lieberkühni* Bütschli in many respects.

KEYS TO THE GENERA AND SPECIES OF MYXOSPORIDIA

No key to the genera and species of Myxosporidia has been published up to the present time. This is due of course to the difficulties which accompany such an attempt. These difficulties lie chiefly in the incompleteness of the observations and descriptions of the majority of the species of Myxosporidia.

The writer has attempted in the following pages to carry out this task. The key is by no means complete, as is unavoidable in the present state of knowledge concerning this particular group of the Protozoa.

Altho the spore is the fundamental factor used in constructing this key, it was necessary to refer also to some other secondary characters such as vegetative form and habitat.

Some authors are inclined to think that the difference in host species gives an ample basis on which to record the parasite as a new species. In some cases the parasite is specific in a certain host species while in other cases a number of different host species are infected by one and the same parasite. Therefore one can not lay much emphasis upon a difference of hosts in fixing the identification of a Myxosporidian.

KEY TO THE GENERA OF MYXOSPORIDIA

- | | | |
|--------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| 1(6) | Spore approximately spherical
Suborder Sphaerosporea Kudo 1919..... | 2 |
| 2(3) | Spore with four polar capsules
Family CHLOROMYXIDAE Thélohan 1890
Genus <i>Chloromyxum</i> Mingazzini 1890..... | (183) |
| 3(2) | Spore with two polar capsules
Family SPHAEROSPORIDAE Davis 1917..... | 4 |
| 4(5) | Sutural line of spore straight
Genus <i>Sphaerospora</i> Thélohan 1892..... | (185) |
| 5(4) | Sutural line of spore sinuous
Genus <i>Sinuolinea</i> Davis 1917..... | (186) |
| 6(1) | Spore not spherical..... | 7 |
| 7(16) | Largest diameter of spore at right angles to sutural line; two polar capsules, one on each side of sutural line
Suborder Eurysporea Kudo 1919
Family CERATOMYXIDAE Doflein 1899..... | 8 |
| 8(11) | Shell-valves prolonged laterally..... | 9 |
| 9(10) | Shell-valves hemispherical or rounded
Genus <i>Leptotheca</i> Thélohan 1895..... | (179) |
| 10(9) | Shell-valves conical; free end tapering to a more or less pointed end
Genus <i>Ceratomyxa</i> Thélohan 1892..... | (180) |
| 11(8) | Shell-valves rather elongated; circular in cross-section..... | 12 |
| 12(13) | Spore rounded oblong; shell longitudinally striated; polar capsules pyriform, with or without long and fine posterior filaments
Genus <i>Mitraspora</i> Fujita 1912 emend. Kudo 1919..... | (183) |

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

- 13(12) Spore angular, not rounded 14
- 14(15) Spore pyramidal in front view; with its base at anterior end; with or without distinct anterior processes; shell smooth
Genus *Myxoproteus* Doflein 1898 (183)
- 15(14) Spore isosceles triangular in front view; anterior end attenuated; polar capsules spherical and large; shell with fine network-like ridges; with posterior fringe-like processes
Genus *Wardia* Kudo 1919 (183)
- 16(7) Largest diameter of spore coincides with or at an acute angle to sutural plane; one or two polar capsules which are in sutural plane
Suborder **Platysporea** Kudo 1919 17
- 17(22) Spore fusiform; two polar capsules, one at each end of spore
Family MYXIDIIDAE Thélohan 1892 18
- 18(21) Spore more or less regularly fusiform; shell-valves symmetrical 19
- 19(20) Polar filament fine and long
Genus *Myxidium* Bütschli 1882 (186)
- 20(19) Polar filament thick and short
Genus *Sphaeromyxa* Thélohan 1892 (188)
- 21(18) Spore semi-circular in front view; polar filament fine
Genus *Zschokkella* Auerbach 1910 (188)
- 22(17) Spore not fusiform; with one or two polar capsules at anterior extremity 23
- 23(26) Sporoplasm without iodophilous vacuole
Family MYXOSOMATIDAE Poche 1913 24
- 24(25) Spore elongated ovoid in front view; anterior end mostly pointed
Genus *Myxosoma* Thélohan 1892 (189)
- 25(24) Spore more or less rounded in front view
Genus *Lentospora* Plehn 1905 (189)
- 26(23) Sporoplasm always with iodophilous vacuole
Family MYXOBOLIDAE Thélohan 1892 27
- 27(30) Spore with posterior process; shell sometimes striated 28
- 28(29) Process more or less long, projecting posteriorly along median line of spore; process either single or double; shell sometimes striated
Genus *Henneguya* Thélohan 1892 (193)
- 29(28) Process short projecting posteriorly from sides; shell longitudinally striated
Genus *Hoferellus* Berg 1898 (173)
- 30(27) Spore without posterior process; shell unstriated; one or two polar capsules
Genus *Myxobolus* Bütschli 1882 (189)

II. KEY TO THE SPECIES

Genus LEPTOTHECA Thélohan 1895

- 1(14) Spore: sutural diameter always more than half of greatest breadth 2
- 2(7) Average sutural diameter less than 10μ 3
- 3(4) Posterior margin of spore concave in front view; sutural diameter 8 to 9μ , breadth 12 to 14μ . Trophozoite usually with a long process
Leptotheca longipes Auerbach 1910 (63)
- 4(3) Posterior margin of spore not concave 5
- 5(6) Posterior margin of spore flattened; polar capsules pyriform; sutural diameter 6 to 7μ , breadth 11 to 12μ . Trophozoite with actively motile long filiform pseudopodia at rounded end
Leptotheca agilis Thélohan 1892 (60)

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

- 6(5) Spore regularly ovoidal; polar capsules short pyriform, opening on opposite sides; sutural diameter 9μ , breadth 16μ . Trophozoite pyriform without any recognizable pseudopodium
Leptotheca fusiformis Davis 1917.....(63)
- 7(2) Average sutural diameter of spore equal to or larger than 10μ 8
- 8(13) Shell-valves symmetrically built; sutural ridge straight..... 9
- 9(10) Posterior margin of spore concave in front view; sutural diameter 10μ , breadth 18 to 20μ ; each spore formed independently
Leptotheca infirmis Auerbach 1910.....(63)
- 10(9) Spore regularly ovoidal..... 11
- 11(12) Trophozoite extremely polymorphous. Spore: sutural diameter 10 to 12μ , breadth 18 to 20μ
Leptotheca polymorpha Th  lohan 1895.....(61)
- 12(11) Typical form of trophozoite elongated; anterior end depressed surrounded by short often branched pseudopodia. Spore: sutural diameter 12 to 15μ , breadth 18 to 20μ
Leptotheca elongata Th  lohan 1895.....(60)
- 13(8) Shell-valves asymmetrically built; sutural ridge sinuous; sutural diameter 9 to 10μ , breadth 16 to 18μ . Trophozoite rounded; movements slow
Leptotheca lobosa Davis 1917.....(64)
- 14(1) Sutural diameter equal to or less than half of greatest breadth..... 15
- 15(18) Average sutural diameter smaller than 10μ 16
- 16(17) Spore arch-shaped in front view; polar capsules pyriform; sutural diameter 3 to 4μ , breadth 8 to 10μ
Leptotheca parva Th  lohan 1895.....(61)
- 17(16) Spore cylindrical; sutural diameter 4.5μ , breadth 9μ
Leptotheca glomerosa Davis 1917.....(65)
- 18(15) Average sutural diameter greater than 10μ 19
- 19(20) Posterior margin of spore slightly concave in front view; anterior end attenuated; polar capsules pyriform; sutural diameter 13μ , breadth 26μ . Trophozoite rounded; with active amoeboid movements
Leptotheca macropsora Auerbach 1909.....(62)
- 20(19) Posterior margin of spore more or less flattened; anterior end smoothly rounded; polar capsules rounded; sutural diameter 11μ , breadth 22μ . Trophozoite elongated; pseudopodia often anastomose
Leptotheca scissura Davis 1917.....(64)
- Incompletely described species
Leptotheca renicola Th  lohan 1895.....(61)
Leptotheca hepseti Th  lohan 1895.....(62)
Leptotheca perlata Gurley 1894.....(62)
Leptotheca sp. Awerinzew 1908.....(62)

Genus CERATOMYXA Th  lohan 1892

- 1(52) Spore constant in form and size..... 2
- 2(21) Sutural diameter equal to or less than one-eighth of total breadth..... 3
- 3(10) Sutural diameter not less than one-tenth of total breadth..... 4
- 4(19) Pseudopodia of vegetative form located at rounded end..... 5
- 5(16) Pseudopodia long filiform; with slow whiplash-like movements toward pointed extremity. Spore: sutural diameter 12μ , breadth 118μ
Ceratomyxa flagellifera Davis 1917.....(77)

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

- 6(5) Pseudopodia short lobose. 7
- 7(8) Extremities of spore attenuated; spore large; sutural diameter 10 to 12 μ , breadth 90 to 100 μ
Ceratomyxa sphaerulosa Thélohan 1892.(66)
- 8(7) Extremities of spore rounded; spore small; sutural diameter 4 μ , breadth 34 to 39 μ
Ceratomyxa streptospora Davis 1917.(79)
- 9(4) Pseudopodia unlocalized; from main part of sporulating trophozoite are branched out from one to six long prolongations. Spore: sutural diameter 5 to 7 μ , breadth 50 μ
Ceratomyxa appendiculata Thélohan 1892.(67)
- 10(3) Sutural diameter equal to or less than one-tenth of total breadth. 11
- 11(14) Shell-valve terminating in a fine thread-like process at distal end. 12
- 12(13) Sutural diameter 7 to 8 μ , breadth 40 to 50 μ , lateral process 250 to 300 μ
Ceratomyxa acadensis Mavor 1915.(71)
- 13(12) Sutural diameter 5 μ , breadth 10 to 12 μ , length of lateral process 20 μ
Ceratomyxa linospora Doflein 1898.(69)
- 14(11) Shell-valves not terminating in thread-like processes. 15
- 15(20) Shell-valve drawn out into a delicate process. 16
- 16(17) Lateral process ribbon-like; sutural diameter 6 μ , breadth 140 to 150 μ
Ceratomyxa taenia Davis 1917.(74)
- 17(16) Lateral process not ribbon-like, but circular in cross-section. 18
- 18(19) Posterior margin of main part of spore flattened; sutural diameter 12 μ , breadth 115 to 140 μ ; trophozoite disporous
Ceratomyxa sphairophora Davis 1917.(73)
- 19(18) Posterior margin of main part of spore rounded; sutural diameter 7 μ , breadth 80 μ . Trophozoite monosporous or disporous
Ceratomyxa spinosa Davis 1917.(80)
- 20(15) Shell-valve tapering gradually to attenuated point; asymmetrical; sutural diameter 9 μ , breadth 115 μ
Ceratomyxa attenuata Davis 1917.(75)
- 21(2) Sutural diameter more than one-eighth of total breadth. 22
- 22(45) Sutural diameter equal to or more than one-fifth of breadth. 23
- 23(34) Shell-valves symmetrically built. 24
- 24(29) Shell-valves attenuated at distal end. 25
- 25(26) Pseudopodia peculiar network-like form. Spore: sutural diameter 12 to 20 μ , breadth 50 to 80 μ
Ceratomyxa ramosa Awerinzew 1907.(69)
- 26(25) Pseudopodia never unite together. 27
- 27(28) Shell-valves curved greatly posteriad; polar capsules rounded; sutural diameter 8 to 9 μ , breadth 16(?) μ . Trophozoite elongated pyriform.
Ceratomyxa recurvata Davis 1917.(75)
- 28(27) Shell-valves not curved; two thickenings on posterior margin equidistant from sutural line; polar capsules pyriform; sutural diameter 40 to 45 μ , thickness 25 to 30 μ , breadth 124 to 140 μ
Ceratomyxa tylosuri Awerinzew 1913.(70)
- 29(24) Shell-valve rounded at distal end. 30
- 30(31) Spore arch-shaped; sutural diameter and thickness 12 to 15 μ , breadth 50 to 60 μ . Trophozoite large amoeboid.
Ceratomyxa spari Awerinzew 1913.(70)
- 31(30) Spore straight. 32

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

32(33)	Shell-valves shorter (sutural diameter: breadth=1:1.6) <i>Ceratomyxa coris</i> Georgévitch 1916.....	(72)
33(32)	Shell-valves longer (sutural diameter: breadth=1:2.6) <i>Ceratomyxa herouardi</i> Georgévitch 1916.....	(72)
34(23)	Shell-valves asymmetrically built.....	35
35(40)	Spore arch-shaped in front view.....	36
36(37)	Sutural diameter equal to one-fifth of breadth; sutural diameter 10 μ , breadth 50 μ <i>Ceratomyxa globulifera</i> Thélohan 1895.....	(67)
37(36)	Sutural diameter more than one-fifth of total breadth.....	38
38(39)	Spore: breadth shorter; sutural diameter 14 μ , breadth 17 μ <i>Ceratomyxa abbreviata</i> Davis 1917.....	(76)
39(38)	Spore: breadth longer; sutural diameter 12 to 15 μ , breadth 45 to 50 μ <i>Ceratomyxa reticularis</i> Thélohan 1895.....	(68)
40(35)	Spore straight.....	41
41(42)	Sutural diameter 11 μ , breadth 27 μ . Trophozoite always rounded <i>Ceratomyxa amorpha</i> Davis 1917.....	(78)
42(41)	Sutural diameter 6 μ	43
43(44)	Trophozoite with active pseudopodia. Spore: sutural diameter 6 μ , breadth 22 to 24 μ <i>Ceratomyxa undulata</i> Davis 1917.....	(79)
44(43)	Trophozoite with inactive pseudopodia. Spore: sutural diameter 6 μ , breadth 31 μ <i>Ceratomyxa inaequalis</i> Doflein 1898.....	(68)
45(22)	Sutural diameter less than one-fifth of total breadth.....	46
46(49)	Breadth of spore equal to or greater than 50 μ	47
47(48)	Shell-valve tapering gradually toward distal end; sutural diameter 8 μ , breadth 50 to 56 μ . Trophozoite usually elongated pyriform <i>Ceratomyxa mesospora</i> Davis 1917.....	(73)
48(47)	Shell-valve rounded at distal end; sutural diameter 6 to 7 μ , breadth 50 μ . Trophozoite usually rounded or irregular form; size small <i>Ceratomyxa aggregata</i> Davis 1917.....	(79)
49(46)	Breadth of spore smaller than 30 μ	50
50(51)	Trophozoite ordinarily spherical, diameter not exceeding 16 to 20 μ ; protoplasm extremely pale looking. Spore: sutural diameter 5 μ , breadth 25 to 30 μ . <i>Ceratomyxa pallida</i> Thélohan 1895.....	(67)
51(50)	Trophozoite pyriform with a long posterior process; movements by wavelike motion of ectoplasm; also active backward movements of pseudopodia. Spore asymmetrically built; sutural diameter 5 μ , breadth 24 to 28 μ <i>Ceratomyxa agglomerata</i> Davis 1917.....	(77)
52(1)	Spore variable in size and form.....	53
53(54)	Variation in number of shell-valves conspicuous; sutural diameter 5 μ , breadth 25 μ <i>Ceratomyxa truncata</i> Thélohan 1895.....	(67)
54(53)	Variable in size and form of spore, but not in number of shell-valve.....	55
55(60)	Trophozoite more or less definite in shape.....	56
56(59)	Trophozoite usually pyriform.....	57
57(58)	Trophozoite disporous. Spore: sutural diameter 7 to 9 μ , breadth 15 to 38 μ <i>Ceratomyxa lunata</i> Davis 1917.....	(76)
58(57)	Trophozoite monosporous or disporous. Spore: sutural diameter 5 to 6 μ , breadth 18 to 25 μ <i>Ceratomyxa monospora</i> Davis 1917.....	(78)

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

- 59(56) Trophozoite always rounded, never pyriform. Spore: sutural diameter 6μ , breadth 16μ
Ceratomyxa navicularia Davis 1917.....(80)
- 60(55) Trophozoite polymorphous..... 61
- 61(62) Shell-valves symmetrically built; sutural diameter 5 to 8μ , breadth 20 to 31μ
Ceratomyxa arcuata Thélohan 1892.....(65)
- 62(61) Shell-valves often asymmetrically built; sutural diameter 8 to 14μ , breadth 50 to 80μ
Ceratomyxa drepanopsettae Awerinzew 1907.....(70)
- Incompletely described species
- Ceratomyxa* sp. (?) Awerinzew 1913.....(71)
- Ceratomyxa* sp. (?) Awerinzew 1913.....(71)
- Ceratomyxa* sp. Georgévitch 1916.....(72)

Genus MYXOPROTEUS Doflein 1898

- 1(2) Spore with two long (5μ) processes extending anteriad from sides; sutural diameter 9μ , breadth 12μ
Myxoproteus cornutus Davis 1917.....(82)
- 2(1) Spore without long process..... 3
- 3(4) Spore with two small spinous processes at anterior end; sutural diameter 25μ , breadth 18 to 20μ
Myxoproteus ambiguus Thélohan 1895.....(81)
- 4(3) Spore without any process; posterior end slightly pointed; sutural diameter 12μ , breadth 10 to 11μ
Myxoproteus cordiformis Davis 1917.....(81)

Genus WARDIA Kudo 1919

- 1 Spore isosceles triangular form; shell with network-like striations which end in fringe-like processes at posterior margin; sutural diameter 9 to 10μ , breadth 10 to 12μ , diameter of polar capsule 4μ .
Wardia ovinocua Kudo 1919.....(82)
- Doubtfully placed species
- Wardia ohlmacheri* (Gurley 1894).....(83)

Genus MITRASPORA Fujita 1912 emend. Kudo 1919

- 1(4) Spore with posterior filaments..... 2
- 2(3) Posterior filaments short (5 to 6μ long); length 10μ , breadth 8 to 9μ , thickness 6 to 8μ , polar capsule 4μ by 1.5 to 2μ
Mitraspora cyprini Fujita 1912.....(84)
- 3(2) Posterior filaments of spore long (up to 28μ); length 10 to 11μ , polar capsule 4 to 4.5μ long
Mitraspora caudata Parisi 1910.....(85)
- 4(1) Spore without posterior filament; anterior end slightly attenuated; posterior end truncate; length 15 to 17μ , breadth 5 to 6μ , thickness 4.5 to 5.5μ , polar capsule 7.5μ by 2μ
Mitraspora elongata Kudo 1919.....(85)

Genus CHLOROMYXUM Mingazzini 1890

- 1(4) Spore with posterior appendage..... 2
- 2(3) Posterior appendage fine and numerous; length 6 to 9μ , breadth 5 to 6μ , polar capsule 2 to 3μ by 1 to 2μ
Chloromyxum leydigi Mingazzini 1890.....(87)

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

- 3(2) Posterior appendage single or bifurcated; length 8μ , breadth 6 to 7μ , appendage 10μ long
Chloromyxum caudatum Thélohan 1895.....(88)
- 4(1) Spore without posterior appendage..... 5
- 5(34) Spore circular or subcircular in front view; parasitic in body cavity of host..... 6
- 6(29) Shell-valves marked with striations or ridges..... 7
- 7(24) Main part of striations or ridges parallel to sutural line..... 8
- 8(11) Shell-valves partially marked..... 9
- 9(10) Ridges on each shell-valve variable in number (six found in original drawing) running closely to sutural line; diameter 10.8μ
Chloromyxum dubium Auerbach 1908.....(91)
- 10(9) Each shell-valve with one ridge from which eight to twelve short ones are directed toward centre of valve; oval in profile; length and breadth 8 to 10μ , thickness 5 to 7μ
Chloromyxum trijugum Kudo 1919.....(96)
- 11(8) Entire shell-valve marked..... 12
- 12(19) Shell-valve marked with fine striations..... 13
- 13(16) Spore oval in lateral view..... 14
- 14(15) Trophozoite larger; size up to 50μ by 20μ ; polysporous (up to eight spores) rarely disporous. Spore: length 8 to 9μ , breadth 6 to 7μ , thickness 5 to 6μ
Chloromyxum misgurni Kudo 1916.....(93)
- 15(14) Trophozoite smaller; size up to 35μ in diameter; polysporous (up to six spores) or disporous. Spore: length 8μ , breadth 7μ , thickness 5 to 6μ
Chloromyxum catostomi Kudo 1919.....(98)
- 16(13) Spore circular in lateral view..... 17
- 17(18) Trophozoite rounded; 40 to 45μ by 28 to 40μ . Spore: diameter 10 to 13μ , polar capsule 4 to 6μ long
Chloromyxum protei Joseph 1905.....(90)
- 18(17) Trophozoite irregular form; 33 to 35μ in average length. Spore: striations thicker and somewhat wavy; diameter 9 to 9.5μ , polar capsule 3μ long
Chloromyxum thymalli Lebzelter 1912.....(92)
- 19(12) Shell-valves marked with ridges..... 20
- 20(21) Trophozoite small (average diameter of adults about 20μ); monosporous, rarely disporous. Spore: shell-valves with ridges marked antero-posteriad; diameter 10 to 11μ
Chloromyxum cristatum Léger 1906.....(91)
- 21(20) Trophozoite large, diameter reaching 40 to 50μ 22
- 22(23) Ridges on shell-valves united into a line at each end and unequal in thickness; spore small; length 10 to 12μ , breadth 8 to 10μ
Chloromyxum fujitai Kudo 1916.....(93)
- 23(22) Shell-valve with two circular and two small ridges; spore large; length 16μ , breadth 10μ
Chloromyxum koi Fujita 1913.....(92)
- 24(7) Striations or ridges not parallel to sutural line..... 25
- 25(26) Striations irregular; posterior margin thickened at sides; diameter 7.5 to 9μ
Chloromyxum wardi Kudo 1919.....(99)
- 26(25) Striations parallel to each other..... 27
- 27(28) Striations forming acute angles with sutural line; diameter 8 to 9μ
Chloromyxum truttiae Léger 1906.....(90)
- 28(27) Four ridges on posterior half of shell-valve converging toward anterior end; diameter 7μ
Chloromyxum granulorum Davis 1917.....(95)

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

29(6)	Shell-valves without marking, beside sutural ridge	30
30(31)	Anterior end of spore rounded; diameter 7 to 8 μ ; one or two short spinous thickenings at posterior margin <i>Chloromyxum fluviatile</i> Th��lohan 1892	(89)
31(30)	Anterior end of spore mucronate or truncate	32
32(33)	Anterior end of spore mucronate; length 8 μ <i>Chloromyxum mucronatum</i> Gurley 1893	(89)
33(32)	Anterior end of spore truncate; spore large; length 40 to 48 μ , breadth 30 to 38 μ <i>Chloromyxum magnum</i> Awerinzew 1913	(92)
34(5)	Spore rounded quadrangular in end view; conical in front view; parasitic in muscular tissue of fish	35
35(36)	Length of spore larger than breadth; length 6 μ , breadth 5 μ <i>Chloromyxum quadratum</i> Th��lohan 1895	(88)
36(35)	Length (sutural diameter) of spore smaller than breadth	37
37(38)	Spore variable in form; anterior end narrower or broader than posterior end; length 4 to 4.75 μ , breadth 5.4 to 6.5 μ <i>Chloromyxum clupeiidae</i> Hahn 1917	(94)
38(37)	Anterior end of spore drawn out; almost circular in end view; length 6 μ , breadth 7.5 μ <i>Chloromyxum funduli</i> Hahn 1915	(93)
Incompletely described species		
	<i>Chloromyxum diploxyi</i> Gurley 1893	(90)
	<i>Chloromyxum</i> sp. Awerinzew 1908	(91)

Genus SPHAEROSPORA Th  lohan 1892

1(8)	Shell-valve of spore without marking except sutural ridge	2
2(7)	Vegetative form amoeboid	3
3(4)	Movements of vegetative form active. Spore: sutural ridge fairly well marked; a pair of short filaments become visible at anterior end on warming; diameter 8 μ <i>Sphaerospora masovica</i> Cohn 1902	(101)
4(3)	Vegetative form without active movements	5
5(6)	Spore: sutural ridge not prominent; polar capsule pyriform; diameter 8 to 13 μ , polar capsule 4 to 5 μ by 2.5 to 3.5 μ <i>Sphaerospora carassii</i> Kudo 1919	(103)
6(5)	Spore: sutural ridge prominent; polar capsule spherical; slightly attenuated at anterior end; diameter 10 to 11 μ <i>Sphaerospora elegans</i> Th��lohan 1892	(100)
7(2)	Vegetative form produces cyst in tissue. Spore: diameter 8 to 9 μ <i>Sphaerospora platessae</i> Woodcock 1904	(102)
8(1)	Shell-valves striated	9
9(10)	Polar capsules divergent; diameter of spore 10 to 12 μ , thickness 8 μ <i>Sphaerospora divergens</i> Th��lohan 1895	(100)
10(9)	Polar capsules not divergent	11
11(14)	Striation marked antero-posteriad	12
12(13)	Spore with a quadrangular lamella at anterior margin; striations ending in small spines at posterior margin; length 12 to 14 μ , breadth 10 to 12 μ <i>Sphaerospora rostrata</i> Th��lohan 1895	(101)
13(12)	Spore smooth-contoured; polar capsules parallel to each other; diameter 7 to 10 μ <i>Sphaerospora polymorpha</i> Davis 1917	(102)

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

- 14(11) Faint concentric striations; pointed at sides and middle part of posterior margin; polar capsules unequal in size; length 7 to 8 μ , breadth 6 to 7 μ , thickness 5 μ
Sphaerospora angulata Fujita 1912.....(102)
- Incompletely described species
Sphaerospora sp. Davis 1917.....(103)
Sphaerospora sp. Southwell et Prashad 1918.....(103)

Genus SINUOLINEA Davis 1917

- 1(4) Spore with two processes..... 2
 2(3) Processes lateral and long (20 μ); spore: 9 to 11 μ long, 9 μ broad, process 18 to 22 μ long
Sinuolinea brachiophora Davis 1917.....(106)
- 3(2) Processes posteriad from sides and short; diameter 12 to 13 μ . Trophozoite opaque
Sinuolinea opacita Davis 1917.....(106)
- 4(1) Spore without process..... 5
 5(6) Trophozoite with active amoeboid movements. Spore: sutural ridge S-shaped at anterior part; length 15 μ , breadth 12 μ , thickness 8 μ
Sinuolinea arborescens Davis 1917.....(105)
- 6(5) Trophozoite with slow amoeboid movements..... 7
 7(8) Sutural plane much twisted on its axis; capsulogenous cells large occupying more than half of sporal cavity; polar capsules opening on opposite sides; diameter 12 to 14 μ
Sinuolinea capsularis Davis 1917.....(105)
- 8(7) Sutural plane not much twisted; diameter 15 μ
Sinuolinea dimorpha Davis 1916.....(104)

Genus MYXIDIUM Bütschli 1882

- 1(16) Breadth of spore equal to or more than half of length..... 2
 2(7) Shell-valves unstriated..... 3
 3(6) Sutural plane curved into an S..... 4
 4(5) Spore small; length 8 to 12 μ , breadth 4 to 6 μ
Myxidium incurvatum Thélohan 1892.....(108)
- 5(4) Spore large; much broader; length 20.8 to 23.4 μ , breadth 13 to 15.6 μ
Myxidium inflatum Auerbach 1909.....(111)
- 6(3) Sutural plane straight; spore cylindrical; surrounded by a gelatinous envelope; length 10 to 11 μ , breadth 6 μ
Myxidium glutinosum Davis 1917.....(115)
- 7(2) Shell-valves striated..... 8
 8(9) Sutural line curved into an S; form oval; circular in cross-section; openings of polar capsules pointed; length 11 to 13 μ , breadth 8 to 9 μ
Myxidium oviforme Parisi 1912.....(114)
- 9(8) Sutural line straight..... 10
 10(13) Sutural line coincides with longitudinal axis of spore..... 11
 11(12) Sutural ridge distinct; extremities mucronate; length 9 to 10 μ , breadth 5 to 5.6 μ , thickness 4.75 to 5 μ . Vegetative form produces cysts, 800 to 900 μ in diameter
Myxidium giardi Cépède 1906.....(110)
- 12(11) Sutural ridge faintly marked; extremities gradually drawn out; length 11 μ , breadth 8 μ . Trophozoite large and leaf-like; diameter up to 1.35 mm.
Myxidium phyllium Davis 1917.....(116)

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

- 13(10) Sutural line forming an acute angle with longitudinal axis of spore 14
- 14(15) Shell thickened at extremities; polar capsules ovoidal; length 10 to 14 μ , breadth 6 to 8 μ , length of polar capsule 4 μ
Myxidium striatum Cunha et Fonseca 1917 (116)
- 15(14) Shell uniformly thick; polar capsules rounded pyriform; length 10 to 12 μ , breadth 6 μ , length of polar capsule 3 to 4 μ
Myxidium macrocapsulare Auerbach 1910 (113)
- 16(1) Breadth of spore less than half of length 17
- 17(34) Breadth more than one-third of length 18
- 18(25) Shell-valves unstriated 19
- 19(22) Extremities of spore pointed 20
- 20(21) Spore: extremities sharply pointed; greatly curved; narrow; length 12 to 14 μ , breadth 5.5 to 6 μ , thickness 2.5 to 3 μ
Myxidium depressum Parisi 1912 (114)
- 21(20) Spore: extremities not so sharply pointed; not greatly curved; broader; length 16.2 to 19 μ , breadth 7 to 9 μ
Myxidium bergense Auerbach 1909 (112)
- 22(19) Extremities of spore not pointed 23
- 23(24) Spore larger; length 15 to 20 μ , breadth 7 to 8 μ
Myxidium sphaericum Th  lohan 1895 (109)
- 24(23) Spore smaller; length 6(?) to 14 μ , breadth 4 to 6 μ . Trophozoite mictosporous
Myxidium gadi Georg  vitch 1916 (115)
- 25(18) Shell-valves striated 26
- 26(33) Spore definite in shape 27
- 27(28) Spore constricted in middle part of length; length 15 μ
Myxidium histophilum Th  lohan 1895 (109)
- 28(27) Spore regularly fusiform 29
- 29(30) Vegetative form produces cyst. Spore: length 12 to 15 μ , breadth 6 μ
Myxidium barbatulae C  p  de 1906 (110)
- 30(29) Vegetative form does not produce cyst 31
- 31(32) Sutural line slightly curved in S-form; length 15 to 16 μ , breadth and thickness 5.5 to 6 μ
Myxidium americanum Kudo 1919 (117)
- 32(31) Sutural line not curved in S-shape, but bent to one side; length 15 to 18 μ , breadth and thickness 6 to 7 μ
Myxidium kagayamai Kudo 1919 (117)
- 33(26) Spore variable in form; straight and constricted; one side concave, the other convex; arch-shaped, etc.; length 13 to 18 μ , breadth 5.2 to 5.8 μ
Myxidium pfeifferi Auerbach 1908 (111)
- 34(17) Breadth of spore equal to or less than one-third of length 35
- 35(40) Shell-valves unstriated 36
- 36(37) Spore greatly elongated (breadth: length = 1:6.2); length 21.6 to 25.2 μ , breadth 3.6 to 4 μ
Myxidium procerum Auerbach 1910 (112)
- 37(36) Spore less elongated (breadth: length = 1:3 or 1:3.4) 38
- 38(39) Spore large; valves asymmetrical; length 28 μ , breadth 8 μ
Myxidium giganteum Doflein 1898 (110)
- 39(38) Spore small; valves symmetrical; length 12 μ , breadth 3 to 4 μ
Myxidium danilewskyi Laveran 1897 (109)
- 40(35) Shell-valves striated 41
- 41(44) Spore definite in shape 42

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

- 42(43) Length of polar capsule more than one-fourth of that of spore; spore 18 to 20 μ long, 5 to 6 μ broad
Myxidium lieberkühni Bütschli 1882.....(107)
- 43(42) Length of polar capsule less than one-seventh of that of spore; length 16 to 17 μ , breadth 5 μ
Myxidium mackiei Bosanquet 1910.....(112)
- 44(41) Spore variable in shape; S-form, straight fusiform, etc.; length 9.1 μ , breadth 2.8 μ . Vegetative form produces cyst
Myxidium anguillae Ishii 1915.....(114)
- Incompletely described species
Myxidium sp. Gurley 1894.....(109)
Myxidium sp. Awerinzew 1908.....(113)
Myxidium sp. Mavor 1915.....(115)

Genus SPHAEROMYXA Thélohan 1892

- 1(6) Spore straight, not arch-shaped..... 2
- 2(5) Shell-valves symmetrical..... 3
- 3(4) Ends of spore truncate; striations longitudinal; length 15 to 20 μ , breadth 5 to 6 μ .
Sphaeromyxa balbianii Thélohan 1892.....(118)
- 4(3) Ends of spore rounded; sutural plane forming some angles with longitudinal axis of spore; striations transverse; length 12 to 14 μ , breadth 9 to 10 μ
Sphaeromyxa immersa Lutz 1889.....(119)
- 5(2) Shell-valves asymmetrical; unstriated; ends less truncate; dimensions about twice or three times larger than those of *Sphaeromyxa balbianii*
Sphaeromyxa gasterostei Georgévitch.....(121)
- 6(1) Spore arch-shaped, not straight..... 7
- 7(8) Shell-valves indistinctly striated; ends truncate; length 22 to 28 μ , breadth 3 to 4.3 μ
Sphaeromyxa sabrazesi Laveran et Mesnil 1900.....(120)
- 8(7) Shell-valves unstriated..... 9
- 9(10) Spore extremely large; length 75 to 80 μ , breadth 18 to 20 μ ; ends slightly tapering
Sphaeromyxa exneri Awerinzew 1913.....(121)
- 10(9) Spore less than 35 μ in length..... 11
- 11(12) Extremities rounded; length 30 to 35 μ , breadth 8 μ
Sphaeromyxa incurvata Doflein 1898.....(119)
- 12(11) Extremities truncate; sutural ridge often twisted in S-form; length 20.8 to 26 μ , breadth and thickness 5.4 μ
Sphaeromyxa hellandi Auerbach 1909.....(121)

Genus ZSCHOKKELLA Auerbach 1910

- 1(4) Shell-valves unstriated..... 2
- 2(3) Openings of polar capsules on flattened side; spore large; length 16 to 28.8 μ , breadth 13 to 18 μ
Zschokkella hildae Auerbach 1910.....(122)
- 3(2) Openings of polar capsules at pointed ends; spore small; length 11 μ , breadth 7 μ
Zschokkella globulosa Davis 1917.....(123)
- 4(1) Shell-valves striated..... 5
- 5(6) Openings of polar capsules at pointed ends; polar capsules spherical; spore larger; length 10 to 14 μ , breadth 6 to 7 μ
Zschokkella acheilognathi Kudo 1916.....(123)

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

- 6(5) Openings of polar capsules on side; polar capsules rounded pyriform; spore smaller; length 9.5 to 11.5 μ , breadth 6.5 to 7 μ
Zschokkella nova Klokacewa 1914.....(122)

Genus MYXOSOMA Thélohan 1892

- 1(2) Spore: shell thickened at anterior end; length 12 to 13 μ , breadth 7 to 8 μ , polar capsule 6 to 7 μ by 2 to 3 μ . Cysts polymorphous
Myxosoma dujardini Thélohan 1892.....(124)
- 2(1) Spore: shell of uniform thickness and with seven to ten folds on sutural edge; length 14 μ , breadth 8 μ , thickness 6 μ , polar capsule 8 μ by 2 μ . Cysts spherical up to 360 μ in largest diameter
Myxosoma funduli Kudo 1918.....(125)

Ambiguous form

Myxosoma lobatum Nemeček 1911.....(124)

Genus LENTOSPORA Plehn 1905

- 1(8) Spore circular in front view..... 2
- 2(3) Vegetative form produces cysts. Spore: length and breadth 6.3 to 7 μ , thickness 4.2 to 4.9 μ
Lentospora dermatobia Ishii 1916.....(127)
- 3(2) Vegetative form does not produce cysts or cysts unobserved..... 4
- 4(5) Spore small; trophozoites found in the blood vessel of the brain. Spore 5 to 5.5 μ in diameter.
Lentospora encephalina Mulsow 1911.....(126)
- 5(4) Spore large, greater than 7.5 μ in average diameter..... 6
- 6(7) Spore slightly pointed at anterior end; length 8 to 10 μ , breadth 7 to 8 μ , thickness 5 to 6 μ
Lentospora acuta Fujita 1912.....(127)
- 7(6) Anterior end of spore rounded; diameter 6 to 10 μ
Lentospora cerebralis Plehn 1905.....(125)
- 8(1) Spore oval in front view..... 9
- 9(10) Spore symmetrically built; length 12 μ , breadth 9.5 μ , thickness 6 μ
Lentospora multiplicata Reuss 1906.....(126)
- 10(9) Spore asymmetrically built; length 10 to 11 μ , breadth 6.5 to 7 μ
Lentospora asymmetrica Parisi 1912.....(126)

Genus MYXOBOLUS Bütschli 1882

- 1(18) Spore with one polar capsule..... 2
- 2(9) Breadth of spore equal to or more than half of length..... 3
- 3(6) Breadth of spore equal to half of length..... 4
- 4(5) Spore larger; often calabash-shaped; anterior end drawn out into a rounded tip; shell thickened at tip; length 15 μ , breadth 7 to 8 μ , thickness 5 to 6 μ
Myxobolus toyamai Kudo 1915.....(131)
- 5(4) Spore smaller; anterior end pointed; shell of uniform thickness; length 9 to 10 μ , breadth 4.5 to 5.5 μ , thickness 3 μ
Myxobolus oculi-leucisci Trojan 1909.....(130)
- 6(3) Breadth of spore more than two-thirds of length..... 7
- 7(8) Polar capsule small and oblique in position
Myxobolus unicapsulatus Gurley 1893.....(129)
- 8(7) Polar capsule long and median in position; spore broader; length 13.2 to 13.6 μ , breadth 10.1 to 10.3 μ
Myxobolus seni Southwell et Prashad 1918.....(132)

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

9(2)	Breadth of spore less than half of length.....	10
10(17)	Breadth of spore more than two-fifths of length.....	11
11(16)	Spore without any process.....	12
12(15)	Shell of uniform thickness.....	13
13(14)	Spore bent to one side; shell thickened at slightly rounded anterior tip; sutural edge without marking; length 16 to 18 μ , breadth 7 to 8 μ , polar capsule 5 to 7 μ long <i>Myxobolus piriformis</i> Th��lohan 1892.....	(129)
14(13)	Spore straight; shell thickened at posterior part; sutural edge with four to six markings; length 18 to 20 μ , breadth 8 μ , thickness 6 μ , polar capsule 9 to 10 μ long <i>Myxobolus fuhrmanni</i> Auerbach 1909.....	(130)
15(12)	Shell of uniform thickness; valves symmetrical; sutural edge with markings up to 12 in number; spore 14 to 15.5 μ long, 6 to 7.3 μ broad, 5 to 6 μ thick, polar capsule 6.3 μ by 2 to 3 μ <i>Myxobolus misgarni</i> Kudo 1919.....	(133)
16(11)	Spore with a posterior process, 5 μ in length and as broad as spore; length 17 to 18 μ , breadth 7.5 to 8 μ , polar capsule 7 μ by 4 μ <i>Myxobolus notatus</i> Mavor 1916.....	(131)
17(10)	Breadth of spore about one-fourth of length; spore large; polar capsule extremely large; length 30 to 32 μ , breadth 7 to 8 μ , length of polar capsule 22 to 23 μ <i>Myxobolus rohita��</i> Southwell et Prashad 1918.....	(132)
18(1)	Spore with two polar capsules.....	19
19(24)	Form of spore variable.....	20
20(23)	Spore with an intercapsular appendix at anterior end.....	21
21(22)	Spore oval; length 10 to 12 μ , breadth 8 to 9 μ , thickness 6 μ , polar capsule 5 μ by 2 to 3 μ <i>Myxobolus m��lleri</i> B��tschli 1882.....	(128)
22(21)	Spore pyriform or elongated oval; length 11 to 16 μ , breadth 8 to 13 μ , polar capsule 6 μ by 4 μ <i>Myxobolus cycloides</i> Gurley 1893.....	(140)
23(20)	Spore without intercapsular appendix; circular form 7 to 8 μ , breadth 8 to 10 μ , thickness 6 μ , polar capsule 4 to 5 μ by 2 μ <i>Myxobolus hyla��</i> Johnston et Bancroft 1918.....	(153)
24(19)	Form of spore definite.....	25
25(28)	Polar capsules in each spore regularly of considerably different size.....	26
26(27)	Spore with an intercapsular appendix; anterior end rounded; sutural edge with folds (3 to 5); length 10 to 12 μ , breadth 8 μ <i>Myxobolus dispar</i> Th��lohan 1895.....	(135)
27(26)	Spore without intercapsular appendix; anterior end pointed; no fold on sutural edge <i>Myxobolus inaequalis</i> Gurley 1893.....	(135)
28(25)	Polar capsules approximately of equal form and size.....	29
29(30)	Sutural diameter smaller than breadth; length 6 to 7 μ , breadth 8 μ <i>Myxobolus transovalis</i> Gurley 1893.....	(139)
30(29)	Length equal to or more than breadth of spore.....	31
31(102)	Length longer than breadth.....	32
32(37)	Breadth of spore less than half of length.....	33
33(34)	Extremities of spore equally pointed; length 13 to 14.5 μ , breadth 6 to 7 μ , polar capsule 4.5 μ long <i>Myxobolus miyaii</i> Kudo 1919.....	(155)
34(33)	Anterior end of spore attenuated; posterior end rounded.....	35

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

35(36)	Shell thickened at posterior margin; spore 12 to 15 μ long, 4 to 6.8 μ broad, polar capsule 5.5 to 7 μ by 2.1 to 2.5 μ <i>Myxobolus anurus</i> Cohn 1895.....	(142)
36(35)	Shell of uniform thickness; spore 14.3 μ long, 6.7 μ broad, polar capsule 6.5 μ by 2 μ <i>Myxobolus funduli</i> Kudo 1919.....	(151)
37(32)	Breadth of spore greater than half of length.....	38
38(41)	Length of spore greater than 20 μ	39
39(40)	Spore large; subcircular and anterior end flattened in front view; sutural edge with markings; length 38 to 45 μ , breadth 32 to 38 μ , thickness 28 to 35 μ , polar capsule 15 to 17 μ long <i>Myxobolus magnus</i> Awerinzew 1913.....	(150)
40(39)	Spore small; extremities equally rounded; length 21 μ , breadth 15 μ , polar capsule 6 μ long <i>Myxobolus cyprini</i> Doflein 1898.....	(143)
41(38)	Length of spore less than 20 μ	42
42(43)	Spore with a wide (2 to 3 μ) membranous posterior process; length 12 μ , breadth 10 μ , length of polar capsule 4.5 μ <i>Myxobolus cordis</i> Keysselitz 1908.....	(148)
43(42)	Normal spore without appendage.....	44
44(49)	Breadth of sutural ridge one-third of thickness of spore.....	45
45(46)	Length of spore smaller than 10 μ ; subcircular in front view; length 7 to 8 μ , breadth 6 to 7 μ , thickness 5 μ <i>Myxobolus globosus</i> Gurley 1893.....	(139)
46(45)	Length of spore greater than 10 μ	47
47(48)	Spore large; elliptical in front view; with an intercapsular appendix, sutural edge with markings; length 16.9 to 21.6 μ , breadth 13 to 16.2 μ , thickness 9 μ <i>Myxobolus gigas</i> Auerbach 1906.....	(145)
48(47)	Spore small; subcircular in front view; markings on entire sutural edge; length 10.8 to 11.7 μ , breadth 9.9 to 10.4 μ , thickness 7.2 to 9 μ <i>Myxobolus aeglefini</i> Auerbach 1906.....	(144)
49(44)	Sutural ridge narrower.....	50
50(69)	Spore with an intercapsular appendix.....	51
51(68)	Intercapsular appendix triangular.....	52
52(57)	Anterior end of spore attenuated in front view.....	53
53(54)	Sutural edge without marking; length 11 to 12 μ , breadth 9.25 to 10.5 μ <i>Myxobolus ballerii</i> Reuss 1906.....	(147)
54(53)	Sutural edge with markings.....	55
55(56)	Spore small; length 8 to 9.5 μ , breadth 6 to 7.5 μ , thickness 5.5 μ <i>Myxobolus exiguus</i> Th��lohan 1895.....	(136)
56(55)	Spore large; often subcircular; length 11.5 μ to 12 μ , breadth 7.5 to 8 μ , thickness 5 μ <i>Myxobolus obesus</i> Gurley 1893.....	(140)
57(52)	Anterior end of spore broadly rounded in front view.....	58
58(59)	Posterior portion of spore narrower; polar capsule rather large; length 11.4 to 13.5 μ , breadth 9.5 to 11 μ , thickness 8.5 to 9.5 μ , polar capsule 5.5 to 6 μ long <i>Myxobolus discrepans</i> Kudo 1919.....	(156)
59(58)	Extremities of spore approximately equal.....	60
60(61)	Sutural edge without markings; spore 10 to 10.5 μ long, 8 to 8.5 μ broad, polar capsule 4.5 μ long <i>Myxobolus squamae</i> Keysselitz 1908.....	(147)
61(60)	Sutural edge with markings.....	62
62(65)	Markings distinct.....	63

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

- 63(64) Marking variable in number along posterior margin of spore; spore more elongated; length 12 to 12.5 μ , breadth 10 to 10.5 μ , polar capsule 5.5 to 6 μ long
Myxobolus pfeifferi Th  lohan 1895. (133)
- 64(63) Sutural edge with four markings around posterior margin; spore rather short; length 11 to 12 μ , breadth 9 to 9.5 μ , thickness 4.5 to 5 μ , polar capsule 5 μ by 2.5 μ
Myxobolus scardinii Reuss 1906 (146)
- 65(62) Markings indistinct. 66
- 66(67) Markings about five at posterior margin; spore larger and shorter; length 11 to 12 μ , breadth 9.25 to 10 μ , thickness 4.5 to 5.5 μ , polar capsule 4 to 5 μ by 2.25 μ
Myxobolus bramae Reuss 1906. (147)
- 67(66) Markings many along entire sutural edge except anterior tip; spore smaller, longer and thicker; length 9.25 to 10 μ , breadth 7 to 7.25 μ , thickness 5 to 5.5 μ , polar capsule 4.5 μ by 2.5 to 3 μ
Myxobolus cyprinicola Reuss 1906. (147)
- 68(51) Intercapsular appendix rounded; sutural edge smooth; length 11 μ , breadth 8 μ , polar capsule 4 to 6 μ long
Myxobolus musculi Keysseltz 1908. (148)
- 69(50) Spore without intercapsular appendix. 70
- 70(75) Length of spore less than 10 μ 71
- 71(72) Spore very much flattened and small; length 6 μ , breadth 4.2 to 5 μ , polar capsule 3 μ by 2 μ
Myxobolus minutus Nemecek 1911. (150)
- 72(71) Thickness of spore about half of length. 73
- 73(74) Shell thick; length 9.25 to 10 μ , breadth 7.25 to 8.25 μ , thickness 4 to 5 μ
Myxobolus sandrae Reuss 1906. (146)
- 74(73) Shell thin; spore 8.25 to 9.5 μ long, 7.25 to 8.25 μ broad, 4.5 to 5.5 μ thick
Myxobolus volgensis Reuss 1906. (145)
- 75(70) Length of spore greater than 10 μ 76
- 76(85) Extremities of spore approximately equal. 77
- 77(80) Spore elongated (breadth: length = 1:1.8 or 1:1.4). 78
- 78(79) Spore larger; length 14 to 17 μ , breadth 8.5 μ , thickness 5 to 6 μ
Myxobolus oblongus Gurley 1893. (139)
- 79(78) Spore smaller; length 12 to 15 μ , breadth 9 to 11 μ
Myxobolus ellipsoides Th  lohan 1892. (136)
- 80(77) Spore shorter (breadth: length = 1:1.3, 1:1.2 or 1:1.1). 81
- 81(82) Sutural edge with markings; slightly truncate at anterior end in front view; spore 9.5 to 11.5 μ long, 8.5 to 9.5 μ broad, 6.5 μ thick, polar capsule 4.75 μ by 1.5 to 2 μ
Myxobolus mesentericus Kudo 1919. (157)
- 82(81) Sutural edge without markings. 83
- 83(84) Polar capsule larger; spore 13.9 μ long, 11 μ broad, 8 μ thick
Myxobolus lintoni Gurley 1893. (138)
- 84(83) Polar capsules smaller; spore 14.5 μ long, 11.9 μ broad, polar capsule 6 μ by 3.7 μ
Myxobolus pleuronectidae Hahn 1917. (152)
- 85(76) Anterior end of spore more attenuated than posterior. 86
- 86(89) Sutural edge with markings. 87
- 87(88) Markings five or six in number; spore 17 to 18 μ long, 10 to 13 μ broad, polar capsule 7 to 8 μ
Myxobolus permagnus Wegener 1910. (149)
- 88(87) Markings sometimes present; spore 13 to 17 μ long, 8 to 10 μ broad, 5 to 7 μ thick, polar capsule 6 to 7 μ long
Myxobolus carassii Klokacewa 1914. (150)

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

- 89(86) Sutural edge without markings. 90
 90(99) Anterior end of spore highly attenuated. 91
 91(96) Length of polar capsule equal to or less than half of that of spore. 92
 92(93) Spore: anterior end pointed; length 12.4 to 13.5 μ , breadth 6.5 to 7.5 μ , thickness 5 μ , polar capsule 6 to 7 μ long. Cysts of bright golden color
Myxobolus aureatus Ward 1919. (154)
 93(22) Anterior tip of spore not pointed. 94
 94(95) Spore greater in thickness (6.5 to 7 μ), length 12 to 13 μ , breadth 8.25 to 9 μ , polar capsule 6 μ by 2.5 μ ; anterior end more rounded
Myxobolus physophilus Reuss 1906. (146)
 95(94) Spore smaller in thickness (5.5 μ), length 11 to 13 μ , breadth 8.25 to 9.25 μ , polar capsule 6 μ by 2.5 to 3 μ ; anterior end less rounded
Myxobolus macrocapsularis Reuss 1906. (146)
 96(91) Length of polar capsule greater than half of that of spore. 97
 97(98) Length of polar capsule greater than two-thirds of that of spore; spore 16 μ long, 10 to 11 μ broad, polar capsule 11 μ by 4 μ
Myxobolus capsulatus Davis 1917. (152)
 98(97) Length of polar capsule less than two-thirds of that of spore; spore 14 to 16 μ long, 8 to 9 μ broad, 5 to 6 μ thick, polar capsule 8 to 9 μ by 2.5 to 3 μ
Myxobolus koi Kudo 1919. (155)
 99(90) Anterior end of spore rounded. 100
 100(101) Cysts: size up to 1.7 mm. by 0.7 mm.; parasitic in various tissues of host. Spore 10 to 12 μ long, 9 μ broad
Myxobolus oviformis Thélohan 1892. (137)
 101(100) Cysts: 0.9 mm. by 0.02 mm.; parasitic in nervous system. Spore 10 to 12 μ long, 8 μ broad, 6 μ thick, polar capsule 6 to 7 μ by 2 μ
Myxobolus neurobius Schuberg et Schröder 1905. (144)
 102(31) Spore almost circular in front view. 103
 103(104) Anterior end somewhat attenuated; sutural edge with four markings; spore 9 to 10 μ long and broad, 6.5 to 7 μ thick, polar capsule 6 to 7.5 μ by 2.5 to 3 μ
Myxobolus orbiculatus Kudo 1919. (155)
 104(103) Spore regularly circular in front view. 105
 105(106) Cysts large, up to 3 mm. by 1.5 mm. Spore 10 μ long, 9.8 μ broad, 3 μ thick, polar capsule 3.8 to 5 μ long
Myxobolus rotundus Nemeček 1911. (149)
 106(105) Cysts small, up to 0.33 mm. in diameter. Spore 9 μ in diameter
Myxobolus sphaeralis Gurley 1893. (141)
 Incompletely described species
Myxobolus sp. Gurley 1894. (142)
Myxobolus sp. Gurley 1894. (142)
Myxobolus sp. Gurley 1894. (143)
Myxobolus sp. Miyairi 1909. (149)
Myxobolus sp. Wegener 1910. (149)
Myxobolus sp. Lebzelter 1912. (150)
Myxobolus sp. Southwell 1915. (151)
Myxobolus sp. Kudo 1918. (132)

Genus HENNEGUYA Thélohan 1892

- 1(10) Parasitic in urinary bladder or urinary tubule of kidney of host. 2
 2(7) Shell-valves striated. 3

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

- 3(6) Length of tail equal to two-thirds of length of main part of spore* 4
- 4(5) Shell-valves asymmetrical; spore smaller; total length 38 to 48 μ , length of main part 15 μ , breadth 6 to 7.5 μ , polar capsule 7.5 to 9 μ by 3 to 3.5 μ . Trophozoite disporous and polysporous
Henneguya gasterosiei Parisi 1912.....(169)
- 5(4) Shell-valves symmetrical; spore broader; length of main part 13.5 to 15 μ , breadth 8 to 9 μ , thickness 6 to 7.5 μ , polar capsule 5 to 6 μ by 3 μ , length of tail 30 to 40 μ . Trophozoite mictosporous
Henneguya mictospora Kudo 1919.....(173)
- 6(3) Length of tail equal to half of length of main part of spore and single(?); length 20 to 24 μ , breadth 5 to 6 μ , length of polar capsule 4 to 5 μ
Henneguya media Th  lohan 1892.....(161)
- 7(2) Shell-valves unstriated..... 8
- 8(9) Spore elongated; polar capsule longer; posterior portion of main part broad; tail wider and bifurcated to same direction; total length 19.5 to 22.5 μ , length of main part 8.5 μ , breadth 6 μ , length of tail 8 to 8.5 μ
Henneguya l  geri C  p  de 1905.....(166)
- 9(8) Spore oval; polar capsule shorter; posterior part of main portion narrow; tail narrower and bifurcated to opposite directions; length of main part 11.5 μ , breadth 7 μ , length of tail 9.6 μ , polar capsule 3.5 μ by 2.5 μ
Henneguya wisconsinensis Mavor et Strasser 1916.....(170)
- 10(1) Parasitic in tissue of host..... 11
- 11(28) Tail always appears as a single process..... 12
- 12(13) Spore small; length 4 μ , breadth 2 μ
Henneguya tenuis Vaney et Conte 1901.....(166)
- 13(12) Spore longer and larger, at least 27 μ long..... 14
- 14(15) Sutural edge with eight to ten markings; tail rather long; length of main part 10 to 11.5 μ , breadth 8 to 8.75 μ , thickness 4 to 5 μ , polar capsule 3 to 4 μ by 2 μ , tail up to 17 μ long
Henneguya brachyura Ward 1919.....(171)
- 15(14) Sutural edge without markings..... 16
- 16(19) Total length of spore greater than 40 μ 17
- 17(18) Anterior end rounded; polar capsule large; shell-valves asymmetrical; tail long; main part 10 to 11 μ long, 6 to 8 μ broad, 4 μ thick, tail 30 to 40 μ long. Cysts elongated and large up to 6 mm. by 2 mm.
Henneguya macrura Gurley 1894.....(164)
- 18(17) Anterior end attenuated; polar capsule smaller; shell-valves symmetrical; tail shorter; length 20 μ , breadth 8 to 9 μ , polar capsule 8 μ by 2 to 3 μ . Cysts elongated oval and smaller, 1.1 mm. by 0.5 mm.
Henneguya creplini Gurley 1894.....(162)
- 19(16) Total length of spore less than 40 μ 20
- 20(21) Tail about one-third of main part; total length 38 μ , main part 26 μ long, 10 to 11 μ broad, 8 μ thick, polar capsule 11 to 14 μ by 2 to 3 μ . Cysts oval, numerous and small (130 μ by 115 μ)
Henneguya minuta Cohn 1895.....(160)
- 21(20) Tail about three-sevenths of main part; total length 29 to 40 μ , main part 15 to 20 μ long, 7 to 8 μ broad, polar capsule 8 μ by 2 μ 22
- 22(25) Cysts large..... 23

*Length of main part of spore denotes in all possible cases the distance between the outer anterior tip and the posterior margin of sporal cavity; and consequently that between the latter and the distal end of the tail is the length of the tail.

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

- 23(24) Cysts spherical up to 6 mm. in diameter; parasitic in ovary
Henneguya oviperda (Cohn 1895) (160)
- 24(23) Cysts elongated up to 2 mm. by 1.5 mm.; parasitic in branchia
Henneguya psorospermica Thélohan 1895. (158)
- 25(22) Cysts rather small or size not observed. 26
- 26(27) Elongated cysts 0.75 mm. by 0.375 mm.; parasitic in branchia
Henneguya texta (Cohn 1895) (159)
- 27(26) Parasitic in intestinal wall
Henneguya peri-intestinalis Cépède 1906. (161)
- 28(11) Tail composed of two processes. 29
- 29(30) Total length reaches 87.5 to 110.5 μ ; length of main part 10.5 to 15 μ , breadth
5 μ , length of polar capsule 5 μ , length of tail 75 to 100 μ
Henneguya gigantea Nemeček 1911. (168)
- 30(29) Total length of spore less than 82 μ 31
- 31(34) Sutural edge with markings. 32
- 32(33) Tail longer and spore larger; anterior end more rounded; total length 47 μ . Cysts
spherical and large, up to 6 mm.
Henneguya salminicola Ward 1919. (171)
- 33(32) Tail shorter and spore smaller; anterior end slightly more attenuated; total
length 32 μ . Cysts lenticular, up to 2 mm. in length
Henneguya nüsslini Schuberg et Schröder 1905. (166)
- 34(31) Sutural edge without markings. 35
- 35(38) Distal end of tail thread-like. 36
- 36(37) Tail 40 to 50 μ in length; total length 50 to 60 μ , main part 8.5 to 9.5 μ long, 8.5 to
9.5 μ broad, polar capsule 4.7 to 5.5 μ by 3 μ . Cysts small, up to 50 μ in
diameter
Henneguya neapolitana Parisi 1912. (170)
- 37(36) Tail 10 to 30 μ in length; main part 12 μ long, 8 μ broad
Henneguya miyairii Kudo 1919 (172)
- 38(35) Distal end of tail tapers to a point and not thread-like. 39
- 39(40) Cysts irregular in shape; size up to 2.5 mm. Spore: total length 30 to 40 μ ,
main part 11.5 to 15 μ long, 5 to 6.5 μ broad, polar capsule 6.5 to 8 μ by 2 to
2.5 μ , tail 22 to 28 μ
Henneguya lobosa (Cohn 1895) (161)
- 40(39) Cysts spherical or oval. 41
- 41(42) Anterior end of spore rounded; tail either single or double processes; total length
55 μ , length of main part 10 μ , breadth 7 μ , length of tail 40 to 50 μ . Cysts
spherical or oval up to 32 mm. by 16 mm.
Henneguya zschokkei (Gurley 1893) (165)
- 42(41) Anterior end attenuated; spore large; main part 20 to 22 μ long, breadth 8 to
9 μ , 6 to 7 μ thick, polar capsule 10 μ by 2 to 3 μ , tail 50 to 60 μ long
Henneguya acerinae Schröder 1906. (167)
- Incompletely described species
Henneguya schizura (Gurley 1893) (162)
Henneguya linearis (Gurley 1893) (163)
Henneguya gurleyi Kudo 1893. (163)
Henneguya monura (Gurley 1893) (164)
Henneguya strongylura (Gurley 1894) (163)
Henneguya kolesnikovii (Gurley 1894) (164)
Henneguya brevis Thélohan 1895. (162)
Henneguya sp. (Gurley 1894) (165)
Henneguya sp. (Gurley 1894) (165)
Henneguya sp. Nemeček 1911. (169)

Numbers enclosed in parentheses refer to the page of the article on which is found the description of the species named.

SUMMARY

1) All species of Myxosporidia which have been observed in various parts of the world, reaching 237 in number, are recorded with figures.

2) A new classification of Myxosporidia is proposed after discussion of those of previous authors.

3) A complete list of the specific names of the hosts that harbor Myxosporidia, is given together with the names of the organ of infection and the localities from which recorded.

4) By study of the geographical distribution of Myxosporidia, it is shown that few species are common both to American and European waters or Asiatic and European waters, while the majority of Myxosporidia are localized in definite and limited regions.

5) The study of the organal distribution of Myxosporidia in the host, shows that the gall-bladder is the organ most frequently invaded by the parasite. The kidney, branchia and urinary bladder have less chances of being attacked.

6) One new genus, *Wardia*, is established.

7) Nine new species; *Wardia ovinocua*, *Mitraspora elongata*, *Chloromyxum trijugum*, *Chloromyxum catostomi*, *Myxidium americanum*, *Myxobolus orbiculatus*, *Myxobolus discrepans*, *Myxobolus mesentericus* and *Henneguya mictospora*, are described from fresh-water fish collected in the vicinity of Urbana, Ill.

8) Six new species; *Sphaerospora carassii*, *Myxidium kagayamai*, *Myxobolus misgurni*, *Myxobolus miyairii*, *Myxobolus koi* and *Henneguya miyairii*, are recorded from fresh-water fish of Nippon.

9) One new species; *Chloromyxum wardi*, is described from Alaska. This is the second species of Myxosporidia from that part of North America.

10) Keys to the genera and species of known Myxosporidia are included.

APPENDIX: NEW MYXOSPORIDIA FROM AUSTRALIA

The following six species described by Johnston and Bancroft did not reach the writer until the page proof was read. For this reason they could not be put in the text and are recorded here separately.

MYXIDIUM THERAPON Johnston et Bancroft

1919 *Myxidium therapon* Johnston and Bancroft 1919 : 520-521

Habitat: In the gall bladder of *Therapon carbo* and *Th. hillii*; Thomson River at Longreach, Australia.

The parasite occurred in one specimen of the former host fish and in nine out of thirteen specimens of the latter. No visible effect of the infection on the part of the host fish was recognized.

Vegetative form: Body pale yellowish to green in color. Form(?). Size varies from 3 to 12mm. in diameter. The protoplasm is differentiated into a clear narrow ectoplasm, about 10μ in width, and a coarsely grained endoplasm. No movements could be seen on slides; but undulations were observed to travel round the margin of the trophozoite. Polysporous.

Spore: Spindle-shaped with slightly pointed extremities. Polar capsules are more or less rounded. Shell with faintly marked longitudinal striation. The sporoplasm is binucleated. Average dimensions: length 9 to 10μ , breadth 4μ , polar capsules 2 to 3μ by 1 to 2μ .

MYXOSOMA OGILBYI Johnston et Bancroft

1919 *Myxosoma ogilbyi* Johnston and Bancroft 1919 : 521-522

Habitat: In the fibrous tissue of the gill arch of *Plectroplites ambiguus*; Thomson River at Longreach, Australia. Three out of nine host specimens examined showed the infection.

Vegetative form: The parasite forms white cysts usually close to the bases of the gill filaments. Cysts are small and rounded, being less than 1mm. in diameter. The authors simply mention that sections revealed the structure usually present in a Myxosporidian cyst.

Spore: Oval with pointed anterior end. The inner margin of the shell is indented posteriad. The sporoplasm contains a single nucleus, but not any iodophilous vacuole. Average dimensions: length 11 to 13μ , breadth 6 to 8μ , thickness 5μ , polar capsules 5 to 6μ by 2μ .

MYXOBOLUS PLECTROPLITES Johnston et Bancroft

1919 *Myxobolus plectroplites* Johnston and Bancroft 1919 : 522-523

Habitat: In the kidney and gall-bladder of *Plectroplites ambiguus*; Thomson River at Longreach, Australia. The parasite was observed in

four out of nine host fish examined; in two cases in the kidney only, in one case only in the gall-bladder, and in one instance in both gall-bladder and kidney. Cysts were found in the kidney, while only spores were recognized in the gall-bladder.

Vegetative form: The cysts which could only be detected in sections, lie in the connective tissues of kidney. They are of minute size, ranging somewhat widely from 36μ in diameter to 144 by 100μ . According to the authors no definite structure could be found.

Spore: Rounded oval. It bears quite a close resemblance to that of *Myxobolus hylae* (page 153), which is slightly longer, and which has a longer polar filament than the present form. The vacuole, however, is apparently not iodophilous(?). Average dimensions: length 10 to 12μ , breadth 7 to 8μ , polar capsules 5 by 2μ , length of polar filament 30 to 40μ .

HENNEGUYA AUSTRALIS Johnston et Bancroft

1919 *Henneguya australis* Johnston and Bancroft 1919 : 523-524

Habitat: In the branchiae of *Plectroplites ambiguus*; Thomson River at Longreach, Australia. The parasite was detected in four out of nine host fish examined. The infection was extremely light in all cases.

Vegetative form: The parasites form cysts. They lie embedded in the spongy mass of the gill filament, and in many cases occupy a relatively large area of the section. Cysts showed two layers in section; the outermost clear ectoplasm and inner endoplasm with developing spores, the central portion of which being filled with mature spores. The spores appear to lie in a definite manner, the long axis of the spore commonly being at right angles to the boundary of the cyst, the anterior end of the spore pointing outwards.

Spore: Elongated ovoidal. Anterior end pointed, posterior end drawn out into a tail. The tail appears single when the spore is removed from the cyst but separates soon afterward into two halves which usually diverge widely. Two polar capsules parallel to each other are quite frequently of different length. The sporoplasm contains two nuclei and a small vacuole (iodophilous?). Average dimensions: length 11 to 15μ , breadth 3 to 5μ , thickness 3 to 4μ , polar capsules 5 to 6μ by 1 to 2μ , length of tail about 20μ .

HENNEGUYA GRACILIS Johnston et Bancroft

1919 *Henneguya gracilis* Johnston and Bancroft 1919 : 524-526

Habitat: In the gill filament of *Therapon hillii*; Thomson River at Longreach, Australia. Out of thirteen specimens examined, eight were infected. Heavy infection was recognized only in one case.

Vegetative form: The cyst is of definite, narrow, pear-shaped form, and lie transversely, i.e., at right angles to the long axis of the gill filament.

Spore: The spore resembles *Henneguya australis*, but is slightly smaller, while the tail is longer in proportion. The spores are arranged with long axis parallel to that of the cyst. Average dimensions: length 10 to 14 μ , breadth 2.5 to 3 μ , thickness 3 μ , polar capsules 5 to 6 μ by 1 to 2 μ , length of tail about 20 to 26 μ .

HENNEGUYA sp. Johnston et Bancroft

1919 *Henneguya* sp.

Johnston and Bancroft 1919 : 526

Habitat: In the branchiae of *Nematalosa elongata*; Thomson River at Longreach, Australia.

The authors state that they observed a number of spores of a *Henneguya* in the scrapings of the gill of one of four host fish.

Vegetative form: Undescribed.

Spore: Undescribed.

BIBLIOGRAPHY

AUERBACH, M.

1906. Ein Myxobolus im Kopfe von *Gadus aeglefini* L. Zool. Anz., 30:568-570; 4 fig.
- 1906a. Weitere Mitteilungen über *Myxobolus aeglefini*. Zool. Anz., 31:115-119; 5 fig.
- 1906b. Eine neuer Myxobolus im Brachsen (*Abramis brama* L.). Zool. Anz., 31:386-391; 5 fig.
1908. Bemerkungen über Myxosporidien heimischer Süßwasserfische. Zool. Anz., 32:456-465; 7 fig.
1909. Bemerkungen über Myxosporidien. Zool. Anz., 34:65-82; 6 fig.
- 1909a. Bericht über eine Studienreise nach Bergen (Norway). Verh. naturw. Ver. Karlsruhe, 21:37-73; 2 pl.
1910. Biologische und morphologische Bemerkungen über Myxosporidien. Zool. Anz., 35:57-63; 3 fig.
- 1910a. Die Sporenbildung von *Zschokkella* und das System der Myxosporidien. Zool. Anz., 35:240-256; 5 fig.
- 1910b. Cnidosporidienstudien. Zool. Anz., 35:767-777; 4 fig.
- 1910c. Die Cnidosporidien (Myxosporidien, Actinomyxidien, Microsporidien). Eine monographische Studie. 261 pp.; 83 fig.
- 1910d. Zwei neue Cnidosporidien aus cyprinoiden Fischen. Zool. Anz., 36:440-441; 1 fig.
1911. Untersuchungen über *Henneguya psorospermica* Thélohan. Verh. naturw. Ver. Karlsruhe, 24:1-25; 2 textfig., 2 pl.
- 1911a. Unsere heutigen Kenntnisse über die geographische Verbreitung der Myxosporidien. Zool. Jahrb., Syst., 30:471-494.
1912. Studien über die Myxosporidien der norwegischen Seefische und ihre Verbreitung. Zool. Jahrb., Syst., 34:1-50; 5 textfig., 5 pl.
- 1912a. Bemerkungen über den Infektionsmodus der Seefische mit Myxosporidien. Zool. Anz., 39:617-623.
- 1912b. Die Sporenbildung der Myxosporidien. Zool. Anz., 40:204-207
1917. Bemerkungen über Myxosporidien. Zool. Anz., 49:145-157; 5 fig.

AWERINZEW, S.

1907. Ueber Myxosporidien aus Gallenblase der Fische. Zool. Anz., 31:831-834.
1908. Studies on parasitic Protozoa. I, II, and III (Russian). Trav. soc. imp. nat. St.-Petersbourg, 28 (fasc. 2), 67 pp.; 3 pl.
1909. Studien über parasitischen Protozoen. I. Die Sporenbildung bei *Ceratomyxa drepanopsettae* mihi. Arch. Protist., 14:74-112, 2 pl.
1911. Studien über parasitischen Protozoen. VII. Ueber die Sporenbildung bei *Myxidium* sp. aus der Gallenblase von *Cottus scorpius*. Arch. Protist., 23:199-204; 7 fig.
1913. *Myxobolus magnus* nov. sp. Zool. Anz., 42:75-76; 2 fig.
- 1913a. Ergebnisse der Untersuchungen über parasitische Protozoen der tropischen Region Afrikas. III. Zool. Anz., 42:151-156; 4 fig.

BALBIANI, G.

1863. Sur l'organisation et la nature des psorospermies. C.R. acad. sci., 57:157-161.
1883. Myxosporidies ou psorospermies des poissons. Journ. Microgr., 7:143-147, 197-204, 270-281; 12 fig.
1884. Leçons sur les Sporozoaires. IV. Les psorospermies des poissons ou Myxosporidies, p. 420-449; 12 fig.

BERG,

*1898. *Communicat. Mus. Nac., Buenos Aires, p. 41.*

BORNE, M. VON DEM, B. BENECKE and E. DALLMER.

*1886. *Handbuch der Fischzucht und Fischerei. 701 pp., 581 fig. Berlin.*

BOSANQUET, W. C.

1910. *Brief notes on two Myxosporidian organisms (Pleistophora hippoglossoides n. sp. and Myxidium mackiei n. sp.). Zool. Anz., 35:434-438; 13 fig.*

BÜTSCHLI, O.

1881. *Beiträge zur Kenntnis der Fischsporospermien. Zeitschr. wiss. Zool., 35:629-651; 1 pl.*

1882. *Myxosporidia. Bronn's Klass. Ordn., Protozoa, 1:590-603; 19 fig.*

CÉPÈDE, C.

1905. *Myxosporidies des poissons des Alpes françaises. C. R. ass. fr. l'avanc. sc. sess., 33:905-913.*

1906. *Myxosporidies des poissons des Alpes françaises. Ann. l'univ. Grenoble, 18: 57-68.*

1906a. *Sur la prétendue immunité des Cobitis à l'égard des infections myxosporidiennes. C.R. soc. biol., 60:15-16.*

1906b. *Myxidium giardi Cépède, et la prétendue immunité des Anguilles à l'égard des infections myxosporidiennes. C.R. soc. biol., 60:170-173; 4 fig.*

1907. *A propos de la déhiscence des spores des Myxosporidies. C.R. soc. biol., 62:135-137.*

1908. *La myxosporidiose des Anguilles dans les eaux douces, saumâtres et salées du Boulonnais. Feuille jenn. natur. (4) 38:93-95; 6 fig.*

1913. *Existence de la plasmotomie hivernale chez Henneguya legeri Cépède. Arch. Parasit., 16:302-305; 26 fig.*

CLAPARÈDE, A. DE

*1874. *Notice sur les psorospermies des poissons. Lunel's Hist. natur. des poissons du bassin du Léman. Geneve, p. 113-114.*

CLELAND, J. B. and JOHNSTON, T. H.

1910. *The Haematozoa of Australian Batrachians. Proc. Roy. Soc. N.S.W., 44: 252-261.*

COHN, L.

1895. *Ueber die Myxosporidien von Esox lucius und Perca fluviatilis. Inaugural-Dissert. Albertus-Univ. Königsberg, 48 pp.; 2 pl.*

1896. *Ueber die Myxosporidien von Esox lucius und Perca fluviatilis. Zool. Jahrb., Anat., 9:229-272; 2 pl.*

1902. *Zur Kenntnis der Myxosporidien. Centralb. Bakt. Parasit., (I) Orig. 32:628-632; 2 fig.*

CREPLIN, J. C. H.

1842. *Beschreibung der Psorospermien des Kaulbarsches nebst einigen Bemerkungen über die der Plötze und andere. Arch. Naturg., 8:61-66; 1 pl.*

CUNHA, A. M. DA and O. DA FONSECA.

1917. *Sobre os myxosporidios dos peixes brasileiros. Brazil-Medico, 31:321.*

DANILEWSKY, B.

1887. *Recherches sur la parasitologie du sang. Arch. slav. biol., 3:35.*

DAVIS, H. S.

1916. *The structure and development of a Myxosporidian parasite of the squeteague, Cynoscion regalis. Journ. Morph., 27:333-377; 7 textfig., 7 pl.*

1917. *The Myxosporidia of the Beaufort Region. A systematic and biologic study. Bull. Bur. Fish., 35:201-243; 8 pl.*

*Original paper not seen.

- DE DROUIN DE BOUVILLE, R.
 1908. Maladie des abcès du Barneau (*Myxoboliasis tuberosa*). Bull. soc. sc. Nancy (3) 9:525-548; 9 textfig., 1 pl.
- DOFLEIN, F.
 1898. Studien zur Naturgeschichte der Protozoen. III. Ueber Myxosporidien. Zool. Jahrb., Anat., 11:281-350; 20 textfig., 7 pl.
 1899. Fortschritte auf dem Gebiete der Myxosporidienkunde. Zusammenfassende Uebersicht. Zool. Centr., 6:361-379.
 1901. Die Protozoen als Parasiten und Krankheitserreger nach biologischen Gesichtspunkten dargestellt. Jena.
 1902. Das System der Protozoen. Arch. Protist., 1:169-192.
 1909. Die Lehrbuch der Protozoenkunde. 2 ed.
 1911. Die Lehrbuch der Protozoenkunde. 3 ed.
 1916. Die Lehrbuch der Protozoenkunde. 4 ed.
- ERDMANN, RH.
 1911. Zur Lebensgeschichte von *Chloromyxum leydigi*, eine miktosporen Myxosporidie. Teil I. Arch. Protist., 24:149-162; 4 textfig., 3 pl.
 1917. New facts and views concerning the occurrence of a sexual process in the Myxosporidian life cycle. Amer. Natur., 51:719-739.
 1917a. *Chloromyxum leydigi* und seine Beziehungen zu anderen Myxosporidien. Teil II. Arch. Protist., 37:276-326; 17 textfig. and 4 pl.
- EVERMANN, B. W.
 1893. A report upon investigations made in Texas in 1891. Bull. U. S. Fish. Comm., 11:76.
- FANTHAM, H. B. and ANNIE PORTER,
 1912. Some effects of the occurrence of Myxosporidia in the gall-bladder of fishes (Prelim. Communic.). Ann. Trop. Med. Parasit., 6:467-481.
- FERMOR, X.
 1913. Wissenschaftliche Ergebnisse einer Reise von S. Awerinzew in die Tropen Afrikas. Zool. Anz., 42:196-199.
- FICKERT, K.
 1895. Ueber die Barbenseuche. Zeitschr. Fischer., 3:209-214.
- FIEBIGER, J.
 1909. Ueber Protozoen als Parasiten der Fische. Verh. zool.-bot. Ges. Wien. 59: 32-48.
- FLETCHER, A. W.
 1888. On a Myxosporidium infesting Australian frogs. Rep. Austr. Ass. Adv. Sc., 1:337.
- FRITSCH, A. J.
 *1902. Ueber Lebensweise, Nahrung und Parasiten der Fische der Elbe. Arch. naturw. Landesforsch. Prag, 11 (No. 3).
- FUHRMANN, O.
 1902. Sur les Myxosporidies des Coregones du lac de Neuchâtel. Arch. sc. phys. nat. Genève, 14:172-173, 16:331-332.
 1904. Ueber eine Krankheit der weiblichen Geschlechtsorgane des Hechtes. Allg. Fischer.-Zeit., 29:469-471.
- FUJITA, T.
 1912. Notes on new Sporozoan parasites of fishes. Zool. Anz., 39:259-262; 3 fig.
 1913. On a new species of *Chloromyxum* from the gall-bladder of the carp. Annot. Zool. Japon., 8:257-259; 1 fig.
- GEORGÉVITCH, J.
 1914. Sur le cycle évolutif chez les myxosporidies. C.R. acad. sci., 158:190-192.

*Original paper not seen.

- 1914a. Étude de cycle évolutif chez les Myxosporidies. Arch. zool. exp., 54:387-409; 3 pl.
1916. Note sur les Myxosporidies recueillies à Roscoff. Bull. soc. zool. France, 41:86-95.
- 1916a. Note sur les Myxosporidies des poissons de la baie de Villefranche et de Monaco. Bull. l'inst. ocean., no. 322; 5 fig.
- 1916b. Sur les diverses formes de *Ceratomyxa herouardi* Georgévitch. C.R. acad. sci., 163:717-719.
- 1916c. Sur le cycle évolutif de *Ceratomyxa herouardi* Georgévitch. C.R. acad. sci., 163:983-985.
1917. Recherches sur le développement de *Ceratomyxa herouardi* Georgévitch. Arch. zool. exp., 56:375-399; 3 pl.
- 1917a. Esquisses protistologiques. Bull. l'inst. océan, no. 328; 50 fig.
- 1917b. Esquisses protistologiques. Bull. soc. zool. France, 42:99-107.
- 1917c. Sur le cycle évolutif de *Myxidium gadi* Georgévitch. C.R. acad. sci., 165:797-799.
1918. Sur *Chloromyxum leydigi*. Bull. soc. zool. France, 42:182-189; 18 textfig.
1919. Études sur le développement de *Myxidium gadi* Georgévitch. Arch. zool. exper., 58:251-289; 3 pl. and 4 textfig.
- GLUGE, G.
1841. Beobachtungen zahlreicher Balggeschwulste als epidemische Krankheit bei Fischen. Anat.-Mikr. Unters. allg. u. sp. Path. 2:202-204; 1 pl.
- GURLEY, R. R.
1893. On the classification of Myxosporidia, a group of protozoan parasites infesting fishes. Bull. U. S. Fish Comm., 11:407-420.
1894. The Myxosporidia, or psorosperms of fishes, and the epidemic produced by them. Rep. U. S. Fish Comm., 26:65-304; 47 pl.
- HAHN, C. W.
1915. Sporozoon parasites of certain fishes in the vicinity of Woods Hole, Mass. Bull. Bur. Fish., 33:193-214; 2 pl.
1917. On the Sporozoon parasites of the fishes of Woods Hole and vicinity. I. Further observations on *Myxobolus musculi* from Fundulus. Journ. Parasit., 3:91-104; 3 fig.
- 1917a. On the Sporozoon parasites of the fishes of Woods Hole and vicinity. II. Additional observations upon *Myxobolus musculi* of Fundulus and a nearly related species, *Myxobolus pleuronectidae* of *Pleuronectes americanus*. Journ. Parasit., 3:150-162; 1 pl.
- 1917b. On the Sporozoon parasites of the fishes of Woods Hole and vicinity. III. On the *Chloromyxum clupeidae* of *Clupea harengus* (Young), *Pomolobus pseudo-harengus* (Young), and *P. aestivalis* (Young). Journ. Parasit., 4:13-20.
- HARTMANN, M.
1909. Autogamie bei Protisten und ihre Bedeutung für das Befruchtungsproblem. Arch. Protist., 14:286-292.
- HASWELL, W. A.
1890. A remarkable flat worm in the golden frog. Proc. Linn. Soc. N.S.W., 5:661-666.
- HECKEL, J. and R. KNER.
- *1858. Die Süßwasserfische der oestereichischen Monarchie mit Rücksicht auf die angrenzenden Länder. Leipzig. 388 pp., 204 fig.
- HOFER, B.
1893. Ueber die Drehkrankheit der Regenbogenforelle. Allg. Fischereiz., 18:7-8.
1895. Ueber Fischkrankheiten. Zeitschr. Fisch., 3:179-209; 23 fig.
1896. Die sogenannte Pockenkrankheit der Karpfen. Allg. Fischereiz., 21:2, 186, etc.
- 1896a. Die Infektion der Fische mit Myxosporidien. Allg. Fischereiz., 21:38-39.
1904. Handbuch der Fischkrankheiten. Stuttgart. 359 pp., 222 fig.
- *Original paper not seen.

- HUBER, J. C.
1888. Ueber Piesbergens Fischsporospermien. Centralbl. Bakt. u. Parasit., 3:663-664.
- ISHII, S.
1915. Myxosporidiosis of Nipponese eel (Nipponese). Jour. Zool., 27:372-382; 1 pl.
1915a. Lentospora-disease of the eel (Nipponese). Jour. Zool. 27:471-474; 4 fig.
- JAMESON, P.
1913. A note on some Myxosporidia collected at Monaco. Bull. l'inst. océan., no. 273.
- JOHNSTON, T. H.
1909. Exhibit of Myxobolus. Proc. Roy. Soc. N.S.W., 43:29.
- JOHNSTON, T. H. and M. J. BANCROFT.
1918. A parasite, *Myxobolus hylae* sp. nov., of the reproductive organs of the golden swamp frog, *Hyla aurea*. Australian Zoologist, 1:171-175; 5 textfig.; 1 pl.
1919. Some new sporozoan parasites of Queensland freshwater fish. Jour. Proc. Roy. Soc. N.S.W., 52:520-528; 5 pl.
- JOHNSTONE, J. and H. M. WOODCOCK.
1907. On a Myxosporidian infection of *Gadus esmarkii*, with a note on the identification of the parasite. Trans. Biol. Soc. Liverpool, 21:204-208; 1 pl.
- JOSEPH, H.
1905. *Chloromyxum protei* n. sp. Zool. Anz., 29:450-451.
1907. *Chloromyxum protei* n. sp., ein in der Niere des Grottenolms parasitierendes Myxosporidium. Arch. Protist., 8:398-412; 1 textfig., 2 pl.
- KEYSSELITZ, G.
1908. Ueber durch Sporozoen (Myxosporidien) hervorgerufene pathologische Veränderungen. Verh. Ges. deutsch. Natur. u. Ärzte, Vers. 79:452-453.
1908a. Die Entwicklung von *Myxobolus pfeifferi*. Teils I, II, Arch. Protist., 11:252-308; 14 textfig., 4 pl.
- KLOKACEWA, S.
1914. Ueber die Myxosporidien der Karausche. Zool. Anz., 44:182-186; 2 fig.
- KOLESNIKOFF, N. F.
*1886. O psorospermiakh (mikrosporidiakh) v muskulature rib. Vet. Vestnik, Kharkoff, 5:242-248; 1 pl.
- KUDO, R.
1915. On a new Myxosporidian from a carp (Nipponese). Journ. Zool., 27:517-523; 2 pl.
1916. Contributions to the study of parasitic protozoa. III, Notes on some Myxosporidia found in some fresh-water fishes of Japan, with the description of three new species. Journ. Parasit., 3:3-9; 4 fig.
1917. Contributions to the study of parasitic protozoa. II. *Myxobolus toyamai* nov. spec., a new Myxosporidian parasite in *Cyprinus carpio* L. Journ. Parasit., 3:163-170; 2 pl.
1918. Experiments on the extrusion of polar filaments of Cnidosporidian spores. Journ. Parasit., 4:141-147; 1 pl.
1918a. Contributions to the study of parasitic protozoa. IV. Note on some Myxosporidia from certain fish in the vicinity of Woods Hole. Journ. Parasit., 5:11-16; 4 textfig., 2 pl.
- LABBÉ, A.
1899. Sporozoa. Das Tierreich. 5 Lief. 180 pp.
- LAVERAN, A.
1897. Sur une Myxosporidie des reins de la tortue. C.R. soc. biol. 49:725-726.
1898. Sur le *Myxidium danilewskyi*. C.R. soc. biol., 50:27-30; 9 fig.
- LAVERAN, A. and F. MÉSNIL.
1900. Sur une Myxosporidie des voies biliaires de l'Hippocampe (*Sphaeromyxa sabrazesi* nov. spec.). C.R. soc. biol., 52:380-382; 4 fig.

*Original paper not seen.

1902. Sur la multiplication endogène des Myxosporidies. C.R., soc. biol., 54:469-472; 5 fig.
- LEBZELTER, V.
1912. Ueber Protozoen aus der Gallenblase von *Thymallus thymallus* L. Zool. Anz., 40:295-297.
- LÉGER, L.
1906. Myxosporidies nouvelles parasites des poissons. I. Sur une nouvelle maladie myxosporidienne de la truite indigne. II. Sur une nouvelle Myxosporidie de la tanche commune. Ann. l'univ. Grenoble, 18:267-272; 3 fig.
- LÉGER, L. and E. HESSE.
1906. Sur la structure de la paroi sporale des Myxosporidies. C.R. acad. sci., 142: 720-722; 8 fig.
- LEUCKART, R.
1852. Parasitismus und Parasiten. Arch. physiol. Heilkde, 11:434-436.
1879. Die Parasiten des Menschen. 336 pp.; 130 fig.
- LEYDIG, F.
1851. Ueber Psorospermien und Gregarinen. Arch. Anat., Phys. u. wiss. Med., 18: 221-234; 1 pl.
- LIEBERKÜHN, N.
1854. Notice sur les psorospermies. Bull. ac. roy. Belg., 21:21-23.
1855. Les psorospermies des poissons. Mém. cour. et mém. sav. étrang. acad. roy. Belg., 26:36-38; 2 pl.
- LINTON, E.
1891. On certain wart-like excrescences, occurring on the short minnow, *Cyprinodon variegatus*, due to psorosperms. Bull. U.S. Fish Comm., 9:99-102; 1 pl.
1891a. Notice of the occurrence of protozoan parasites (psorosperms) on Cyprinoid fishes in Ohio. Bull. U. S. Fish Comm., 9:359-361; 1 pl.
1901. Parasites of fishes of the Woods Hole region. Bull. U. S. Fish Comm., 19:407-492; 34 pl.
- LO GIUDICE,
1912. Studi sui Cnidosporidi. Pavia, 91 pp. ; 1 pl.
- LUDWIG, H.
*1888. Ueber die Myxosporidienkrankheit der Barben in der Mosel. Jahresb. rhein. Fisch. Ver., Bonn, p. 27-36.
- LÜHE, M.
1896. Die Infektion der Fische mit Myxosporidien. Die Natur, 45:284-285; 5 fig.
1899. *Cystodiscus immersus*. Verh. deutsch. zool. Ges., 9:291-293; 1 fig.
- LUTZ, A.
1889. Ueber eine Myxosporidium aus der Gallenblase brasilianischer Batrachier (*Cystodiscus immersus*). Centralbl. Bakt. u. Parasit., 5:84-88; 10 fig.
- MAVOR, J. W.
1915. Studies on the Sporozoa of the fishes of the St. Andrew's Region. Ann. Rep. Dept. Marine Fishery, 47, no. 39b:25-38; 6 textfig.; 1 pl.
1916. On the life-history of *Ceratomyxa acadensis*, a new species of Myxosporidia from the eastern coast of Canada. Proc. Amer. Acad. Arts and Sci., 51:551-574; 3 textfig., 3 pl.
1916a. Studies on the protozoan parasites of the fishes of the Georgian Bay. Trans. Roy. Soc. Canada., (3) 10:63-74; 6 fig.
1916b. On the occurrence of a parasite of the pike in Europe, *Myxidium lieberkühni* Bütschli, in the pike on the American continent and its significance. Biol. Bull., 31:373-378; 1 pl.
- *Original paper not seen.

- MAVOR, J. W., and W. STRASSER.
 1916. On a new Myxosporidian, *Henneguya wisconsinensis* n. sp., from the urinary bladder of the yellow perch, *Perca flavescens*. Trans. Wisc. Acad. Sci., Arts and Letters, 18:676-682; 3 fig.
- MAZZARELLI, G.
 1905. Sulla pseudodiferite degli Agoni. Atti soc. ital. sc. nat. mus. civ. stor. nat. Milano, 44:71-72.
 1906. Le condizioni della pesca nella provincia di Milano. Riv. mens. pesca, Milano, 8:301-325.
 *1908. Di alcune malattie di pesci e gamberi osservate in Lombardia. Atti terzo congresso naz. di pesca, Milano, p. 272.
- MÉGNIN, T. P.
 1885. Sur le rôle pathogénique de certaines psorospermies. Bull. soc. zool. France, 10:351-352.
- MERCIER, L.
 1906. Phénomènes de sexualité chez *Myxobolus pfeifferi*. C.R. soc. biol., 110:427-428.
 1906a. Contribution à l'étude du développement des spores chez *Myxobolus pfeifferi*. C. R. soc. biol., 110:763-764.
 1908. Notes sur les Myxosporidies. Arch. zool. exp., 8:LIII-LXII; 5 figs.
 1909. Contribution à l'étude de la sexualité chez les Myxosporidies et chez les Microsporidies. Mém. class. sc. l'acad. roy. Belg., 2:3-30; 1 pl.
- MILEWSKI, A.
 1917. Myxoboliasis tuberosa, die Barbenseuche oder Beulkrankheit der Barbe. Wochenschr. Aquar.-Terrar.-Kde., 14:14-17.
- MINCHIN, E. A.
 1912. An Introduction to the study of the protozoa. 517 pp.; 194 textfig.
- MINGAZZINI, P.
 1890. Sullo sviluppo dei Myxosporidi. Boll. soc. nat. Napoli, 4:160-164; 1 pl.
- MIVAIRI, K.
 1909. Guide to the study of parasitic protozoa (Nipponese). 170 pp.; 8 pl.
- MOROFF, T.
 1906. Ueber die Gelbsucht der Bachforelle. Osterr. Fischereiz. 3:285.
- MÜLLER, J.
 1841. Ueber eine eigenthümliche krankhafte parasitische Bildung mit specifisch organisierten Samenkörperchen. Arch. Anat. Phys. u. wiss. Med., 5:477-488; 1 pl.
 1841a. Krankhafter Hautausschlag mit specifisch organisierten Körperchen (Psorospermien). Ber. preuss. Akad. Wiss. Berlin, p. 212-221, 246-250, 232 pp., 4 pl.
- MULSOW, K.
 1911. Ein neuer Gehirnparasit des Karpfens. Allg. Fisch. Zeit., 36:483-485; 3 fig.
- NEMECZEK, A.
 1911. Beiträge zur Kenntnis der Myxo- und Microsporidien der Fische. Arch. Protist., 22:143-169; 19 textfig., 2 pl.
- NUFER, W.
 1905. Die Fische des Vierwaldstättersees und ihre Parasiten. Inaug.-Diss. Basel, 230 pp.; 21 fig.
 *1905a. Die Parasiten der Fische des Vierwaldstättersees. Festschr. 50-jähr. Besth. naturf. Gesellsch., Luzern, pp. 65-232; 4 pl.
- OHLMACHER, A.P.
 1893. Myxosporidia in the common toad, with preliminary observations on two chromophile substances in their spores. Jour. Amer. Med. Assoc., 20:561-567; 1 pl.
- PARISI, B.
 1910. *Sphaerospora caudata* n. sp. Zool. Anz., 36:253-254; 3 figs.
 *Original paper not seen.

1912. Primo contributo alla distribuzione geografica dei missosporidi in Italia. *Atti soc. ital. sc. nat.*, 50:283-290; 11 fig.
1913. Sulla *Sphaerospora caudata* Parisi. *Atti soc. ital. sc. nat.* 51:5-12; 1 pl.
- PERUGIA, A.
 *1890. Sulle Myxosporidie dei pesci marini. *Boll. scient. Pavia*, 12:134-139.
 *1891. Sulle Myxosporidie dei pesci marini. *Boll. scient. Pavia*, 13:22-25; 1 pl.
- PFEIFFER, L.
 1890. Die Protozoen als Krankheitserreger. 1 ed. 100 pp.; 34 fig.
 1891. Die Protozoen als Krankheitserreger. 2 ed. pp. 216; 91 fig.
 1893. Die Parasitismus des Epithelcarcinoms sowie der Sarco-, Micro- und Myxosporidien in Muskelgewebe. *Centralbl. Bakt. u. Parasit.*, 14:118-130.
 1895. Die Protozoen als Krankheitserreger. *Nachträge. Jena.* pp. 122; 52 fig.
- PIESBERGEN, F.
 1886. Eine neue Form von Fischpsorospermien aus *Perca fluviatilis*. *Jahresb. Ver. vater. Naturk. Württemberg*, 42:73.
- PLEHN, M.
 1904. Woher kommt die Drehkrankheit der Salmoniden? *Allg. Fischereiz.*, 29:151-153.
 1904a. Weiteres über die Drehkrankheit. *Allg. Fischereiz.*, 29:183-184.
 1905. Ueber die Drehkrankheit der Salmoniden (*Lentospora cerebralis* [Hofer] Plehn). *Arch. Protist.*, 5:145-166; 7 textfig., 1 pl.
 1906. Die Drehkrankheit der Salmoniden. *Allg. Fischereiz.*, 31:465-470.
 1906a. Nochmals über die Drehkrankheit der Salmoniden. *Allg. Fischereiz.*, 31:516-519.
 1910. Die pathogene Bedeutung der Myxoboliden für die Fische. *Sitz. Ges. Morph. u. Physiol. München*, 26:20-27; 3 fig.
- POCHE, F.
 1913. Das System der Protozoen. *Arch. Protist.*, 30:125-321; 1 fig.
- PRENANT, A.
 1902. Striation et ciliation de la partie adhérente du *Myxidium lieberkühni* Bütschli. *C.R. soc. biol.*, 54:844-846.
 1902a. Notes cytologiques. VII. Contribution a l'étude de la ciliation. Striation et ciliation de la partie adhérente du *Myxidium lieberkühni*. *Arch. anat. micr.*, Paris, 5:212; 7 textfig.
- RAILLIET, A. *
 1886. Maladie des Barbeaux causée par des psorospermies. *Bull. Mém. soc. centr. méd. vét. Paris*, 4:134-137.
 1890. La maladie des Barbeaux de la Marne. *Bull. soc. centrale d'aquic. Paris*, 2:117-120.
- REMAK, R.
 1852. Ueber runde Blutgerinsel und über pigmentgukelhaltige Zellen. *Arch. Anat., Phys. u. wiss. Med.*, 144-146; 1 pl.
- REUSS, H.
 1906. Neue Myxosporidien von Süßwasserfischen. *Bull. l'acad. imp. sc. St.-Petersbourg*, 25:199-205; 1 pl.
- RYDER, J. A.
 1880. The psorosperms found in *Aphredoderus sayanus*. *Amer. Nat.*, 14:211-212; 2 figs.
- SANDEMAN, G.
 1894. On the multiple tumors in Plaice and Flounders. *Ann. Rep. Scott. Fish. Board*, 11:391-392; p. 391-392.
- *Original paper not seen.

SCHRÖDER, O.

1906. Eine neue Myxosporidienart aus den Kiemen von *Acerina cernua* (*Henneguya acerinae* n. sp.). Arch. Protist., 7:186-196; 1 pl.
1907. Beiträge zur Entwicklungsgeschichte der Myxosporidien. *Sphaeromyxa sabrazesi* Laveran et Mesnil. Arch. Protist., 9:359-381; 3 textfig., 2 pl.
1910. Ueber die Anlage der Sporocyste (Pansporoblast) bei *Sphaeromyxa sabrazesi* Laveran et Mesnil. Arch. Protist., 19:1-5; 10 fig.
1912. Myxosporidien. Prowazek's Handbuch Path. Protoz., 1:324-336.

SCHUBERG, A. and O. SCHRÖDER.

1905. Myxosporidien aus dem Nervensystem und der Haut der Bachforelle (*Myxobolus neurobius* n. sp. and *Henneguya nüsslini* n. sp.). Arch. Protist., 6:45-60; 1 pl.

SOUTHWELL, T.

1915. Notes from the Bengal Fisheries Laboratory, Indian Museum. On some Indian Parasites of Fish with a note on Carcinoma in Trout. Rec. Ind. Mus., 11:311-330; 2 pl.

SOUTHWELL, T. and B. PRASHAD.

1918. On some Indian Myxosporidia. Rec. Ind. Mus., 15:344-348; 1 pl.

STAZZI, P.

1906. Psorospermosi o myxoboliosi di Barbi. Riv. mens. pesca., Milano, 8:14-19; 3 fig.

SURBECK, G.

1900. Eine neue Krankheit beim Bachsaibling (*Salvelinus fontinalis*). Allg. Fischereiz., 25:367-368.
1911. Eine grosse Sporencyste von *Henneguya zschokkei*. Schweiz. Fisch.-Zeitg., 19:163-165; 1 fig.
1913. Ueber eine eigenartige Form des Auftretens von *Henneguya zschokkei* Gurley. Schweiz. Fisch.-Zeitg., 21:30-31.
1914. Beitrag zur Fischpathologie. I u. II. Schweiz. Fisch.-Zeitg., 22:296-299.

THÉLOHAN, P.

1889. Sur la constitution des spores des Myxosporidies. C.R. acad. sci., 109:919-922.
1890. Recherches sur le développement des spores chez les Myxosporidies. C.R. soc. biol., 2:602-604.
- 1890a. Nouvelles recherches sur les spores des Myxosporidies (structure et développement). C.R. acad. sci., 111:692-695.
- 1890b. Contribution à l'étude des Myxosporidies. Microgr., 2:193-213; 1 pl. Annal.
1891. Sur deux Sporozoaires nouveaux, parasites des muscles des poissons. C.R. soc. biol., 3:27-29.
1892. Observation sur les myxosporidies et essai de classification de ces organismes. Bull. soc. philom., 4:165-178.
- 1892a. Myxosporidies de la vésicule biliaire des poissons. C.R. acad. sci., 115:961-964, 1091-1094.
1893. Altération du tissu musculaire due à la présence de Myxosporidies et de microbes. C.R. soc. biol., 5:267-270.
1894. Sur les affinités réciproques des Myxosporidies. C.R. acad. sci., 118:428-430.
1895. Recherches sur les Myxosporidies. Bull. sc. France et Belg., 26:100-394; 6 fig., 3 pl.

TROJAN, E.

1909. Ein Myxobolus im Auge von *Leuciscus rutilus*. Zool. Anz., 34:679-682; 3 figs.

TYZZER, E. E.

1900. Tumors and Sporozoa in fishes. Journ. Boston Soc. Med. Sc., 5:62-68; 1 pl.

VANEY, C. and A. CONTE,

1901. Sur deux nouveaux Sporozoaires endospores parasites de l'*Acerina cernua* Cuv. Ann. Soc. Linn. Lyon, 47: 103-106; 4 fig.

- WARD, H. B.
1919. Notes on North American Myxosporidia. Journ. Parasit., 6:49-64, 1 pl.
- WASIELEWSKY, VON
1896. Sporozoenkunde. Jena. 162 pp.; 111 textfig.
- WEGENER, G.
1910. Die Ektoparasiten der Fische Ostpreussens. Inaugur.-Dissert. Königsberg, p. 72-90; 4 figs.
- WELTNER, W.
1892. Ueber Myxosporidien Sporen in den Eiern von *Esox lucius*. Sitz. Ges. nat. Fr. Berlin, p. 7-41; 16 fig.
- WHINERY, J. B.
1893. Some additional notes on a Myxosporidian infection in the common toad. N. Y. Med. Journ., 18:660-662; 1 fig.
- WOODCOCK, H. M.
1904. On Myxosporidia in flat-fish. Trans. Biol. Soc. Liverpool, 18:46-62; 1 pl.
1907. On a Myxosporidian infection of *Gadus esmarkii*. With Johnstone.
- ZSCHOKKE, F.
1884. Psorospermies de *Coregonus fera*. Arch. biol. gand, Leipzig et Paris, 5:234-235; 1 pl.
1898. Die Myxosporidien der Gattung *Coregonus*. Centralbl. Bakt. u. Parasit., Abt. I. Orig., 23:602-607, 646-655, 699-703; 4 fig.
1898a. Die Myxosporidien in der Muskulature der Gattung *Coregonus*. Zool. Anz., 21:213-214.
1898b. *Myxobolus bicaudatus* n. sp., ein Parasit der Coregoniden des Vierwaldstättersees. Mitt. Naturf. Ges., Luzern, p. 205-217.
1900. *Myxobolus psorospermicus* Thélohan im Vierwaldstättersee. Mitt. Naturf. Ges. Luzern, p. 441-442.
- ZSCHOKKE, F. and A. HEITZ.
1914. Entoparasiten aus Salmoniden von Kamtschatka. Revue suisse zool., 22:195-256.

GENERAL EXPLANATION OF FIGURES

For the type species of each genus, both the vegetative form and the spore are illustrated. For the other species, except those which are new, figures of the spore are given, unless the vegetative forms are different from those of the type species or the species were reported in papers which seem to be of less universal distribution.

The original drawings were made with the Abbe drawing apparatus. The combinations used were Zeiss apochromatic objectives 16, 8, 3, and homogeneous oil immersion 2 mm. with compensation oculars 2, 4, 6, 8, 12 and 18. All the other drawings were copied from the original figures of the respective observers, an exact citation of which is given in each case, and were also made with the same drawing apparatus on the same scale except that a few figures were enlarged among those that were taken from other authors.

Magnifications were also calculated and given for those quoted figures, for which the respective authors failed to mention the scale at which the drawings were made.

PLATE I

EXPLANATION OF PLATE

Figs. 1 to 5. *Leptotheca agilis*.

Fig. 1. A typical trophozoite *in vivo*. After Thélohan (1895, Fig. 29); $\times 750$.

Fig. 2. A young form. After Thélohan (1895, Fig. 31). $\times 750$.

Fig. 3. A trophozoite in motion. After Doflein (1898, Fig. 5).

Fig. 4. A trophozoite in contracted condition. After Doflein (1898, Fig. 7).

Fig. 5. A fresh spore. After Thélohan (1895, Fig. 30). $\times 1500$.

Fig. 6. A fresh mature spore of *Leptotheca elongata*. After Thélohan (1895, Fig. 38). $\times 1500$.

Fig. 7. A fresh mature spore of *Leptotheca parva*. After Thélohan (1895, Fig. 25). $\times 1500$.

Fig. 8. A fresh spore of *Leptotheca perlata*. After Balbiani (1884, Fig. 40).

Fig. 9. A spore of *Leptotheca macrospora*. After Auerbach (1909, Fig. 2a). $\times 1350$.

Fig. 10. A spore of *Leptotheca informis*, preserved in formol. After Auerbach (1910b, Fig. 1a). \times about 2000.

Fig. 11. A spore of *Leptotheca longipes*, preserved in formol. After Auerbach (1910b, Fig. 1d). \times about 2200.

Fig. 12. A fresh spore of *Leptotheca fusiformis*. After Davis (1917, Fig. 1). $\times 1500$.

Fig. 13. A fresh spore of *Leptotheca scissura*. After Davis (1917, Fig. 8). $\times 1500$.

Fig. 14. A fresh spore of *Leptotheca lobosa*. After Davis (1917, Fig. 11). $\times 1500$.

Fig. 15. A fresh spore of *Leptotheca glomerosa*. After Davis (1917, Fig. 13). $\times 1500$.

Fig. 16. A trophozoite of *Leptotheca* sp. After Awerinzew (1908, Pl. 2, Fig. 14). 1/12 and comp. oc. 12.

Fig. 17. Another trophozoite of the same. After Awerinzew (1908, Pl. 2, Fig. 17). 1/12 and comp. oc. 12.

Figs. 18 to 20. *Ceratomyxa arcuata*. After Thélohan (1895).

Figs. 18 and 19. Typical young form. After Thélohan (1895, Figs. 16 and 17).

Fig. 20. A trophozoite with two spores. After Thélohan (1895, Fig. 18).




PLATE II

EXPLANATION OF PLATE

- Fig. 21. A sporulating trophozoite of *Ceratomyxa arcuata*. After Parisi (1912, Fig. 6a).
Fig. 22. A spore of *Ceratomyxa arcuata* treated with KOH. After Thélohan (1895, Fig. 19).
×1500.
Figs. 23 and 24. *Ceratomyxa sphaerulosa*. After Thélohan.
Fig. 23. A part of the trophozoite (1895, Fig. 2). ×750.
Fig. 24. A fresh spore (1895, Fig. 3). ×750.
Fig. 25. A fresh spore of *Ceratomyxa globulifera*. After Thélohan (1895, Fig. 43). ×1500.
Fig. 26. An adult trophozoite of *Ceratomyxa appendiculata*. After Thélohan (1895, Fig. 4).
× about 400.
Fig. 27. A spore of *Ceratomyxa truncata*. After Thélohan (1895, Fig. 51). ×1500.
Fig. 28. A spore of *Ceratomyxa reticularis*. After Thélohan (1895, Fig. 27). ×1500.
Fig. 29. A spore of *Ceratomyxa inaequalis*. After Doflein (1898, Fig. 10). ×1250.
Figs. 30 and 31. *Ceratomyxa linospora*. After Doflein (1898). × about 1900.
Fig. 30. A spore (1898, Fig. 39).
Fig. 31. A trophozoite with spores (1898, Fig. 43).
Figs. 32 and 33. *Ceratomyxa ramosa*. After Awerinzew (1908).
Fig. 32. A trophozoite (1908, Pl. 2, Fig. 20). Zeiss obj. D and comp. oc. 4.
Fig. 33. A spore (1908, Pl. 2, Fig. 19). Zeiss obj. E and comp. oc. 4.




PLATE III

EXPLANATION OF PLATE

- Figs. 34 to 39. *Ceratomyxa drepanopsettae*. Awerinzew (1908).
 Fig. 34 and 35. Trophozoites (1908, Pl. 2, Figs. 7 and 9). Obj. D and oc. 4.
 Fig. 36. The part of a trophozoite attached to the epithelium of the gall-bladder of the host (1908, Pl. 2, Fig. 10). Obj. E and oc. 4.
 Figs. 37 to 39. Spores (1908, Pl. 1, Figs. 2, 3 and 1). Obj. D and oc. 4.
 Figs. 40 and 41. Two different views of the spore of *Ceratomyxa tylosuri*. After Awerinzew (1913a, Fig. 1). \times about 350.
 Figs. 42 and 43. *Ceratomyxa(?) spari*. After Awerinzew (1913a, Fig. 2).
 Fig. 42. A trophozoite.
 Fig. 43. A spore. \times about 345.
 Figs. 44 to 47. Spores of *Ceratomyxa acadensis*. After Mavor (1916). Figs. 44 and 45 (1916, Fig. B). \times 270. Fig. 46 (1916, Fig. A) \times 1800. Fig. 47 (1916, Fig. 40) \times 2950.
 Fig. 48. A spore of *Ceratomyxa coris*. After Georgévitch (1916a, Fig. 1).
 Fig. 49. A spore of *Ceratomyxa herouardi*. After Georgévitch (1917, Fig. 1).
 Fig. 50. A spore of *Ceratomyxa mesospora*. After Davis (1917, Fig. 15). \times 1500.
 Fig. 51. A spore of *Ceratomyxa sphairophora*. After Davis (1917, Fig. 23). \times 950.
 Figs. 52 and 53. Spores of *Ceratomyxa taenia*. After Davis (1917, Figs. 26 and 25). \times 700.




PLATE IV

EXPLANATION OF PLATE

- Fig. 54. A spore of *Ceratomyxa attenuata*. After Davis (1917, Fig. 28). $\times 950$.
- Figs. 55 and 56. Spores of *Ceratomyxa recurvata*. After Davis (1917, Figs. 32 and 33). $\times 1500$.
- Figs. 57 to 60. Spores of *Ceratomyxa lunata*. After Davis (1917, Figs. 34 to 37). $\times 1500$.
- Fig. 61. A spore of *Ceratomyxa abbreviata*. After Davis (1917, Fig. 41). $\times 1500$.
- Fig. 62. A spore of *Ceratomyxa flagellifera*. After Davis (1917, Fig. 42). $\times 1500$.
- Fig. 63. A spore of *Ceratomyxa agglomerata*. After Davis (1917, Fig. 45). $\times 1500$.
- Fig. 64. A spore of *Ceratomyxa amorphia*. After Davis (1917, Fig. 47). $\times 1500$.
- Figs. 65 to 67. Spores of *Ceratomyxa monospora*. After Davis (1917, Figs. 57, 56 and 55). $\times 1500$.
- Figs. 68 and 69. Spores of *Ceratomyxa streptospora*. After Davis (1917, Figs. 59 and 60). $\times 1500$.
- Fig. 70. A spore of *Ceratomyxa aggregata*. After Davis (1917, Fig. 63). $\times 1400$.
- Fig. 71. A spore of *Ceratomyxa undulata*. After Davis (1917, Fig. 66). $\times 1500$.
- Figs. 72 and 73. Spores of *Ceratomyxa navicularia*. After Davis (1917, Figs. 69 and 68). $\times 1500$.
- Fig. 74. A spore of *Ceratomyxa spinosa*. After Davis (1917, Fig. 72). $\times 1500$.




PLATE V

EXPLANATION OF PLATE

- Figs. 75 to 80. *Myxoproteus ambiguus*. After Doflein (1898).
Figs. 75 and 76. Trophozoites of typical forms (1898, Figs. 12 and 52).
Figs. 77 and 78. Young trophozoites produced by budding (1898, Figs. 55 and 56).
Figs. 79 and 80. Spores (1898, Figs. 54 and 64). \times about 800 and 1080.
Figs. 81 to 83. Spores of *Myxoproteus cordiformis*. After Davis (1917, Figs. 79, 80 and 78).
 \times 1500.
Fig. 84. A spore of *Myxoproteus cornutus*. After Davis (1917, Fig. 85). \times 1400.
Figs. 85 to 95. *Wardia ovinocua*. Original.
Fig. 85. A portion of the cross-section thru an infected ovary of *Lepomis humilis*, showing the parasite in one ovum and the hypertrophied nurse cells and several connective tissue layers. \times 160.
Fig. 86. A portion of the cross-section of a cyst. \times 640.
Figs. 87 to 89. Three different views of fresh spore. \times 2000.
Figs. 90 and 91. Stained spores. \times 1700.
Fig. 92. A spore mechanically pressed and stained with Giemsa's mixture, showing the escaping polar capsules without extruding polar filament, and the sporoplasm. \times 1700.
Figs. 93 to 95. Front and lateral views of the shell valves, exhibiting the network-like fine ridges on the surface and the posterior processes. \times 1700.
Figs. 96 and 97. Spores of *Wardia ohlmacheri*. After Ohlmacher (1893, Figs. 4a and 2).
2mm. and oc. 4.




PLATE VI

EXPLANATION OF PLATE

- Figs. 98 to 104. *Mitraspora cyprini*.
- Figs. 98 and 99. Trophozoites from the ureter of *Cyprinus carpio in vivo*. Original. \times about 1500.
- Figs. 100 and 101. Different views of fresh spores. Original. \times about 2000.
- Figs. 102 to 104. Spores. After Fujita (1912, Figs. 1a to 1c).
- Figs. 105 to 107. *Mitraspora caudata*. After Parisi (1910 and 1913).
- Fig. 105. A trophozoite (1910, Fig. 1).
- Figs. 106 and 107. Front and lateral views of the spore (1913, Fig. 20 and 1910, Fig. 2). \times about 1600.
- Figs. 108 to 113. *Chloromyxum leydigi*.
- Figs. 108 and 109. Trophozoites. After Th  lohan (1895, Figs. 7 and 6). \times 750.
- Figs. 110 and 111. Trophozoites in division. After Doflein (1898, Figs. 57 and 58).
- Figs. 112 and 113. Different views of spores. After Th  lohan (1895, Figs. 10 and 9). \times 1500.
- Fig. 114. A spore of *Chloromyxum caudatum*. After Th  lohan (1895, Fig. 36). \times 1500.
- Figs. 115 and 116. Different views of the spore of *Chloromyxum quadratum*. After Th  lohan (1895, Figs. 100a and 100b). \times 1500.
- Fig. 117. A spore of *Chloromyxum quadratum* treated with nitric acid. After Th  lohan (1895, Fig. 100c). \times 1500.
- Fig. 118. A spore of *Chloromyxum fluviatile*. After Th  lohan (1892, Fig. 2). \times 1500.
- Figs. 119 and 120. Different views of the spore of *Chloromyxum mucronatum*. After Lieberk  hn from Gurley (1894, Pl. 39, Fig. 5). \times about 1750.
- Figs. 121 and 122. The same after Balbiani (1884, Fig. 41). \times about 1200.
- Figs. 123 and 125. Spores of *Chloromyxum diploxys*. After Balbiani from Gurley (1894 Pl. 42, Figs. 13a, 13b and 13c).




PLATE VII

EXPLANATION OF PLATE

- Fig. 126. Trophozoite of *Chloromyxum truttae*. After Léger (1906, Fig. 4). $\times 1000$.
- Figs. 127 and 128. Trophozoites of *Chloromyxum cristatum*. After Léger (1906, Fig. 7). $\times 1000$.
- Figs. 129 to 133. *Chloromyxum dubium*. After Auerbach (1908).
- Figs. 129 and 130. Trophozoites (1908, Figs. 2 and 1).
- Figs. 131 and 132. Spores (1908, Figs. 3 and 4). \times about 2250.
- Fig. 133. A stained young spore (1908, Fig. 5).
- Fig. 134. Trophozoite of *Chloromyxum* sp. After Awerinzew (1908, Pl. 2: Fig. 12). Objective D and ocular 4.
- Fig. 135. *Chloromyxum koi*. After Fujita (1913). \times about 800.
- Figs. 136 to 138. *Chloromyxum magnum*. After Awerinzew (1913a, Fig. 4).
- Figs. 136 and 137. Trophozoites.
- Fig. 138. A spore. \times about 320.
- Figs. 139 and 140. Spores of *Chloromyxum funduli*. After Hahn (1915, Figs. 34 and 33). $\times 2000$.
- Figs. 141 to 146. *Chloromyxum misgurni*. After Kudo (1916). $\times 1750$.
- Figs. 141 and 142. Trophozoites (1916, Figs. 3f and 3g).
- Figs. 143 to 145. Different views of fresh spore (1916, Figs. 3a, 3c and 3b).
- Fig. 146. A spore treated with potassium hydrate (1916, Fig. 3e).
- Figs. 147 to 152. *Chloromyxum fujitai*. After Kudo (1916). $\times 1750$.
- Fig. 147. A trophozoite (1916, Fig. 4a).
- Fig. 148. A fresh spore (1916, Fig. 4e).
- Fig. 149. A spore stained with Giemsa's mixture (1916, Fig. 4g).
- Figs. 150 to 152. Two surface views and an optical cross-section of a stained spore, showing the ridges on the shell valves (1916, Figs. 4b, 4d and 4c).
- Figs. 153 to 156. Spores of *Chloromyxum clupeiidae*.
- Figs. 153 to 155. Fresh spores. After Linton (1901, Fig. 3). "Variously magnified."
- Fig. 156. A spore. After Hahn (1917b, Fig. 10). $\times 1650$.
- Figs. 157 and 158. Two views of *Chloromyxum granulosum*. After Davis (1917, Figs. 137 and 138). $\times 1500$.




PLATE VIII

EXPLANATION OF PLATE

- Figs. 159 to 182. *Chloromyxum trijugum*. Original.
- Figs. 159 to 174. Trophozoites of various form and age.
- Figs. 159 and 160. Trophozoites of medium size. $\times 640$.
- Fig. 161. A trophozoite, ten minutes after it was removed from the host. $\times 1700$.
- Figs. 162 and 163. The movements of pseudopodia of the same specimen in ten minutes. $\times 1700$.
- Fig. 164. The same specimen after thirty minutes. $\times 1700$.
- Figs. 165 to 167. A trophozoite, showing the change of pseudopodia in five and ten minutes. $\times 1700$.
- Figs. 168 to 172. Small trophozoite with different numbers of the nuclei. Fig. 172 is probably a disporous form. $\times 2350$.
- Fig. 173. A monosporous trophozoite with a young spore. $\times 2350$.
- Fig. 174. A young spore. $\times 2350$.
- Figs. 175 to 177. Different views of fresh spores. $\times 1700$.
- Fig. 178. A Giemsa stained spore. $\times 1700$.
- Figs 179 and 180. Side views of the valves showing the ridges by Giemsa staining. $\times 1700$.
- Fig. 181. A spore treated with potassium hydrate solution, and stained with Giemsa solution. $\times 1700$.
- Fig. 182. A spore from which the sporoplasm is leaving the shell. From the Giemsa smear of the infected bile. $\times 1700$.




PLATE IX

EXPLANATION OF PLATE

Fig. 183 to 186. *Sphaerospora divergens*.

Fig. 183. Trophozoite. After Thélohan (1895, Fig. 12). $\times 750$.

Figs. 184 and 185. Two views of spore. After Auerbach (1912, Pl. 5, Fig. 4). \times about 1500.

Fig. 186. A spore treated with nitric acid. After Thélohan (1895, Fig. 13). $\times 1500$.

Figs. 187 and 88. Spores of *Sphaerospora elegans*. After Thélohan (1890b, Fig. 1). \times about 1000.

Fig. 189. A spore of *Sphaerospora rostrata*. After Thélohan (1895, Fig. 93). \times about 1635.

Figs. 190 to 192. Spores of *Sphaerospora masovica*. After Cohn (1902, Fig. 3). $\times 1000$.

Fig. 192. Spore with extruded polar filaments and "starren Fäden" by warming.

Figs. 193 and 194. Spores of *Sphaerospora platessae*. After Woodcock (1904, Fig. 7d). $\times 900$.

Figs. 195 to 197. Spores of *Sphaerospora angulata*. After Fujita (1912, Fig. 3). \times about 2800.

Figs. 198 and 199. Spores of *Sphaerospora polymorpha*. After Davis (1917, Figs. 91 and 92) $\times 1500$.

Figs. 200 to 204. *Sphaerospora carassii*. Original.

Fig. 200. A trophozoite. $\times 2250$.

Figs. 201 to 203. Different views of spores. $\times 1800$.

Fig. 204. A young spore. $\times 2250$.


Figs. 205 to 209. *Sinuolinea dimorpha*. After Davis (1916).

Fig. 205. A fresh trophozoite (1916, Fig. 1). $\times 1400$.

Figs. 206 and 207. Trophozoites with erythrocytes in different stages of disintegration (1916, Figs. 2 and 57). $\times 640$.

Fig. 208. A stained disporous trophozoite (1916, Fig. 41).

Fig. 209. A stained gemmule (1916, Fig. 72).



KUDO

STUDIES ON MYXOSPORIDIA

PLATE IX

PLATE X

EXPLANATION OF PLATE

Figs. 210 to 213. *Sinuolinea dimorpha*. After Davis (1916 and 1917).

Fig. 210. A living trophozoite (1916, Fig. 56). $\times 640$.

Fig. 211. A living trophozoite from which a gemmule is just escaping (1916, Fig. 60). $\times 640$.

Figs. 212 and 213. Spores (1917, Figs. 99 and 100). $\times 1400$.

Figs. 214 to 216. Spores of *Sinuolinea capsularis*. After Davis (1917, Figs. 105 to 107). $\times 1500$.

Figs. 217 and 218. Spores of *Sinuolinea arborescens*. After Davis (1917, Figs. 109 to 110). $\times 1500$.

Fig. 219. Spore of *Sinuolinea opacita*. After Davis (1917, Fig. 112). $\times 1500$.


Fig. 220. Spore of *Sinuolinea brachiophora*. After Davis (1917, Fig. 113). $\times 1500$.

Figs. 221 to 224. *Myxidium lieberkühni*. After Bütschli (1881 and 1882).

Fig. 221. A trophozoite (1882, Pl. 38, Fig. 12). \times about 60.

Fig. 222. A trophozoite (1882, Pl. 38, Fig. 13). $\times 160$.

Figs. 223 and 224. Trophozoites (1881, Figs. 27 and 26).



KUDO

STUDIES ON MYXOSPORIDIA

PLATE X

PLATE XI

EXPLANATION OF PLATE

Figs. 225 to 240. *Myxidium lieberkühni*.

Fig. 225. A trophozoite. After Lieberkühn from Gurley (1894, Pl. 43, Fig. 1a). $\times 330$.

Figs. 226 and 227. Trophozoites. After Bütschli (1881, Figs. 25 and 31).

Figs. 228 to 230. Stained trophozoites. After Thélohan (1895, Figs. 44, 46 and 45). $\times 750$.

Fig. 231. A trophozoite forming daughter individuals by budding. After Cohn (1895 Fig. 5).

Fig. 232. Four figures showing the separation of a bud. After Cohn (1895, Figs. 6a, 6b, 6d and 6e).

Fig. 233. A cross-section of a trophozoite, showing the ectoplasm, mesoplasm and endoplasm. After Cohn (1895, Fig. 2).

Fig. 234. A trophozoite. After Laveran and Mesnil (1902, Fig. 3). $\times 500$.

Fig. 235. A part of a trophozoite. After Laveran and Mesnil (1902, Fig. 3). $\times 800$.

Figs. 236 and 237. Young trophozoites undergoing division. After Laveran and Mesnil (1902, Figs. 4 and 5). $\times 1000$.

Fig. 238. An isolated spore. After Bütschli (1881, Fig. 32). \times about 2500.

Figs. 239 and 240. Fresh and stained spores. After Thélohan (1895, Figs. 47 and 48). $\times 1500$.

Figs. 241 to 251. *Myxidium incurvatum*.

Fig. 241. A monosporous trophozoite. After Parisi (1912, Fig. 1).

Fig. 242. A spore. After Thélohan (1895, Fig. 54). $\times 1500$.

Figs. 243 and 244. Different views of spore. After Parisi (1912, Fig. 1).

Fig. 245. A young spore. After Georgévitch (1916, Fig. 11).


Figs. 246 to 248. Spores. After Georgévitch (1916, Figs. 10, 9 and 8).

Figs. 249 to 251. Spores. After Davis (1917, Figs. 119 to 121). $\times 1500$.

Fig. 252. Trophozoite of *Myxidium sphaericum*. After Thélohan (1895, Fig. 28). $\times 1500$.

Fig. 253. A spore of *Myxidium histophilum*. After Thélohan (1895, Fig. 49). $\times 1500$.

Fig. 254. A spore of *Myxidium* sp. After Leydig from Gurley (1894, Pl. 47, Fig. 6).



KUDO

STUDIES ON MYXOSPORIDIA

PLATE XI

PLATE XII

EXPLANATION OF PLATE

- Figs. 255 to 257. *Myxidium danilewskyi*. After Laveran (1898).
 Figs. 255 and 256. Longitudinal and transverse sections thru renal tubules, showing the trophozoites (1898, Figs. 1 and 2). Fig. 255. $\times 350$.
 Fig. 257. Spores (1898, Figs. 4, 5 and 6). $\times 800$.
 Fig. 258. Spores of *Myxidium giganteum*. After Doflein (1898, Fig. 48). \times about 500.
 Figs. 259 to 261. Spores of *Myxidium giardi*. After Cépède (1906a, Figs. 1, 3 and 2). $\times 2000$.
 Figs. 262 to 265. Spores of *Myxidium pfeifferi*. After Auerbach (1908, Figs. 6 and 7). \times about 2000, except Fig. 265.
 Fig. 266. A spore of *Myxidium inflatum*. After Auerbach (1909, Fig. 3a). \times about 1500.
 Fig. 267. A spore of *Myxidium bergense*. After Auerbach (1910a, Fig. 57). \times about 1820.
 Fig. 268. A spore of *Myxidium procerum*. After Auerbach (1910a, Fig. 58). \times about 2000.
 Figs. 269 to 271. Spores of *Myxidium mackiei*. After Bosanquet (1910, Fig. 12). \times about 1250.
 Figs. 272 and 273. Spores of *Myxidium macrocapsulare*. After Auerbach (1910d, Figs. 1a and 1b). $\times 3000$.
 Figs. 274 to 276. *Myxidium* sp. After Awerinzew (1908 and 1911).
 Fig. 274. A monosporous trophozoite (1911, Fig. C).
 Fig. 275. A disporous trophozoite (1908, Pl. 2, Fig. 6). Obj. E and oc. 4.
 Fig. 276. A spore (1908, Pl. 1a, Fig. 17). \times about 1000.
 Figs. 277 and 278. Spores of *Myxidium depressum*. After Parisi (1912, Figs. 2a and 2b). \times about 1600.
 Figs. 279 and 280. Spores of *Myxidium oviforme*. After Parisi (1912, Fig. 3). \times about 1600.
 Figs. 281 to 284. Spores of *Myxidium anguillae*. After Ishii (1915, Fig. 3a). $\times 1450$.
 Figs. 285 and 286. *Myxidium* sp. After Mavor (1915).
 Fig. 285. A spore treated with ammonia water (1915, Fig. 3a). $\times 660$.
 Fig. 286. A spore (1915, Fig. 3b). $\times 1320$.
 Figs. 287 to 290. Spores of *Myxidium gadi*. After Georgévitch (1916, Figs. 1, 4, 3 and 2).
 Fig. 228. A spore from *Solea vulgaris*.
 Figs. 289 and 290. Young spores.




PLATE XIII

EXPLANATION OF PLATE

- Fig. 291. A spore of *Myxidium glutinosum*. After Davis (1917, Fig. 124). $\times 1400$.
- Figs. 292 and 293. Spores of *Myxidium phyllium*. After Davis (1917, Figs. 126 and 127). $\times 2000$.
- Figs. 294 and 295. Spores of *Myxidium kagayamai*. After Kudo (1916, Fig. 2). $\times 1750$ and $\times 1000$ respectively.
- Figs. 296 to 307. *Sphaeromyxa balbianii*.
- Fig. 296. A trophozoite. After Thélohan (1895, Fig. 57). $\times 3$.
- Fig. 297. A trophozoite. After Davis (1917, Fig. 128). $\times 640$.
- Fig. 298. A trophozoite in plasmotomy. After Georgévitch (1916, Fig. 15).
- Figs. 299 and 300. Spores. After Thélohan (1895, Figs. 58 and 59). $\times 1500$.
- Fig. 301. An end of a spore. After Thélohan (1895, Fig. 60).
- Fig. 302. A spore treated with nitric acid. After Thélohan (1895, Fig. 61).
- Fig. 303. A spore. After Parisi (1912, Fig. 4). \times about 1750.
- Fig. 304. A spore. After Davis (1917, Fig. 130). $\times 2100$.
- Figs. 305 to 307. Mature and young spores (Figs. 306 and 307). After Georgévitch (1916, Figs. 17, 20 and 19).
- Figs. 308 to 311. *Sphaeromyxa immersa*. After Lutz (1889: 301).
- Fig. 308. An infected gall-bladder of *Bufo aqua* (1889, Fig. 1). $\times 1$.
- Fig. 309. Spores (1889, Figs. 4, 5 and 6).
- Fig. 310. A spore (1889, Fig. 10). $\times 600$.
- Fig. 311. A spore with extruded polar filaments (1889, Fig. 7).
- Figs. 312 to 314. Spores of *Sphaeromyxa incurvata*. After Doflein (1898, Fig. 49). \times about 1000.
- Figs. 315 and 316. *Sphaeromyxa sabrazesi*. After Schröder (1907).
- Fig. 315. A trophozoite (1907, Fig. 1). $\times 15$.
- Fig. 316. A cross section thru a trophozoite (1907, Fig. 3). $\times 1500$.





PLATE XIV

EXPLANATION OF PLATE

- Figs. 317 to 322. Spores of *Sphaeromyxa sabrazesi*.
 Figs. 317 to 319. After Laveran and Mesnil (1900, Figs. 1, 3 and 4). $\times 1500$.
 Fig. 318. A spore treated with nitric acid (1900, Fig. 3).
 Figs. 320 and 321. Spores. After Schröder (1907, Figs. 39 and 41). $\times 1500$.
 Fig. 322. A polar capsule. After Schröder (1907, Fig. 45).
 Figs. 323 and 324. Spores of *Sphaeromyxa hellandi*. After Auerbach (1909, Fig. 5). \times
 about 1500.
 Figs. 325 and 326. Spores of *Sphaeromyxa exneri*. After Awerinzew (1913a, Fig. 3). \times
 about 365.
 Fig. 327. A spore of *Sphaeromyxa gasterostei*. After Georgévitch (1916, Fig. 22).
 Figs. 328 to 331. *Zschokkella hildae*. After Auerbach (1910a and 1912).
 Fig. 328. A monosporous trophozoite (1912, Pl. 5, Fig. 2).
 Figs. 329 to 331. Spores (1910a, Fig. 62).
 Figs. 332 and 333. Spores of *Zschokkella nova*. After Klokacewa (1914, Fig. 2). \times about
 2500.
 Figs. 334 to 338. Spores of *Zschokkella acheilognathi*. After Kudo (1916).
 Figs. 334 to 336. Different views of fresh spores (1916, Figs. 3d, 3e and 3f). $\times 2250$.
 Fig. 337. A young spore. Original. $\times 2785$.
 Fig. 338. A stained spore (1916, Fig. 3h). $\times 2800$.
 Figs. 339 and 340. Spores of *Zschokkella globulosa*. After Davis (1917, Figs. 135 and 134).
 $\times 1500$.
 Figs. 341 to 343. Spores of *Myxosoma dujardini*. After Thélohan (1895, Figs. 90, 91, and
 89). $\times 1500$.
 Figs. 344 to 347. Spores of *Myxosoma funduli*. After Kudo (1918a, Fig. A). $\times 1500$.



KUDO

STUDIES ON MYXOSPORIDIA

PLATE XIV

PLATE XV

EXPLANATION OF PLATE

- Fig. 348. A spore of *Myxosoma(?) lobatum*. After Nemeček (1911, Fig. 18). $\times 1000$.
Fig. 349 to 354. *Lentospora cerebralis*. After Plehn (1905). $\times 1200$.
Fig. 349. Various young ameboid forms; stained (1905, Textfig. 5).
Fig. 350. A stained larger form (1905, Textfig. 5).
Fig. 351. A trophozoite with two spores (1905, Textfig. 4).
Fig. 352. Various spores (1905, Textfig. 2).
Fig. 353. A stained spore (1905, Textfig. 3).
Fig. 354. A spore with extruded polar filaments (1905, Textfig. 2).
Fig. 355. A spore of *Lentospora multiplicata*. After Reuss (1906, Fig. 8). $\times 1500$.
Figs. 356 to 359. Spores of *Lentospora asymmetrica*. After Parisi (1912, Fig. 7). \times about 1500.
Figs. 360 to 362. Spores of *Lentospora acuta*. After Fujita (1912, Fig. 2). $\times 2000$.
Figs. 363 and 364. Spores of *Myxobolus piriformis*. After Thélohan (1895, Figs. 117 and 116). $\times 1500$.
Figs. 365 and 366. Spores of *Myxobolus unicasulatus*. After Müller (1841, Fig. 5).
Fig. 367. A spore of *Myxobolus fuhrmanni*. After Auerbach (1909, Fig. 1b). $\times 1840$.
Fig. 368. A spore of *Myxobolus oculi-leucisci*. After Trojan (1909, Fig. 3). $\times 2000$.
Figs. 369 and 370. Spores of *Myxobolus toyamai*. After Kudo (Original and 1917, Fig. 40). $\times 2500$.
Figs. 371 and 372. Spores of *Myxobolus notatus*. After Mavor (1916, Figs. 6C and 6B). $\times 2600$.
Figs. 373 and 374. Spores of *Myxobolus rohita*. After Southwell and Prashad (1918, Figs. 26 and 27). \times about 1720 and 700 respectively.
Figs. 375 and 376. Spores of *Myxobolus seni*. After Southwell and Prashad (1918, Figs. 29 and 30). \times about 1700.
Figs. 377 and 378. Spores of *Myxobolus misgurni*. Original. $\times 1500$.





PLATE XVI

EXPLANATION OF PLATES

- Figs. 379 to 385. *Myxobolus pfeifferi*.
 Figs. 379 and 380. Parts of section thru cyst. After Keysselitz (1908a, Pl. 15, Figs. 1 and 2).
 Fig. 381. A spore. After Thélohan (1895, Fig. 77). $\times 1500$.
 Fig. 382. An optical section of spore. After Keysselitz (1908a, Fig. A).
 Fig. 383. A spore treated with Lugol's solution. After Keysselitz (1908a, Pl. 14: Fig. 92).
 Fig. 384. A spore with extruded filaments. After Keysselitz (1908a, Fig. C).
 Fig. 385. A stained young spore. After Keysselitz (1908a, Pl. 14, Fig. 81).
 Fig. 386. A spore of *Myxobolus dispar*. After Thélohan (1895, Fig. 86). \times about 1500.
 Figs. 387 to 389. Spores of *Myxobolus ellipsoides*. Fig. 389. An abnormal spore with six polar capsules. After Thélohan (1895, Figs. 112, 113 and 115).
 Fig. 390. A part of the infected intestine of *Mugil auratus*, showing cysts of *Myxobolus exiguus*. After Parisi (1912, Fig. 9e). \times about 3.
 Fig. 391. A spore of *Myxobolus exiguus*. After Thélohan (1895, Fig. 98). $\times 1500$.
 Figs. 392 to 395. Spores of *Myxobolus exiguus*. After Parisi (1912, Fig. 9). $\times 2500$.
 Fig. 396. A spore of *Myxobolus oviformis*. After Thélohan (1895, Fig. 81). $\times 1500$.
 Figs. 397 to 398. Spores of *Myxobolus mülleri*. After Thélohan (1895, Figs. 96 and 97). $\times 1500$.
 Figs. 399 and 400. Spores of *Myxobolus mülleri*. After Bütschli (1881, Figs. 1 and 2).
 Fig. 401. A spore of *Myxobolus mülleri* in conc. sulphuric acid. After Bütschli (1881, Fig. 6).
 Figs. 402 and 403. Abnormal spores of *Myxobolus mülleri*. After Bütschli (1881, Figs. 9 and 8).
 Figs. 404 and 405. Spores of *Myxobolus lintoni*. After Linton (1891, Figs. 3 and 5).
 Fig. 406. Diagram of the cross section of a spore of *Myxobolus lintoni*. After Linton (1891, Fig. 8).
 Fig. 407. A spore of *Myxobolus lintoni* with extruded polar filaments. After Linton (1891, Fig. 10).
 Fig. 408. A stained spore of *Myxobolus lintoni*. After Gurley (1894, Pl. 26, Fig. 7). \times about 2000.
 Figs. 409 and 410. Spores of *Myxobolus globosus*. After Gurley (1894, Pl. 26, Fig. 7). \times about 2900.
 Fig. 411. Spores of *Myxobolus inaequalis*. After Müller (1841, Fig. 6).



KUDO


STUDIES ON MYXOSPORIDIA

PLATE XVI

PLATE XVII

EXPLANATION OF PLATE

- Figs. 412 to 416. Spores of *Myxobolus oblongus*. Figs. 412 to 414. After Gurley (1894, Pl. 26, Fig. 6). $\times 2300$. Figs. 415 and 416. After Müller (1841, Fig. 9).
- Figs. 417 and 418. Spores of *Myxobolus transovalis*. After Gurley (1894, Pl. 29, Fig. 1).
- Figs. 419 and 420. Spores of *Myxobolus obesus*. After Balbiani (1884, Fig. 39).
- Fig. 421. Spores of *Myxobolus cycloides*. After Müller (1841, Fig. 3).
- Figs. 422 and 423. Spores of *Myxobolus anurus*. After Cohn (1895, Fig. 25). $\times 1500$.
- Fig. 424. Spores of *Myxobolus* sp. $\times 700$. After Bütschli (1882, Pl. 36, Fig. 23).
- Fig. 425. *Myxobolus* sp. After Gurley (1894, Pl. 28: Fig. 4a). \times about 1500.
- Figs. 426 to 429. Spores of *Myxobolus* sp. After Müller (1841, Fig. 4).
- Fig. 430. A vegetative form of *Myxobolus cyprini*. After Doflein (1898, Fig. 112).
- Figs. 431 and 432. Spores of *Myxobolus cyprini*. After Doflein (1898, Figs. 113 to 115).
- Figs. 433 to 436. *Myxobolus neurobius*. After Schuberg and Schröder (1905).
- Figs. 433 and 434. Longitudinal and transverse sections thru infected nerve fibres (1905, Figs. 2 and 4a). $\times 520$.
- Figs. 435 and 436. Spores (1905, Figs. 5 and 6). Comp. oc. 12 and imm. obj. 2mm.
- Figs. 437 to 441. *Myxobolus aeglefini*. After Auerbach (1906a).
- Fig. 437. A cyst in the sclerotic cartilage of the eye of *Gadus aeglefini* (1906a, Fig. 2).
- Figs. 438 and 439. Spores (1906a, Figs. 5a and 3d). \times about 1320.
- Figs. 440 and 441. Abnormal spores (1906a, Figs. 5b and 5c). \times about 1320.
- Figs. 442 to 445. *Myxobolus gigas*. After Auerbach (1906b).
- Fig. 442. A part of the section of a cyst (1906b, Fig. 1).
- Figs. 443 to 445. Spores (1906b, Figs. 3a, 3c, 5 and 3b). \times about 850.
- Figs. 446 and 447. Spores of *Myxobolus volgensis*. After Reuss (1906, Fig. 1). $\times 2000$.



KUDO


STUDIES ON MYXOSPORIDIA

PLATE XVII


PLATE XVIII

EXPLANATION OF PLATE


- Fig. 448. The branchia of *Lucioperca volgensis* with cysts of *Myxobolus volgensis*. After Reuss (1906, Fig. 2). $\times 2.25$.
- Fig. 449. A spore of *Myxobolus scardinii*. After Reuss (1906, Fig. 3). $\times 1500$.
- Fig. 450. Air bladder of *Scardinius erythrophthalmus* with the cysts of *Myxobolus physophilus*. After Reuss (1906, Fig. 5). $\times 2$.
- Fig. 451. A spore of *Myxobolus physophilus*. After Reuss (1906, Fig. 4). $\times 1500$.
- Fig. 452. A spore of *Myxobolus macrocapsularis*. After Reuss (1906, Fig. 6). $\times 1500$.
- Fig. 453. A spore of *Myxobolus sandrae*. After Reuss (1906, Fig. 7). $\times 2000$.
- Fig. 454. A spore of *Myxobolus bramae*. After Reuss (1906, Fig. 9). $\times 1500$.
- Fig. 455. A spore of *Myxobolus balleri*. After Reuss (1906, Fig. 10). $\times 1500$.
- Fig. 456. A spore of *Myxobolus cyprinicola*. After Reuss (1906, Fig. 11). $\times 1500$.
- Figs. 457-459. *Myxobolus squamae*. After Keysselitz (1908a).
- Fig. 457. A part of the infected scale (1908a, Fig. G.)
- Fig. 458. A spore treated with Lugol's solution (1908a, Pl. 14, Fig. 94).
- Fig. 459. A stained spore (1908a, Pl. 14, Fig. 96).
- Figs. 460 and 461. Spores of *Myxobolus cordis*. After Keysselitz (1908a).
- Fig. 460. A spore treated with Lugol's solution (1908a, Pl. 16, Fig. 16).
- Fig. 461. A strained spore (1908a, Fig. B on page 281).
- Figs. 462 to 464. Spores of *Myxobolus musculi*. After Keysselitz (1908a).
- Fig. 462. A spore treated with Lugol's solution (1908a, Pl. 15, Fig. 13).
- Fig. 463 and 464. Stained spores (1908a, Figs. D and E on page 286).
- Fig. 465. Spores of *Myxobolus* sp. After Wegener (1910, Fig. 44). $\times 1050$.
- Fig. 466. A spore of *Myxobolus permagnus*. After Wegener (1910, Fig. 45). $\times 1050$.
- Fig. 467. Spores of *Myxobolus rotundus*. After Nemeček (1911, Figs. 10 and 11). $\times 1000$.
- Fig. 468. Spores of *Myxobolus minutus*. After Nemeček (1911, Figs. 16 and 17). $\times 1000$.
- Figs. 469 and 470. Spores of *Myxobolus magnus*. After Awerinzew (1913, 76). \times about 340.
- Figs. 471 to 473. Spores of *Myxobolus carassii*. After Klokacewa (1914, Fig. 1). \times about 2400.




448




449




451




455




456




452




454




453




460




458



459




450




457



473



461



462



469



470



463



464



466




465



467



468



471



472

PLATE XIX

EXPLANATION OF PLATE

- Figs. 474 to 476. *Myxobolus funduli*. After Hahn (1915 and 1917).
 Fig. 474. A cyst from the gill filament (1917, Fig. 1).
 Fig. 475. A stained spore (1915, Fig. 28). $\times 2000$.
 Fig. 476. Diagram of the cross section of a spore (1915, Fig. 30).
 Fig. 477. A spore of *Myxobolus pleuronectidae*. After Hahn (1917a, Fig. 2). $\times 1575$.
 Fig. 478. A spore of *Myxobolus capsulatus*. After Davis (1917, Fig. 139). $\times 1500$.
 Figs. 479 and 480. Spores of *Myxobolus nodularis*. After Southwell and Prashad (1918, Pl. 11, Figs. 34 and 35). \times about 1450.
 Fig. 481. A spore of *Myxobolus miyairii*. After Miyairi (1909, Fig. 14).
 Figs. 482 to 485. Spores of *Myxobolus koi*. Original. $\times 1300$.
 Figs. 482 to 484. Different views.
 Fig. 485. A spore stained with Giemsa's mixture.
 Figs. 486 and 487. *Henneguya psorospermica*. After Thélohan (1895).
 Fig. 486. A cross section thru branchial lamella of *Esox lucius* with a cyst (1895, Fig. 82).
 Fig. 487. Two spores (1895, Figs. 83 and 84). \times about 1000.
 Figs. 488 and 489. *Henneguya minuta*. After Cohn (1895).
 Fig. 488. A longitudinal section of an infected branchial lamella (1895, Fig. 29).
 Fig. 489. Two spores. One with two vacuoles(?) (1895, Fig. 30). \times about 450.
 Fig. 490 and 491. Spores of *Henneguya oviperda*. After Cohn (1895, Fig. 31).
 Fig. 491. A spore with extruded "starren Fäden" and polar filaments.
 Figs. 492 and 493. *Henneguya lobosa*. After Cohn.
 Fig. 492. An external view of the parasite on the gill (1895, Fig. 18).
 Fig. 493. Two spores and one unseparated young spores (1895, Fig. 21).
 Figs. 494 and 495. *Henneguya media*. After Thélohan (1890b).
 Fig. 494. A sporoblast in the ovary of *Gasterosteus*, with one spore (1890b, Fig. 18).
 Fig. 495. Spores (1890b, Fig. 1).
 Fig. 496. The peripheral portion of a section of a cyst of *Henneguya psorospermica*, showing the characteristic structure. After Thélohan (1895: 237).
 Figs. 497 to 499. Spores of *Henneguya schizura*. After Müller (1841, Fig. 1).
 Figs. 500 to 503. Spores of *Henneguya creplini*. After Creplin (1842, Figs. B, E, A and C).
 Fig. 504. Spores of *Henneguya linearis*. After Müller (1841, Fig. 10).




PLATE XX

EXPLANATION OF PLATE

- Fig. 505. Spores of *Henneguya Gurleyi*. After Gurley (1894, Pl. 33, Figs. 8c, 6 and 7).
× about 3100.
- Fig. 506. Spores of *Henneguya strongylura*. After Müller (1841, Fig. 2).
- Fig. 507. Spores of *Henneguya monura*. After Ryder (1880, Figs. 1c and 2d).
- Fig. 508. Spores of *Henneguya kolesnikovii*. After Kolesnikov from Gurley (1894, Pl. 35, Fig. 7).
- Figs. 509 to 512. *Henneguya macrura*. After Gurley (1894). × about 2100.
- Figs. 509 and 510. Spores (1894, Pl. 32, Fig. 5, Pl. 33, Fig. 1).
- Fig. 511. A spore treated with iodine, showing the "beading of the tail" (1894, Pl. 33, Fig. 3).
- Fig. 512. A tail separated from the main part by iodine (1894, Pl. 33, Fig. 4).
- Fig. 513. Spores of *Henneguya zschokkei*. After Zschokke (1898, Figs. 2 and 1).
- Fig. 514. Spores of *Henneguya* sp. After Benecke from Gurley (1894, Pl. 29, Fig. 8).
- Fig. 515. Spore of *Henneguya tenuis*. After Vaney and Conte (1901, Fig. 2).
- Figs. 516 and 517. Spores of *Henneguya nüsslini*. After Schuberg and Schröder (1905, Figs. 13 and 14). Comp. oc. 12 and obj. 2mm.
- Figs. 518 to 523. *Henneguya legeri*. After Cépède (1913).
- Figs. 518 to 521. Trophozoites (1913, Figs. 2, 23, 15 and 25). ×900.
- Fig. 519. A trophozoite in division (1913, Fig. 23). ×900.
- Fig. 521. A trophozoite stained with iron hematoxylin (1913, Fig. 25). ×900.
- Fig. 522. An elongated spore (1913, Fig. 26). ×900.
- Fig. 523. An ovoidal spore (1913, Fig. 24). ×450.
- Fig. 524. A spore of *Henneguya miyairii*. After Miyairi (1909, Fig. 11).





PLATE XXI

EXPLANATION OF PLATE

- Figs. 525 and 526. Spores of *Henneguya acerinae*. After Schröder (1906, Figs. 5 and 6).
×1650.
- Figs. 527 to 535. *Henneguya gigantea*. After Nemeček (1911). ×1000.
- Fig. 527. A mature spore with extruded polar filaments (1911, Fig. 1).
- Figs. 528 to 535. Stages in development of spores (1911, Figs. 2 to 9).
- Figs. 536 to 539. Spores of *Henneguya(?)* sp. After Nemeček (1911, Figs. 12 to 15).
×1000.
- Figs. 540 to 543. *Henneguya gasterostei*. After Parisi (1912). × about 1500.
- Fig. 540. A disporous trophozoite (1912, Fig. 10d).
- Figs. 541 and 542. Two spores (1912, Figs. 10f and 10e).
- Fig. 543. A young spore (1912, Fig. 10c).
- Figs. 544 and 545. Spores of *Henneguya neapolitana*. After Parisi (1912, Fig. 11). ×
about 1500.
- Figs. 546 to 549. Various trophozoites of *Henneguya mictospora*. Original.
- Fig. 546. A monosporous trophozoite with a young spore. ×950.
- Figs. 547 and 549. Trophozoites in vivo. ×650.
- Fig. 548. A stained binucleated young trophozoite. ×1700.



KUDO


STUDIES ON MYXOSPORIDIA

PLATE XXI

PLATE XXII

EXPLANATION OF PLATE

- Figs. 550 to 557. *Henneguya mictospora*. Original.
- Fig. 550. A stained trophozoite. $\times 1700$.
- Figs. 551 to 553. Three different stages of development of disporous trophozoites. Giemsa. $\times 1700$.
- Figs. 554 and 555. Two stages of monosporous trophozoites. Giemsa. $\times 1700$.
- Figs. 556 and 557. Different views of mature spores in vivo. $\times 2000$.
- Figs. 558 and 559. *Henneguya wisconsinensis*. After Mavor and Strasser (1916).
- Fig. 558. A trophozoite in vivo (1916, Fig. 1a). $\times 570$.
- Fig. 559. A fresh spore (1916, Fig. 3d). $\times 4000$.
- Figs. 560 and 561. Trophozoites of *Chloromyxum catostomi*. Original. $\times 1500$.
- Figs. 562 to 565. *Chloromyxum clupeiidae*. Original drawn from Dr. Tyzzer's smears which were restained. $\times 2360$.
- Fig. 562. Anterior end view of two spores in preserved and decolorized smears.
- Fig. 563. The same views of three spores restained with Giemsa mixture.
- Fig. 564. Front view of a preserved and decolorized spore.
- Fig. 565. The same views of two spores restained with Giemsa's mixture.
- Figs. 566 to 572. Spores of *Myxobolus orbiculatus*. Original.
- Figs. 566, 569 and 570. Different views of normal spores in preserved specimen. $\times 1500$.
- Figs. 567 and 568. Abnormal spores. $\times 1500$.
- Fig. 571. A spore stained with Lugol's solution. $\times 1500$.
- Fig. 572. A spore stained with Giemsa's mixture. $\times 2360$.



KUDO

STUDIES ON MYXOSPORIDIA

PLATE XXII


PLATE XXIII

EXPLANATION OF PLATE

- Figs. 573 to 576. *Myxobolus orbiculatus*. Original.
- Fig. 573. Young vegetative forms in the muscle fibres. From a section stained with Heidenhain's iron hematoxylin. $\times 900$.
- Figs. 574 and 575. Two vegetative forms under higher magnifications. $\times 1500$.
- Fig. 576. A cross section thru the infected muscle fibres, stained with Heidenhain's iron hematoxylin. $\times 900$.
- Figs. 577 to 581. *Hoferellus cyprini*. After Doflein (1898).
- Figs. 577 and 578. Two vegetative forms (1898, Figs. 106 and 105).
- Figs. 579 to 581. Spores (1898, Figs. 108 and 107).
- Figs. 582 and 583. Genus incert. *merlucii*. After Perugia from Gurley (1894, Pl. 29, Figs. 2 and 7).
- Figs. 584 and 585. Genus incert. *congru*. After Perugia from Gurley (1894, Pl. 6, Figs. 7 and 3).
- Figs. 586 and 587. Genus et species incertae. After Mavor (1915).
- Fig. 586. A trophozoite with attached *Ceratomyxa acadensis* (1915, Fig. 8). $\times 830$.
- Fig. 587. A trophozoite (1915, Fig. 6).
- Figs. 588 and 589. Trophozoites of genus et species incertae. After Mavor (1916a, Figs. 3d and 3b). $\times 660$.
- Fig. 586. A trophozoite with attached *Ceratomyxa acadensis* (1915, Fig. 8). $\times 830$.
- Fig. 587. A trophozoite (1915, Fig. 6).
- Figs. 588 and 589. Trophozoites of genus et species incertae. After Mavor (1916a, Figs. 3d and 3b). $\times 660$.
- Fig. 590. Spores of genus et species incertae. After Linton (1891a, Fig. 2).
- Figs. 591 to 593. *Myxobolus hylae*. After Johnston and Bancroft (1918).
- Fig. 591. A transverse section of a heavily infected testis of *Hyla aurea*, (1918, Fig. 1). \times about 11.
- Fig. 592. Different views of normal spores, stained (1918, Fig. 3). \times about 800.
- Fig. 593. Abnormal spores (1918, Fig. 4). \times about 800.
- Figs. 594 to 596. *Lentospora dermatobia*. After Ishii (1915).
- Fig. 594. A part of the infected skin of the host (1915, Fig. 2). $\times 140$.
- Figs. 595 and 596. Different views of spore (1915, Figs. 4 and 3). $\times 1450$.
- Figs. 597 to 601. *Myxobolus discrepans*. Original.
- Fig. 597. An infected branchial lamella showing the cysts of various size and form. \times about 4.
- Figs. 598 to 601. Unstained preserved spores, showing different views. \times about 1500.




573




574




575




577




578



576



579




580




581




582



583




584




585




586



587




588




589




590




591




592




593




594




595




596




597




598



599



600




601

PLATE XXIV

EXPLANATION OF PLATE

- Figs. 602 to 621. *Mitraspora elongata*. Original. $\times 1500$, except Figs. 620 and 621, $\times 2500$. Giemsa staining, unless otherwise stated.
- Fig. 602. A trophozoite showing various nuclei and sporoblasts at different stages of development.
- Fig. 603. A trophozoite with a mature and a young spore. Iron hematoxylin.
- Fig. 604. A trophozoite with two mature spores.
- Figs. 605 to 609. Formation and development of sporoblasts.
- Figs. 610 to 613. Developing spores. Fig. 612; Delafield's hematoxylin.
- Fig. 614. A surface view of a preserved spore.
- Fig. 615. An optical section of a preserved spore.
- Fig. 616. Front view of a stained spore.
- Fig. 617. Lateral view of a spore stained with Heidenhain's iron hematoxylin.
- Fig. 618. A slightly elongated spore.
- Fig. 619. An abnormal spore.
- Fig. 620. A longitudinal section thru a polar capsule.
- Fig. 621. An oblique view of a polar capsule, showing the spirally coiled polar filament.
- Figs. 622 to 627. *Myxidium americanum*. Original. $\times 1500$.
- Figs. 622 and 623. Two young trophozoites. Giemsa.
- Fig. 624. A sporulating trophozoite in unstained preserved state.
- Fig. 625. A spore in preserved state.
- Fig. 626. A fresh spore treated with potassium hydrate solution.
- Fig. 627. A spore stained with Giemsa's mixture.
- Figs. 628 to 631. *Myxobolus mesentericus*. Original. $\times 1500$.
- Figs. 628 and 629. Front and lateral views of unstained preserved spores.
- Fig. 630. A Giemsa stained spore.
- Fig. 631. A spore with extruded polar filament (potassium hydrate), stained with Giemsa's mixture.



KUDO


STUDIES ON MYXOSPORIDIA

PLATE XXIV

PLATE XXV

EXPLANATION OF PLATE

- Figs. 632 to 642. *Chloromyxum wardi*. Original.
- Fig. 632. A young trophozoite. Smear and Giemsa. $\times 1500$.
- Fig. 633. A sporulating trophozoite. Giemsa. $\times 1500$.
- Figs. 634 to 637. Surface views and optical sections of four unstained spores, showing the sutural ridge and the fine striations on the shell. $\times 1500$.
- Fig. 638. A polar view of unstained spore. $\times 2360$.
- Figs. 640. Surface view and optical section of a single spore. $\times 2360$.
- Fig. 641. A front view of an unstained spore. $\times 2360$.
- Fig. 642. A Giemsa stained spore. $\times 1500$.
- Figs. 643 to 649. *Myxobolus aureatus*. After Ward (1919).
- Fig. 643. A fresh spore (1919, Fig. Aa).
- Figs. 644 and 645. Fresh spores kept for 24 hours or more in water (1919, Fig. Aa).
- Fig. 646. A preserved unstained spore (1919, Fig. Ba).
- Fig. 647. A spore stained with Delafield's hematoxylin (1919, Fig. Bd). $\times 1500$.
- Fig. 648. A spore with an extruded polar filament from section preparation stained with Giemsa's mixture (1919, Fig. Bg). $\times 1500$.
- Fig. 649. A young spore, stained with Giemsa's mixture (1919, Fig. Bi). $\times 1500$.
- Figs. 650 to 653. *Henneguya brachyura*. After Ward (1919). $\times 1500$.
- Fig. 650. A young spore stained with Giemsa's mixture (1919, Fig. Ce).
- Figs. 651 and 652. Front and lateral views of spore (1919: Figs. Cb and Cc).
- Fig. 653. A detached tail (1919, Fig. Cf).
- Figs. 654 to 656. *Henneguya salminicola*. After Ward (1919). $\times 1500$.
- Fig. 654. A young spore stained with Giemsa's mixture (1919, Fig. Fc).
- Figs. 655 and 656. Front and lateral views of unstained preserved spores.



INDEX

- African Myxosporidia, 14
 Air bladder, Myxosporidia parasitic in, 39
 Alaskan Myxosporidia, 16
 Alimentary canal, Myxosporidia parasitic in,
 39, 41
 Amphibian hosts of Myxosporidia, 36
 Annelida as host of Myxosporidian, 25
 Anterior end of spore, 47, 50
 process of spore, 47, 50, 51
 Asiatic Myxosporidia, 13
 Australian Myxosporidian, 14
 Austrian Myxosporidia, 24
 Blood vessel, Myxosporidia parasitic in, 39
 Body cavity, Myxosporidia parasitic in, 37,
 43, 45
 Bone, Myxosporidia parasitic in, 39
 Branchia, Myxosporidia parasitic in, 38
 Breadth of spore, 50
 Burma, Myxosporidia of, 14
 Canadian Myxosporidia, 16
 Capsulogenous cell of spore, 50
 Cartilage, Myxosporidia parasitic in, 39
 Ceratomyxa, 9, 56, 65, 178, 180
Ceratomyxa abbreviata, 76
 acadiensis, 71
 agglomerata, 77
 aggregata, 79
 amorpha, 78
 appendiculata, 67
 arcuata, 65
 attenuata, 75
 coris, 72
 drepanopsettae, 70
 flagellifera, 77
 globurifera, 67
 herouardi, 72
 inaequalis, 68
 linospora, 69
 lunata, 76
 mesospora, 73
 monospora, 78
 navicularia, 80
 pallida, 67
 ramosa, 69
 recurvata, 75
 reticularis, 68
 spati, 70
 sp. (?) Awerinzew, 71
 sp. Georgévitch, 72
 sphaerulosa, 66
 sphaerophora, 73
 spinosa, 80
 streptospora, 79
 taenia, 74
 truncata, 67
 tylosuri, 70
 undulata, 79
 Ceratomyxidae, 9, 56, 60, 178
 Chloromyxidae, 10, 57, 87, 178
 Chloromyxum, 10, 57, 87, 178, 183
Chloromyxum catostomi, 98
 caudatum, 88
 clupeiadae, 94
 cristatum, 91
 diploxys, 90
 dubium, 91
 fluviatile, 89
 fujitai, 93
 funduli, 93
 granulosum, 96
 koi, 92
 leydigi, 87
 magnum, 92
 misgurni, 93
 mucronatum, 89
 protei, 90
 quadratum, 88
 sp. Awerinzew, 91
 thymalli, 92
 trijugum, 96
 truttae, 90
 wardi, 99
 Classification of Myxosporidia after
 Auerbach, 52
 Davis, 54
 Doflein, 52
 Kudo, 56
 Parisi, 54
 Poche, 54
 Thélohan, 52
 Connective tissue, Myxosporidia parasitic in,
 37
 Cyst, 50
 Disporea, 52, 54
 Disporous, 50

- England, Myxosporidia of, 22
 European, Myxosporidia, 17
 Eurysporea, 9, 56, 60
 Eye, Myxosporidia parasitic in, 38
 Fish as hosts of Myxosporidia, 25
 fresh water, 44
 marine, 44
 Foramen of polar capsule, 47, 48, 50
 France, Myxosporidia of, 18
 Front view of spore, 47, 50
 Gall bladder, Myxosporidia parasitic in, 39
 Gemmule, 50
 German Myxosporidia, 20
 Heart, Myxosporidia parasitic in, 39
 Henneguya, 11, 59, 158, 179, 193
 Henneguya acerinae, 167
 brachyura, 171
 brevis, 162
 creplini, 162
 gasterostei, 169
 gigantea, 168
 gurleyi, 163
 kolesnikovi, 164
 legeri, 166
 linearis, 163
 lobosa, 161
 macrura, 164
 media, 161
 mictospora, 173
 minuta, 160
 miyairii, 172
 monura, 164
 neapolitana, 170
 nüsslini, 166
 oviperda, 160
 peri-intestinalis, 161
 psorospermica, 158
 salmonicola, 171
 schizura, 162
 sp. (Gurley) Labbé, 165
 sp. (Gurley) Labbé, 165
 sp. (?) Nemeček, 169
 strongylura, 163
 tenuis, 166
 texta, 159
 wisconsinensis, 170
 zschokkei, 165
 Hoferellus, 12, 59, 173, 179
 Hoferellus cyprini, 174
 Indian Myxosporidia, 14
 Insect as host of Myxosporidia, 25
 Integument, Myxosporidia parasitic in, 37
 Intercapsular appendix of spore, 47
 Intestine, Myxosporidia parasitic in, 41
 Iodinophilous vacuole, 47, 50
 Italian Myxosporidia, 17
 Kamtschatka, Myxosporidian of, 14
 Kidney, Myxosporidia parasitic in, 42
 Lateral process of spore, 50
 Length of spore, 50
 Lentospora, 11, 58, 125, 179, 189
 Lentospora acuta, 127
 asymmetrica, 126
 cerebralis, 125
 dermatobia, 127
 encephalina, 126
 multiplicata, 126
 Leptotheca, 9, 56, 60, 178, 179
 Leptotheca agilis, 60
 elongata, 60
 fusiformis, 63
 glomerosa, 65
 hepseti, 62
 informis, 63
 lobosa, 64
 longipes, 63
 macrospora, 62
 parva, 61
 perlata, 62
 polymorpha, 61
 renicola, 61
 scissura, 64
 sp. Awerinzew, 62
 Liver, Myxosporidia parasitic in, 39
 Longitudinal striation on spore, 50
 Mesentery, Myxosporidia parasitic in, 43
 Mesoplasm, 50
 Mictosporaea, 52, 54
 Mictosporous, 50
 Mitraspora, 9, 56, 84, 178, 183
 Mitraspora caudata, 85
 cyprini, 84
 elongata, 85
 Monaco, Myxosporidia of, 17
 Monosporous, 50
 Muscle, Myxosporidia parasitic in, 38
 Myxidiidae, 10, 57, 106, 179
 Myxidium, 10, 58, 107, 179, 186
 Myxidium americanum, 117
 anguillae, 114
 barbatulae, 110
 bergense, 112
 danilewskyi, 109
 depressum, 114

- gadi*, 115
giardi, 110
giganteum, 110
glutinosum, 115
histophilum, 109
incurvatum, 108
inflatum, 111
kagayamai, 117
lieberkühni, 107
mackiei, 112
macrocapsulare, 113
oviforme, 114
pfeifferi, 111
phyllium, 116
 sp. Awerinzew, 113
 sp. Gurley, 109
 sp. Mavor, 115
sphaericum, 109
striatum, 116
- Myxobolidae, 11, 58, 128, 179
 Myxobolus, 11, 58, 128, 179, 189
Myxobolus aeglefini, 144
anurus, 142
aureatus, 154
balleri, 147
bramae, 147
capsulatus, 152
carassii, 150
cordis, 148
cycloides, 140
cyprini, 143
cyprinicola, 147
discrepans, 156
dispar, 135
ellipsoides, 136
exiguus, 136
fuhmanni, 130
funduli, 151
gigas, 145
globosus, 139
hylae, 153
inaequalis, 135
koi, 155
linloni, 138
macrocapsularis, 146
magnus, 150
mesentericus, 157
minutus, 150
misgurni, 133
miyairii, 155
mülleri, 128
musculi, 148
neurobins, 144
nodularis, 153
notatus, 131
obesus, 140
oblongus, 139
oculi-leucisci, 130
orbiculatus, 155
oviformis, 137
permagnus, 149
pfeifferi, 133
physophilus, 146
piriformis, 129
pleuronectidae, 152
rohita, 132
rotundus, 149
sandrae, 146
scardinii, 146
seni, 132
 sp. Gurley, 142
 sp. Gurley, 142
 sp. Gurley, 143
 sp. Kudo, 132
 sp. Lebzelter, 150
 sp. Miyairi, 149
 sp. Southwell, 151
 sp. Wegener, 149
sphaeralis, 141
squamae, 147
toyamai, 131
transovalis, 139
unicapsulatus, 129
volgensis, 145
- Myxoproteus, 9, 56, 81, 179, 183
Myxoproteus ambiguus, 81
cordiformis, 81
cornutus, 82
- Myxosoma, 11, 58, 123, 179, 189
Myxosoma dujardini, 124
funduli, 125
 ? *lobatum*, 124
- Myxosomatidae, 11, 58, 123, 179
 Myxosporidia in air bladder, 39
 alimentary canal, 39, 41
 Amphibia, 36
 Annelida, 25
 blood vessel, 39
 body cavity, 37, 43, 45
 bone, 39
 branchia, 38
 cartilage, 39
 connective tissue, 37
 eye, 38

- gall bladder, 39
 heart, 39
 Insecta, 25
 integument, 37
 intestine, 41
 kidney, 41
 liver, 39
 mesentery, 41
 muscle, 38
 nervous tissue, 39
 organs of host, 37
 ovary, 41
 Pisces, 25
 pyloric coecum, 39
 Reptilia, 36
 spleen, 41
 stomach, 39
 testis, 41
 tissue, 37, 43, 45
 unknown seat, 41
 urinary bladder, 41
 of Africa, 14
 Asia, 13
 Burma, 14
 India, 14
 Japan, 13
 Kamtschatka, 14
 Nippon, 13
 Australia, 14
 Europe, 17
 Austria, 24
 England, 22
 France, 18
 Germany, 20
 Italy, 17
 Monaco, 17
 Netherland, 22
 Norway, 22
 Russia, 24
 Serbia, 24
 Switzerland, 23
 North America, 15
 Alaska, 16
 Canada, 16
 United States, 15
 South America, 16
 unknown genera and species,
 12, 174
 Nervous tissue, Myxosporidia parasitic in, 39
 Netherland, Myxosporidian of, 22
 New species listed, 196
 Nippon, Myxosporidia of, 13
 North American Myxosporidia, 15
 Norwegian Myxosporidia, 22
 Organs of host infected by Myxosporidia, list
 of, 37
 Ovary, Myxosporidia parasitic in, 41
 Pansporoblast, 50
 Plasmogamy, 51
 Plasmotomy, 51
 Platysporea, 10, 57, 106, 179
 Polar capsule, 48, 51
 filament, 48, 51
 Polysporea, 52, 54
 Polysporous, 51
 Posterior filament, 47, 51
 process, 47, 51
 Pyloric coecum, Myxosporidia parasitic in, 39
 Reptilian hosts of Myxosporidia, 36
 Ridge of spore, 51
 Russian Myxosporidia, 24
 Serbian Myxosporidian, 24
 Shell of spore, 47, 51
 -valve, 47, 51
 Sinuolinea, 10, 57, 104, 178, 186
Sinuolinea arborescens, 105
 brachiophora, 106
 capsularis, 105
 dimorpha, 104
 opacita, 106
 South American Myxosporidia, 16
 Species, scheme of description of, 7
 Sphaeromyxa, 10, 58, 118, 179, 188
Sphaeromyxa balbianii, 118
 exneri, 121
 gasterostei, 121
 hellandi, 121
 immersa, 119
 incurvata, 119
 sabrazesi, 120
 Sphaerospora, 10, 57, 100, 178, 185
Sphaerospora angulata, 102
 carassii, 103
 divergens, 100
 elegans, 100
 masovica, 101
 platessae, 102
 polymorpha, 102
 rostrata, 101
 sp. Davis, 103
 sp. Southwell et Prashad, 103
 Sphaerosporea, 10, 57, 86, 178

- Sphaerosporidae, 10, 57, 99, 178
 Spleen, Myxosporidia found in, 41
 Spore, description of, 47
 Sporoplasm, 51
 Stomach, Myxosporidia parasitic in, 39
 Sutural diameter, 51
 edge, 51
 line, 51
 plane, 51
 ridge, 51
 Switzerland, Myxosporidia of, 23
 Tail of spore, 51
 Testis, Myxosporidia parasitic in, 41
 Thickness of spore, 51
 Tissue, Myxosporidia parasitic in, 37, 43, 45
 Trophozoite, 51
 United States, Myxosporidia of, 15
 Urinary bladder, Myxosporidia parasitic in,
 41
 Vegetative form, 51
 Wardia, 9, 56, 82
Wardia ohlmacheri, 83
 ovinocua, 82
 Zschokkella, 10, 58, 122
Zschokkella acheilognathi, 123
 globulosa, 123
 hildae, 122
 nova, 122

ILLINOIS BIOLOGICAL
MONOGRAPHS

Vol. V

July-October, 1919

Nos. 3 and 4

STUDIES ON MYXOSPORIDIA

A SYNOPSIS OF GENERA AND SPECIES
OF MYXOSPORIDIA

WITH 25 PLATES AND 2 TEXTFIGURES

BY

ROKUSABURO KUDO

Price \$3.00

PUBLISHED BY THE UNIVERSITY OF ILLINOIS
UNDER THE AUSPICES OF THE GRADUATE SCHOOL
URBANA, ILLINOIS

UNIVERSITY OF ILLINOIS STUDIES PUBLISHED
ILLINOIS BIOLOGICAL MONOGRAPHS

THE ILLINOIS BIOLOGICAL MONOGRAPHS is a serial published quarterly by the University of Illinois. It is the purpose of the publication to present in monographic form research contributions of especial merit and of somewhat extended character. Each volume will include about 500 pages and 20 or more full page plates. The series may be obtained at a subscription price of three dollars yearly. Individual numbers will be sold separately at prices determined by the size and amount of illustration contained in the particular number.

Vol. I

- Nos. 1 and 2. A Revision of the Cestode family Proteocephalidae. With 16 plates. By G. R. La Rue. \$2.00.
No. 3. Studies on the Cestode family Anoplocephalidae. With 6 plates. By H. Douthitt. 80 cts.
No. 4. Some North American Larval Trematodes. With 8 plates. By W. W. Cort. \$1.20.

Vol. II

- No. 1. Classification of Lepidopterous larvae. With 10 plates. By S. B. Fracker. \$1.50.
No. 2. On the Osteology of some of the Loricati. With 5 plates. By J. E. Gutberlet. 50 cts.
No. 3. Studies on Gregarines, including twenty-one new species and a synopsis of the Eugregarine records from the Myriapoda, Coleoptera, and Orthoptera of the world. With 15 plates. By Minnie E. Watson. \$2.00.
No. 4. The Genus *Meliola* in Porto Rico. With 5 plates. By Frank L. Stevens. 75 cts.

Vol. III

- No. 1. Studies on the factors controlling the rate of regeneration. By Charles Zeleny. \$1.25.
No. 2. The head-capsule and mouth-parts of Diptera. With 25 plates. By Alvah Peterson. \$2.00.
No. 3. Studies on North American Polystomidae, Aspidogastridae, and Paramphistomidae. With 11 plates. By Horace W. Stunkard. \$1.25.
No. 4. Color and color-pattern mechanism of Tiger Beetles. With 29 black and 3 colored plates. By Victor E. Shelford. \$2.00.

Vol. IV

- No. 1. Life history studies on Montana Trematodes. By E. C. Faust. With 9 plates. \$2.00.
No. 2. The Goldfish as a test animal in the study of toxicity. By E. B. Powers. \$1.00.
No. 3. Morphology and biology of some Turbellaria from the Mississippi basin. With 3 plates. By Ruth Higley. \$1.25.
No. 4. North American pseudophyllidean Cestodes from fishes. With 13 plates. By A. R. Cooper. \$2.00.

Vol. V

- No. 1. The Skull of *Amiurus*. With 8 plates. By J. E. Kindred. \$1.25.
No. 2. Contributions to the Life Histories of *Gordius robustus* Leidy and *Paragordius varius* (Leidy). By Henry Gustav May. With 21 plates. \$1.50.
Nos. 3 and 4. Studies on Myxosporidia. A Synopsis of Genera and Species of Myxosporidia. By Rokusaburo Kudo. With 25 plates and 2 textfigures. \$3.00.

Vol. VI

- No. 1. The Nasal Organ in Amphibia. By G. M. Higgins. With 10 plates. *In press.*
Nos. 2 and 3. Revision of the North American and West Indian Species of *Cuscuta*. With 13 plates. By Truman George Yuncker. *In press.*
No. 4. The Larvae of the Coccinellidae. With 6 plates. By J. Howard Gage. *In press.*

Entered as second-class matter July 27, 1915, at the post-office at Urbana, Illinois, under the Act of August 24, 1912. Acceptance for mailing at the special rate of postage provided for in section 1102, Act of October 3, 1917, authorized July 31, 1918

UNIVERSITY OF ILLINOIS STUDIES IN LANGUAGE AND LITERATURE

The Studies in Language and Literature are designed to include monographs in general linguistics and comparative literature; the classical languages and Sanskrit; the Romance languages; and English, the Scandinavian, and other Germanic languages. The title of the series will be so construed as to admit the publication of such researches in the history of culture as may throw light on the processes of language and the interpretation of literature. This series is published quarterly; the annual subscription price is three dollars.

Vol. III

- No. 1. The *Ad Deum Vadit* of Jean Gerson. By David Hobart Carnahan. \$1.75.
- No. 2. Tagalog Texts: Part I. Texts and translation. By Leonard Bloomfield. \$1.50.
- No. 3. The same. Part II. Grammatical analysis. By Leonard Bloomfield. \$3.00.
- No. 4. The same. Part III. List of formations and glossary. By Leonard Bloomfield. \$1.50.

Vol. IV

- No. 1. Madame de Staël's literary reputation in England. By Robert C. Whitford. 75 cts.
- No. 2. Index verborum quae in Senecae fabulis necnon in Octavia Praetexta reperiuntur a Guilielmo Abbott Oldfather, Arthuro Stanley Pease, Howardo Vernon Canter confectus. \$1.50.
- No. 3. The same. Part II. \$1.50.
- No. 4. The same. Part III. \$1.50.

VOL. V

- No. 1. The influence of Christianity on the Vocabulary of Old English poetry. Part I. By Albert Keiser. 75 cents.
- No. 2. The same. Part II. 75 cents.
- No. 3. Spencer's Defense of Lord Gray. By H. S. V. Jones. \$1.00.
- No. 4. Ysoper and Avionet. Latin and French texts. Kenneth McKenzie and W. A. Oldfather. (*In press.*)

Vol. VI

- No. 1. La Colección Cervantina de la Sociedad Hispánica de América (The Hispanic Society of America): Ediciones de Don Quijote. By Homero Seris. \$1.50. (*In press.*)

UNIVERSITY OF ILLINOIS STUDIES IN THE SOCIAL SCIENCES

The University of Illinois Studies in the Social Sciences are designed to afford a means of publishing monographs prepared by graduate students or members of the faculty in the departments of history, economics, political science, and sociology. Each volume will consist of about 600 printed pages annually. The subscription price is three dollars a year.

Vol. II

- No. 1. Taxation of corporations in Illinois, other than railroads, since 1872. By J. R. Moore. 55 cts.
- Nos. 2 and 3. The West in the diplomatic negotiations of the American Revolution. By P. C. Phillips. \$1.25.
- No. 4. The development of banking in Illinois, 1817-1863. By G. W. Dowrie. 90 cts.

Vol. III

- Nos. 1 and 2. The history of the general property tax in Illinois. By R. M. Halg. \$1.25.

Vol. IV

- Nos. 1. The Illinois Whigs before 1846. By C. M. Thompson. 95 cts.
- Nos. 3-4. The history of the Illinois Central railway to 1870. By H. G. Brownson. \$1.25.

UNIVERSITY OF ILLINOIS STUDIES IN THE SOCIAL SCIENCES.—Continued.

Vol. V

- No. 1. Enforcement of international law by municipal law. By P. Q. Wright. \$1.25.
No. 2. The life of Jesse W. Fell. By Frances M. Morehouse. 60 cts.
No. 4. Mine taxation in the United States. By L. E. Young. \$1.50.

Vol. VI

- Nos. 1 and 2. The veto power of the Governor of Illinois. By N. H. Debel. \$1.00.
No. 3. Wage bargaining on the vessels of the Great Lakes. By H. E. Hoagland. \$1.50.
No. 4. The household of a Tudor nobleman. By P. V. B. Jones. \$1.50.

Vol. VII

- No. 4. The Journeymen Tailors' Union of America. A study in trade union policy. By Charles J. Stowell. \$1.00.

Vol. VIII

- No. 2. Cumulative voting and minority representation in Illinois. Revised edition. By B. F. Moore. 75 cents.
Nos. 3 and 4. Labor problems and labor administration in the United States during the World War. By Gordon Watkins. \$2.00.

UNIVERSITY STUDIES

GENERAL SERIES, Vol. I, II, III AND IV

Partly out of print. A detailed list of these numbers will be sent on request.

PUBLICATIONS OF THE UNIVERSITY OF ILLINOIS

Following is a partial list of the publications issued at the University:

1. THE UNIVERSITY OF ILLINOIS STUDIES IN LANGUAGE AND LITERATURE. Published quarterly. Three dollars a year.
2. THE UNIVERSITY OF ILLINOIS STUDIES IN THE SOCIAL SCIENCES. Monographs in history, economics, political science, and sociology. Quarterly. Three dollars a year.
3. THE ILLINOIS BIOLOGICAL MONOGRAPHS. Quarterly. Three dollars a year.
4. THE UNIVERSITY STUDIES. A series of monographs on miscellaneous subjects.
5. THE JOURNAL OF ENGLISH AND GERMANIC PHILOLOGY. Quarterly. Three dollars a year.

For any of the above address 161 Administration Building, Urbana, Illinois.

6. THE BULLETIN OF THE ENGINEERING EXPERIMENT STATION. Reports of the research work in the Engineering Experiment Station. Address Director of Engineering Experiment Station, University of Illinois.
7. THE BULLETIN OF THE AGRICULTURAL EXPERIMENT STATION. Address Director of Agricultural Experiment Station, University of Illinois.
8. THE BULLETIN OF THE STATE LABORATORY OF NATURAL HISTORY. Address Director of State Laboratory of Natural History, University of Illinois.
9. THE BULLETIN OF THE STATE GEOLOGICAL SURVEY. Address Director of State Geological Survey, University of Illinois.
10. THE BULLETIN OF THE STATE WATER SURVEY. Address Director of State Water Survey, University of Illinois.
11. THE REPORT OF THE STATE ENTOMOLOGIST. Address State Entomologist, University of Illinois.
12. THE BULLETIN OF THE ILLINOIS ASSOCIATION OF TEACHERS OF ENGLISH. Address 301 University Hall, University of Illinois.
13. THE BULLETIN OF THE SCHOOL OF EDUCATION. Address 203 University Hall, University of Illinois.
14. THE LAW BULLETIN. Address 161 Administration Building, University of Illinois.
15. The general series, containing the University catalog and circulars of special departments. Address The Registrar, University of Illinois.

UNIVERSITY OF ILLINOIS-URBANA



3 0112 065097237